



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:48 AM UTC

PDB ID : 3DA6 / pdb_00003da6
Title : Crystal Structure of human JNK3 complexed with N-(3-methyl-4-(3-(2-(methylamino)pyrimidin-4-yl)pyridin-2-yloxy)naphthalen-1-yl)-1H-benzo[d]imidazole-2-amine
Authors : Cee, V.J.; Cheng, A.C.; Romero, K.; Bellon, S.; Mohr, C.; Whittington, D.A.; Bready, J.; Caenepeel, S.; Coxon, A.; Deak, H.L.; Hodous, B.L.; Kim, J.L.; Lin, J.; Nguyen, H.; Olivieri, P.R.; Patel, V.F.; Wang, L.; Hughes, P.; Geuns-Meyer, S.
Deposited on : 2008-05-28
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.49

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BZ9	A	1	-	-	X	-

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.48Å 71.28Å 106.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.72 – 2.00 22.72 – 2.00	Depositor EDS
% Data completeness (in resolution range)	91.5 (22.72-2.00) 91.4 (22.72-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.256 , 0.320 0.253 , 0.314	Depositor DCC
R_{free} test set	1279 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2757	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BZ9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.61	28/2723 (1.0%)	1.48	26/3679 (0.7%)

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	229	TYR	N-CA	8.69	1.56	1.46
1	A	178	LYS	N-CA	7.84	1.55	1.46
1	A	348	ILE	CA-CB	7.79	1.61	1.54
1	A	204	LYS	C-O	7.28	1.32	1.23
1	A	140	GLN	CA-C	7.01	1.59	1.52
1	A	211	ALA	N-CA	6.79	1.53	1.46
1	A	203	LEU	C-O	6.70	1.32	1.23
1	A	123	ILE	C-O	-6.60	1.17	1.24
1	A	196	VAL	C-O	6.57	1.31	1.23
1	A	327	LEU	C-O	-6.24	1.17	1.24
1	A	59	THR	CA-CB	6.05	1.62	1.53
1	A	82	TYR	C-O	-6.01	1.16	1.24
1	A	205	ILE	C-O	-5.96	1.17	1.24
1	A	255	GLU	CA-C	5.93	1.60	1.52
1	A	269	ILE	C-O	-5.80	1.17	1.24
1	A	257	VAL	CA-C	5.75	1.59	1.52
1	A	241	LYS	CA-C	5.54	1.58	1.53
1	A	177	ILE	CA-C	5.51	1.59	1.52
1	A	247	TRP	CB-CG	5.47	1.67	1.50
1	A	259	HIS	N-CA	5.35	1.52	1.46
1	A	300	ASN	N-CA	5.33	1.52	1.46
1	A	124	ILE	C-O	5.32	1.30	1.24
1	A	178	LYS	CB-CG	5.31	1.68	1.52
1	A	244	VAL	CA-C	5.31	1.59	1.52
1	A	93	LYS	N-CA	5.30	1.52	1.46
1	A	394	ILE	CA-CB	5.26	1.60	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	THR	CA-CB	5.17	1.60	1.53
1	A	350	VAL	C-O	5.05	1.29	1.24

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	MET	N-CA-C	10.94	126.46	112.89
1	A	283	CYS	CA-C-N	10.22	132.61	119.84
1	A	283	CYS	C-N-CA	10.22	132.61	119.84
1	A	137	GLU	N-CA-C	-6.80	103.66	112.23
1	A	394	ILE	N-CA-C	-6.56	104.36	110.53
1	A	103	THR	N-CA-C	6.45	118.11	111.14
1	A	118	VAL	N-CA-C	6.14	122.10	109.34
1	A	281	THR	N-CA-C	-5.99	102.72	109.60
1	A	291	GLN	CA-C-N	5.91	126.33	119.47
1	A	291	GLN	C-N-CA	5.91	126.33	119.47
1	A	339	MET	N-CA-C	-5.72	105.97	113.12
1	A	301	ARG	CA-C-N	5.68	126.57	119.98
1	A	301	ARG	C-N-CA	5.68	126.57	119.98
1	A	312	LEU	N-CA-C	5.56	118.06	111.33
1	A	63	ARG	NE-CZ-NH2	5.56	124.20	119.20
1	A	303	LYS	N-CA-C	5.47	118.58	110.48
1	A	156	VAL	CA-C-N	5.39	128.13	120.53
1	A	156	VAL	C-N-CA	5.39	128.13	120.53
1	A	140	GLN	N-CA-C	5.28	119.95	112.93
1	A	64	TYR	N-CA-C	5.20	116.78	108.41
1	A	97	ARG	CA-C-O	5.18	124.40	120.26
1	A	278	GLN	N-CA-C	-5.16	107.05	113.55
1	A	289	LYS	N-CA-C	-5.13	106.42	113.30
1	A	175	CYS	CA-C-N	-5.12	114.36	120.00
1	A	175	CYS	C-N-CA	-5.12	114.36	120.00
1	A	179	HIS	N-CA-C	5.00	117.11	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2666	0	2698	78	0
2	A	36	0	23	10	0
3	A	55	0	0	4	0
All	All	2757	0	2721	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:CYS:HB2	1:A:284:PRO:HD2	1.43	1.00
1:A:283:CYS:CB	1:A:284:PRO:HD2	1.98	0.88
1:A:226:THR:HG21	3:A:449:HOH:O	1.75	0.86
1:A:162:ASP:OD2	1:A:164:GLU:HG3	1.78	0.84
1:A:283:CYS:HB2	1:A:284:PRO:CD	2.09	0.81
1:A:71:GLY:HA3	2:A:1:BZ9:H23	1.61	0.80
1:A:102:GLN:OE1	1:A:385:HIS:N	2.15	0.79
1:A:93:LYS:NZ	2:A:1:BZ9:HN1	1.82	0.76
1:A:115:MET:HE2	1:A:124:ILE:HD13	1.69	0.74
1:A:115:MET:HE2	1:A:124:ILE:HG21	1.71	0.72
1:A:120:HIS:HD2	1:A:122:ASN:H	1.36	0.72
1:A:77:ILE:HD13	1:A:77:ILE:N	2.05	0.71
1:A:120:HIS:CD2	1:A:122:ASN:H	2.09	0.71
1:A:399:MET:HE3	3:A:441:HOH:O	1.92	0.69
1:A:297:TYR:CE1	1:A:301:ARG:NH1	2.61	0.68
1:A:291:GLN:HA	1:A:291:GLN:HE21	1.56	0.67
1:A:131:THR:CG2	1:A:399:MET:CE	2.73	0.66
1:A:399:MET:O	1:A:400:ASN:HB2	1.94	0.66
1:A:155:GLN:HA	1:A:158:GLN:HE21	1.61	0.66
1:A:297:TYR:HE1	1:A:301:ARG:NH1	1.95	0.64
1:A:276:ILE:O	1:A:280:GLY:N	2.31	0.63
1:A:131:THR:CG2	1:A:399:MET:HE2	2.29	0.63
1:A:283:CYS:CB	1:A:284:PRO:CD	2.73	0.62
1:A:61:LEU:HB3	3:A:428:HOH:O	1.99	0.61
1:A:77:ILE:HD13	1:A:77:ILE:H	1.65	0.61
1:A:116:LYS:HG2	1:A:117:CYS:N	2.17	0.60
1:A:106:LYS:HE2	1:A:213:THR:HG21	1.84	0.59
1:A:97:ARG:NH1	1:A:140:GLN:OE1	2.31	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:LYS:HE2	2:A:1:BZ9:H8	1.85	0.58
1:A:78:VAL:HG22	1:A:93:LYS:HB2	1.85	0.58
1:A:114:LEU:HD22	1:A:185:ILE:HD11	1.86	0.56
1:A:110:ARG:NH1	1:A:114:LEU:HD11	2.21	0.56
1:A:131:THR:HG22	1:A:399:MET:HE2	1.88	0.55
1:A:283:CYS:O	1:A:286:PHE:HB3	2.07	0.55
1:A:62:LYS:N	3:A:428:HOH:O	2.41	0.54
1:A:357:PRO:HA	1:A:360:ASN:ND2	2.22	0.54
1:A:95:LEU:CD1	2:A:1:BZ9:H7	2.38	0.53
1:A:71:GLY:HA3	2:A:1:BZ9:C23	2.37	0.53
1:A:115:MET:HE2	1:A:124:ILE:CG2	2.37	0.52
1:A:276:ILE:O	1:A:280:GLY:CA	2.58	0.51
1:A:267:ASP:OD2	1:A:270:ASP:HB2	2.10	0.51
1:A:135:THR:OG1	1:A:138:GLU:HG3	2.10	0.51
1:A:321:ASP:OD1	1:A:321:ASP:N	2.40	0.51
1:A:394:ILE:O	1:A:398:VAL:HG23	2.10	0.51
1:A:93:LYS:HZ3	2:A:1:BZ9:HN1	1.57	0.50
1:A:188:ARG:HG2	1:A:244:VAL:HG11	1.94	0.50
1:A:95:LEU:HD13	2:A:1:BZ9:H7	1.94	0.49
1:A:99:PHE:O	1:A:387:ILE:HG23	2.14	0.48
1:A:93:LYS:HE2	2:A:1:BZ9:C8	2.44	0.48
1:A:120:HIS:HB2	1:A:179:HIS:CD2	2.49	0.48
1:A:287:MET:O	1:A:290:LEU:HB2	2.14	0.48
1:A:46:ASN:HD22	1:A:47:GLN:N	2.12	0.47
1:A:270:ASP:OD1	1:A:274:LYS:NZ	2.46	0.47
1:A:254:GLY:HA3	1:A:262:LEU:HD11	1.96	0.47
1:A:250:GLY:HA2	1:A:339:MET:HE3	1.95	0.47
1:A:131:THR:HG22	1:A:399:MET:CE	2.43	0.47
1:A:152:ASN:OD1	1:A:152:ASN:C	2.58	0.46
1:A:115:MET:HE3	1:A:208:PHE:CE1	2.51	0.46
1:A:83:ASP:O	1:A:87:ASP:N	2.49	0.46
1:A:49:TYR:O	1:A:59:THR:HA	2.16	0.45
1:A:65:GLN:OE1	1:A:84:ALA:HB2	2.16	0.45
1:A:95:LEU:HD11	2:A:1:BZ9:C7	2.47	0.45
1:A:63:ARG:HB2	1:A:85:VAL:HG23	1.99	0.45
1:A:253:MET:O	1:A:257:VAL:HG23	2.18	0.44
1:A:281:THR:HG21	1:A:303:LYS:HE3	1.99	0.44
1:A:307:LEU:HB2	1:A:312:LEU:HG	1.99	0.44
1:A:246:ILE:HD13	1:A:246:ILE:HA	1.90	0.44
1:A:287:MET:HE1	1:A:298:VAL:HG12	1.99	0.43
1:A:255:GLU:O	1:A:259:HIS:HA	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:HIS:HA	1:A:357:PRO:HD3	1.86	0.42
1:A:264:PRO:HD2	1:A:274:LYS:HD2	2.02	0.42
1:A:233:GLU:HB2	1:A:238:MET:HB2	2.02	0.41
1:A:315:ASP:OD1	1:A:315:ASP:N	2.50	0.41
1:A:77:ILE:H	1:A:77:ILE:CD1	2.25	0.41
1:A:93:LYS:CE	2:A:1:BZ9:HN1	2.34	0.41
1:A:392:GLU:OE2	1:A:396:LYS:HE2	2.21	0.41
1:A:102:GLN:CD	1:A:390:TRP:HE1	2.30	0.40
1:A:356:HIS:CE1	1:A:357:PRO:HD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	320/364 (88%)	302 (94%)	15 (5%)	3 (1%)	14 9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	CYS
1	A	118	VAL
1	A	284	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/329 (90%)	271 (91%)	26 (9%)	9 6

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	VAL
1	A	56	SER
1	A	68	LYS
1	A	72	SER
1	A	77	ILE
1	A	95	LEU
1	A	96	SER
1	A	102	GLN
1	A	115	MET
1	A	116	LYS
1	A	118	VAL
1	A	134	LYS
1	A	164	GLU
1	A	166	MET
1	A	224	VAL
1	A	225	VAL
1	A	226	THR
1	A	274	LYS
1	A	288	LYS
1	A	291	GLN
1	A	293	THR
1	A	296	ASN
1	A	301	ARG
1	A	307	LEU
1	A	315	ASP
1	A	384	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	47	GLN
1	A	66	ASN
1	A	89	ASN
1	A	120	HIS
1	A	158	GLN
1	A	271	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	291	GLN
1	A	385	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BZ9	A	1	-	40,41,41	1.64	8 (20%)	54,58,58	3.30	15 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BZ9	A	1	-	-	4/14/14/14	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	BZ9	C21-C28	-4.09	1.38	1.48
2	A	1	BZ9	C3-N1	-3.31	1.32	1.38
2	A	1	BZ9	C4-N2	-2.74	1.34	1.39
2	A	1	BZ9	C9-N16	-2.66	1.31	1.36
2	A	1	BZ9	C20-N16	-2.61	1.33	1.41
2	A	1	BZ9	C21-C26	-2.51	1.38	1.41
2	A	1	BZ9	C14-C15	2.21	1.41	1.36
2	A	1	BZ9	C7-C6	2.17	1.43	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	BZ9	C17-O33-C26	12.52	129.87	116.84
2	A	1	BZ9	C28-N27-C32	11.46	125.63	116.81
2	A	1	BZ9	N31-C32-N27	-9.52	117.21	126.42
2	A	1	BZ9	N34-C32-N27	6.10	122.82	116.96
2	A	1	BZ9	C30-N31-C32	5.55	120.05	115.42
2	A	1	BZ9	C30-C29-C28	-4.66	112.69	117.20
2	A	1	BZ9	C11-C20-N16	4.48	125.83	118.35
2	A	1	BZ9	C19-C20-N16	-3.50	113.25	121.90
2	A	1	BZ9	C29-C28-N27	-3.10	117.88	121.97
2	A	1	BZ9	N34-C32-N31	2.90	119.81	117.00
2	A	1	BZ9	C13-C14-C15	-2.61	116.90	120.40
2	A	1	BZ9	C7-C6-C5	-2.43	117.24	120.24
2	A	1	BZ9	C6-C7-C8	2.40	123.20	120.24
2	A	1	BZ9	C4-C3-N1	2.14	108.28	105.71
2	A	1	BZ9	C15-C10-C11	2.02	120.92	118.42

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	BZ9	N1-C9-N16-C20
2	A	1	BZ9	N2-C9-N16-C20
2	A	1	BZ9	C11-C20-N16-C9
2	A	1	BZ9	C19-C20-N16-C9

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	BZ9	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/364 (90%)	0.67	31 (9%) 14 12	21, 37, 60, 71	4 (1%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	GLY	6.0
1	A	117	CYS	4.8
1	A	115	MET	4.5
1	A	284	PRO	3.8
1	A	365	PRO	3.7
1	A	71	GLY	3.3
1	A	306	GLY	3.2
1	A	213	THR	3.0
1	A	307	LEU	2.8
1	A	224	VAL	2.8
1	A	302	PRO	2.8
1	A	319	PRO	2.8
1	A	65	GLN	2.7
1	A	299	GLU	2.7
1	A	366	ALA	2.7
1	A	300	ASN	2.5
1	A	304	TYR	2.5
1	A	363	TYR	2.5
1	A	107	ARG	2.5
1	A	201	CYS	2.4
1	A	55	ASP	2.4
1	A	320	ALA	2.4
1	A	303	LYS	2.4
1	A	330	SER	2.3
1	A	305	ALA	2.3
1	A	212	ARG	2.1
1	A	52	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	383	ARG	2.1
1	A	76	GLY	2.1
1	A	56	SER	2.1
1	A	57	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BZ9	A	1	36/36	0.92	0.08	23,30,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.