



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 12:57 AM UTC

PDB ID : 4DA5 / pdb_00004da5
Title : Choline Kinase alpha acts through a double-displacement kinetic mechanism involving enzyme isomerisation, as determined through enzyme and inhibitor kinetics and structural biology
Authors : Brown, K.; Hudson, C.; Charlton, P.; Pollard, J.
Deposited on : 2012-01-12
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	
1	B	457	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5951 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Choline kinase alpha.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2926	1893	492	525	16			
1	B	330	Total	C	N	O	S	0	0	0
			2742	1784	453	490	15			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is (3R)-1-azabicyclo[2.2.2]oct-3-yl[bis(5-chlorothiophen-2-yl)]methanol (CCD ID: 0H7) (formula: C₁₆H₁₇Cl₂NOS₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			22	16	2	1	1	2		
3	B	1	Total	C	Cl	N	O	S	0	0
			22	16	2	1	1	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total	O	0	0
			123	123		
4	B	106	Total	O	0	0
			106	106		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.84Å 121.69Å 131.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.50 – 2.40 18.50 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.6 (18.50-2.40) 98.4 (18.50-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.01 (at 2.40Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.196 , 0.259 0.205 , 0.264	Depositor DCC
R_{free} test set	1758 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5951	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, OH7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/3002	1.32	9/4042 (0.2%)
1	B	0.77	0/2813	1.33	6/3787 (0.2%)
All	All	0.77	0/5815	1.32	15/7829 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	372	LYS	CA-C-N	6.12	128.39	120.44
1	A	372	LYS	C-N-CA	6.12	128.39	120.44
1	A	128	SER	N-CA-C	5.62	117.81	108.99
1	A	430	ILE	N-CA-C	5.36	119.91	112.35
1	B	432	SER	CA-C-N	5.30	127.60	120.50
1	B	432	SER	C-N-CA	5.30	127.60	120.50
1	A	293	LEU	CA-C-N	5.17	127.21	120.28
1	A	293	LEU	C-N-CA	5.17	127.21	120.28
1	B	452	LYS	CA-C-N	5.15	127.45	120.65
1	B	452	LYS	C-N-CA	5.15	127.45	120.65
1	A	249	LEU	N-CA-C	5.10	117.24	111.02
1	B	417	HIS	CA-C-N	5.03	127.28	120.44
1	B	417	HIS	C-N-CA	5.03	127.28	120.44
1	A	116	ILE	N-CA-C	-5.02	107.25	111.56
1	A	437	TYR	N-CA-C	5.02	116.75	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2926	0	2897	11	0
1	B	2742	0	2718	4	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	22	0	17	0	0
3	B	22	0	17	1	0
4	A	123	0	0	1	0
4	B	106	0	0	0	0
All	All	5951	0	5649	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:ILE:HD11	1:A:126:GLN:HB2	1.85	0.57
1:B:254:GLU:OE2	1:B:290:ARG:NH2	2.30	0.54
1:A:387:GLN:HG3	1:A:390:PHE:HB2	1.92	0.52
1:A:254:GLU:OE2	1:A:290:ARG:NH2	2.43	0.51
1:A:249:LEU:HG	1:A:253:MET:HE2	1.93	0.50
1:A:128:SER:HB3	1:A:142:LYS:HB3	1.95	0.49
1:B:390:PHE:HZ	1:B:402:LYS:HE3	1.79	0.47
1:A:352:TYR:O	4:A:721:HOH:O	2.20	0.47
1:A:268:GLU:HG3	1:A:272:LYS:HE2	1.97	0.46
3:B:502:OH7:H8	3:B:502:OH7:O1	2.16	0.46
1:A:192:LEU:HD12	1:A:238:MET:SD	2.59	0.43
1:A:253:MET:HG3	1:A:419:LEU:HD12	2.01	0.42
1:A:122:ASN:HA	1:A:147:LEU:O	2.20	0.41
1:A:201:PRO:HG3	1:B:175:GLU:HG3	2.03	0.41
1:B:195:LYS:HB2	1:B:207:GLN:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/457 (76%)	337 (97%)	11 (3%)	0	100	100
1	B	322/457 (70%)	307 (95%)	15 (5%)	0	100	100
All	All	670/914 (73%)	644 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/400 (79%)	301 (95%)	15 (5%)	23	41
1	B	297/400 (74%)	279 (94%)	18 (6%)	17	30
All	All	613/800 (77%)	580 (95%)	33 (5%)	20	35

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	GLU
1	A	96	LYS
1	A	195	LYS
1	A	199	ILE
1	A	206	GLU
1	A	231	LYS
1	A	238	MET
1	A	258	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	278	LEU
1	A	329	ILE
1	A	332	GLU
1	A	366	ILE
1	A	367	ARG
1	A	399	SER
1	A	429	LYS
1	B	96	LYS
1	B	147	LEU
1	B	175	GLU
1	B	180	GLU
1	B	216	THR
1	B	238	MET
1	B	264	LYS
1	B	272	LYS
1	B	288	ASN
1	B	291	SER
1	B	322	GLU
1	B	329	ILE
1	B	332	GLU
1	B	357	GLU
1	B	368	LYS
1	B	407	LEU
1	B	419	LEU
1	B	427	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	HIS
1	A	324	GLN
1	A	427	GLN
1	B	126	GLN
1	B	202	GLN
1	B	288	ASN
1	B	345	ASN
1	B	427	GLN
1	B	442	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	OH7	B	502	-	23,25,25	1.09	1 (4%)	28,38,38	1.88	8 (28%)
2	SO4	A	501	-	4,4,4	0.36	0	6,6,6	0.18	0
3	OH7	A	502	-	23,25,25	1.24	3 (13%)	28,38,38	1.79	8 (28%)
2	SO4	B	501	-	4,4,4	0.53	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OH7	A	502	-	-	5/18/36/36	0/5/4/4
3	OH7	B	502	-	-	0/18/36/36	0/5/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	OH7	C2-C3	2.89	1.57	1.54
3	B	502	OH7	C20-S22	2.48	1.76	1.71
3	A	502	OH7	C14-S16	2.15	1.76	1.71
3	A	502	OH7	C20-S22	2.10	1.76	1.71

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	OH7	C19-C20-S22	-4.30	109.38	113.30
3	A	502	OH7	C19-C20-S22	-4.28	109.39	113.30
3	B	502	OH7	C11-C2-C17	-3.98	103.84	110.10
3	B	502	OH7	CL2-C20-S22	3.44	125.16	119.72
3	A	502	OH7	CL1-C14-S16	3.30	124.93	119.72
3	A	502	OH7	C13-C14-S16	-3.21	110.37	113.30
3	B	502	OH7	C13-C14-S16	-3.10	110.47	113.30
3	A	502	OH7	CL2-C20-S22	3.09	124.60	119.72
3	B	502	OH7	CL1-C14-S16	2.95	124.38	119.72
3	A	502	OH7	C11-C2-C17	-2.84	105.64	110.10
3	B	502	OH7	C18-C19-C20	2.38	114.75	111.43
3	B	502	OH7	C3-C4-N5	-2.37	108.95	111.36
3	A	502	OH7	C18-C19-C20	2.32	114.66	111.43
3	A	502	OH7	C12-C13-C14	2.18	114.47	111.43
3	B	502	OH7	C12-C13-C14	2.16	114.44	111.43
3	A	502	OH7	C13-C14-CL1	-2.04	124.70	126.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	OH7	C18-C17-C2-O1
3	A	502	OH7	S22-C17-C2-O1
3	A	502	OH7	C18-C17-C2-C3
3	A	502	OH7	S22-C17-C2-C3
3	A	502	OH7	S16-C11-C2-C3


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	OH7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/457 (77%)	0.11	8 (2%) 61 57	22, 40, 70, 84	0
1	B	330/457 (72%)	0.24	17 (5%) 33 29	21, 40, 76, 97	0
All	All	682/914 (74%)	0.17	25 (3%) 45 41	21, 40, 74, 97	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	ALA	3.8
1	A	120	LEU	3.8
1	B	176	ALA	3.6
1	B	149	GLY	3.6
1	B	137	GLY	3.4
1	B	388	ASN	3.4
1	A	121	SER	3.4
1	B	133	THR	3.3
1	A	119	GLY	3.2
1	B	136	LEU	3.1
1	B	391	GLU	3.0
1	B	132	THR	2.8
1	A	388	ASN	2.8
1	B	180	GLU	2.7
1	B	389	ASP	2.6
1	B	115	VAL	2.6
1	B	128	SER	2.6
1	A	118	GLY	2.5
1	B	106	LEU	2.5
1	A	279	SER	2.4
1	B	175	GLU	2.4
1	B	134	ALA	2.2
1	B	399	SER	2.1
1	B	93	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	177	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	0H7	B	502	22/22	0.70	0.20	59,78,89,95	0
3	0H7	A	502	22/22	0.74	0.18	38,50,71,83	0
2	SO4	B	501	5/5	0.86	0.14	43,48,53,53	0
2	SO4	A	501	5/5	0.97	0.09	32,33,41,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around OH7 B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 0H7 A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.