



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:53 PM UTC

PDB ID : 6DA7 / pdb_00006da7
Title : Crystal structure of the TtnD decarboxylase from the tautomycetin biosynthesis pathway of *Streptomyces griseochromogenes* with apo form at 1.83 Å resolution (I222)
Authors : Han, L.; Rudolf, J.D.; Chang, C.-Y.; Miller, M.D.; Soman, J.; Phillips Jr., G.N.; Shen, B.; Enzyme Discovery for Natural Product Biosynthesis (NatPro)
Deposited on : 2018-05-01
Resolution : 1.83 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	9% 91% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UbiD-like decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	463	Total	C	N	O	S	0	10	0
			3639	2306	641	674	18			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	initiating methionine	UNP C6ZCR8
A	-14	GLY	-	expression tag	UNP C6ZCR8
A	-13	SER	-	expression tag	UNP C6ZCR8
A	-12	SER	-	expression tag	UNP C6ZCR8
A	-11	HIS	-	expression tag	UNP C6ZCR8
A	-10	HIS	-	expression tag	UNP C6ZCR8
A	-9	HIS	-	expression tag	UNP C6ZCR8
A	-8	HIS	-	expression tag	UNP C6ZCR8
A	-7	HIS	-	expression tag	UNP C6ZCR8
A	-6	HIS	-	expression tag	UNP C6ZCR8
A	-5	SER	-	expression tag	UNP C6ZCR8
A	-4	GLN	-	expression tag	UNP C6ZCR8
A	-3	ASP	-	expression tag	UNP C6ZCR8
A	-2	PRO	-	expression tag	UNP C6ZCR8
A	-1	ASN	-	expression tag	UNP C6ZCR8
A	0	SER	-	expression tag	UNP C6ZCR8

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O 9 9	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	233	Total O 237 237	0	6

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UbiD-like decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	52.35Å 114.39Å 194.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.77 – 1.83 26.77 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.9 (26.77-1.83) 99.9 (26.77-1.83)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.83Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.189 , 0.213 0.182 , 0.207	Depositor DCC
R_{free} test set	1999 reflections (3.84%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4009	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNL, EPE, GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/3726	0.97	1/5087 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	453	ASN	CA-CB-CG	5.08	117.68	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3639	0	3624	3	0
2	A	1	0	0	0	0
3	A	15	0	17	0	0
4	A	108	0	144	1	0
5	A	9	0	0	0	0
6	A	237	0	0	0	0
All	All	4009	0	3785	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:THR:O	1:A:471:THR:HG22	2.13	0.49
1:A:135:ALA:HB2	4:A:517:GOL:O1	2.15	0.47
1:A:325:THR:HG22	1:A:391:MET:HE1	1.99	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/501 (93%)	459 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/405 (95%)	383 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 1 is monoatomic and 1 is unknown - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	520	-	5,5,5	0.05	0	5,5,5	0.15	0
3	EPE	A	502	2	15,15,15	0.41	0	19,20,20	0.62	0
4	GOL	A	507	-	5,5,5	0.11	0	5,5,5	0.21	0
4	GOL	A	516	-	5,5,5	0.11	0	5,5,5	0.28	0
4	GOL	A	512	-	5,5,5	0.19	0	5,5,5	0.64	0
4	GOL	A	506	-	5,5,5	0.09	0	5,5,5	0.45	0
4	GOL	A	510	-	5,5,5	0.05	0	5,5,5	0.16	0
4	GOL	A	505	-	5,5,5	0.07	0	5,5,5	0.17	0
4	GOL	A	514	-	5,5,5	0.06	0	5,5,5	0.14	0
4	GOL	A	519	-	5,5,5	0.14	0	5,5,5	0.13	0
4	GOL	A	517	-	5,5,5	0.24	0	5,5,5	0.36	0
4	GOL	A	509	-	5,5,5	0.27	0	5,5,5	0.46	0
4	GOL	A	513	-	5,5,5	0.02	0	5,5,5	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	504	-	5,5,5	0.08	0	5,5,5	0.23	0
4	GOL	A	515	-	5,5,5	0.14	0	5,5,5	0.18	0
4	GOL	A	511	-	5,5,5	0.08	0	5,5,5	0.15	0
4	GOL	A	508	-	5,5,5	0.11	0	5,5,5	0.22	0
4	GOL	A	503	-	5,5,5	0.09	0	5,5,5	0.14	0
4	GOL	A	518	-	5,5,5	0.26	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	520	-	-	0/4/4/4	-
3	EPE	A	502	2	-	0/9/19/19	0/1/1/1
4	GOL	A	507	-	-	1/4/4/4	-
4	GOL	A	516	-	-	2/4/4/4	-
4	GOL	A	512	-	-	0/4/4/4	-
4	GOL	A	506	-	-	0/4/4/4	-
4	GOL	A	510	-	-	0/4/4/4	-
4	GOL	A	505	-	-	0/4/4/4	-
4	GOL	A	514	-	-	1/4/4/4	-
4	GOL	A	519	-	-	0/4/4/4	-
4	GOL	A	517	-	-	2/4/4/4	-
4	GOL	A	509	-	-	2/4/4/4	-
4	GOL	A	513	-	-	2/4/4/4	-
4	GOL	A	504	-	-	2/4/4/4	-
4	GOL	A	515	-	-	4/4/4/4	-
4	GOL	A	511	-	-	2/4/4/4	-
4	GOL	A	508	-	-	0/4/4/4	-
4	GOL	A	503	-	-	0/4/4/4	-
4	GOL	A	518	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	504	GOL	C1-C2-C3-O3
4	A	509	GOL	O1-C1-C2-C3
4	A	511	GOL	C1-C2-C3-O3
4	A	513	GOL	O1-C1-C2-C3
4	A	515	GOL	O1-C1-C2-C3
4	A	516	GOL	O1-C1-C2-C3
4	A	518	GOL	C1-C2-C3-O3
4	A	509	GOL	O1-C1-C2-O2
4	A	515	GOL	C1-C2-C3-O3
4	A	517	GOL	C1-C2-C3-O3
4	A	504	GOL	O2-C2-C3-O3
4	A	511	GOL	O2-C2-C3-O3
4	A	513	GOL	O1-C1-C2-O2
4	A	518	GOL	O2-C2-C3-O3
4	A	517	GOL	O2-C2-C3-O3
4	A	515	GOL	O1-C1-C2-O2
4	A	515	GOL	O2-C2-C3-O3
4	A	516	GOL	O1-C1-C2-O2
4	A	507	GOL	O1-C1-C2-O2
4	A	514	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	517	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	463/501 (92%)	0.28	44 (9%) 14 14	10, 37, 79, 126	9 (1%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443[A]	ARG	6.7
1	A	199	GLY	5.3
1	A	140[A]	GLU	5.0
1	A	198	LEU	4.9
1	A	469[A]	ARG	4.8
1	A	112	VAL	4.5
1	A	482[A]	GLU	4.5
1	A	266	PRO	4.2
1	A	197	ASN	4.0
1	A	122	VAL	4.0
1	A	141	GLY	3.6
1	A	111	VAL	3.5
1	A	160	GLY	3.1
1	A	125	GLY	3.1
1	A	113	VAL	3.0
1	A	158	PRO	3.0
1	A	248	VAL	3.0
1	A	460	ALA	2.8
1	A	159	ASP	2.8
1	A	124	LEU	2.8
1	A	200	GLN	2.7
1	A	2	LYS	2.7
1	A	196	ASP	2.7
1	A	132	ARG	2.7
1	A	194	LEU	2.7
1	A	193	LYS	2.7
1	A	290	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	188	LEU	2.5
1	A	195	TRP	2.5
1	A	162	PHE	2.5
1	A	185	THR	2.5
1	A	108	VAL	2.5
1	A	201	PRO	2.5
1	A	368	GLY	2.4
1	A	265	MET	2.4
1	A	246	LYS	2.3
1	A	157	THR	2.3
1	A	459	MET	2.3
1	A	184	PRO	2.1
1	A	462	ARG	2.1
1	A	244	ARG	2.1
1	A	247	THR	2.0
1	A	123	LEU	2.0
1	A	429	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	511	6/6	0.59	0.20	60,65,67,67	0
4	GOL	A	518	6/6	0.61	0.24	54,55,58,60	0
4	GOL	A	516	6/6	0.62	0.24	57,70,72,74	0
4	GOL	A	510	6/6	0.62	0.24	80,81,81,81	0
4	GOL	A	513	6/6	0.70	0.20	78,82,82,83	0
4	GOL	A	512	6/6	0.72	0.20	57,60,60,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	504	6/6	0.75	0.23	61,68,69,70	0
4	GOL	A	519	6/6	0.76	0.15	47,56,57,60	0
4	GOL	A	509	6/6	0.78	0.17	21,44,45,45	0
4	GOL	A	517	6/6	0.78	0.15	59,61,66,66	0
4	GOL	A	505	6/6	0.79	0.21	66,67,67,69	0
4	GOL	A	514	6/6	0.80	0.17	77,80,81,81	0
4	GOL	A	515	6/6	0.83	0.17	33,63,68,69	0
4	GOL	A	520	6/6	0.83	0.19	93,93,93,93	6
2	NA	A	501	1/1	0.86	0.11	68,68,68,68	0
4	GOL	A	507	6/6	0.86	0.17	49,57,61,62	0
4	GOL	A	508	6/6	0.86	0.14	64,65,67,70	0
3	EPE	A	502	15/15	0.88	0.13	41,47,57,59	0
4	GOL	A	503	6/6	0.90	0.10	28,45,49,53	0
4	GOL	A	506	6/6	0.91	0.10	29,42,50,57	0
5	UNL	A	521	9/-	0.91	0.07	39,40,41,42	9

6.5 Other polymers [i](#)

There are no such residues in this entry.