



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:41 AM UTC

PDB ID : 8DAX / pdb_00008dax
Title : New insights into the P186 flip and oligomeric state of Staphylococcus aureus exfoliative toxin E: implications for the exfoliative mechanism
Authors : Gismene, C.; Nascimento, A.F.Z.; Hernandez-Gonzalez, J.E.; Santisteban, A.R.N.; de Moraes, F.R.; Arni, R.K.; Mariutti, R.B.
Deposited on : 2022-06-14
Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	12% 85% 9% .
1	B	259	13% 87% 8% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exfoliative toxin E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	1	0
			1926	1224	321	380	1			
1	B	248	Total	C	N	O	S	0	1	0
			1930	1225	324	380	1			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	initiating methionine	UNP L0RUV7
A	-8	GLY	-	expression tag	UNP L0RUV7
A	-7	SER	-	expression tag	UNP L0RUV7
A	-6	SER	-	expression tag	UNP L0RUV7
A	-5	HIS	-	expression tag	UNP L0RUV7
A	-4	HIS	-	expression tag	UNP L0RUV7
A	-3	HIS	-	expression tag	UNP L0RUV7
A	-2	HIS	-	expression tag	UNP L0RUV7
A	-1	HIS	-	expression tag	UNP L0RUV7
A	0	HIS	-	expression tag	UNP L0RUV7
A	1	MET	-	expression tag	UNP L0RUV7
B	-9	MET	-	initiating methionine	UNP L0RUV7
B	-8	GLY	-	expression tag	UNP L0RUV7
B	-7	SER	-	expression tag	UNP L0RUV7
B	-6	SER	-	expression tag	UNP L0RUV7
B	-5	HIS	-	expression tag	UNP L0RUV7
B	-4	HIS	-	expression tag	UNP L0RUV7
B	-3	HIS	-	expression tag	UNP L0RUV7
B	-2	HIS	-	expression tag	UNP L0RUV7
B	-1	HIS	-	expression tag	UNP L0RUV7
B	0	HIS	-	expression tag	UNP L0RUV7
B	1	MET	-	expression tag	UNP L0RUV7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	63	Total 63	O 63	0	0
2	B	82	Total 82	O 82	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Exfoliative toxin E



• Molecule 1: Exfoliative toxin E



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.74Å 97.74Å 116.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.92 – 1.61 48.92 – 1.61	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.92-1.61) 99.9 (48.92-1.61)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.214 , 0.248 0.224 , 0.255	Depositor DCC
R_{free} test set	3650 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 30.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4001	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	0/1966	1.38	8/2658 (0.3%)
1	B	1.10	2/1969 (0.1%)	1.29	2/2661 (0.1%)
All	All	1.08	2/3935 (0.1%)	1.34	10/5319 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	171	GLU	C-O	-5.45	1.17	1.24
1	B	208	GLY	C-O	5.01	1.27	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLU	O-C-N	-7.64	114.83	121.32
1	A	69	ARG	NE-CZ-NH1	6.84	128.34	121.50
1	A	69	ARG	CB-CG-CD	6.60	126.48	111.30
1	A	225	SER	CA-C-N	6.45	128.92	120.28
1	A	225	SER	C-N-CA	6.45	128.92	120.28
1	A	111	GLY	CA-C-O	-5.94	117.48	122.29
1	A	69	ARG	CD-NE-CZ	5.56	132.19	124.40
1	B	157	PRO	CA-C-N	5.45	127.58	120.28
1	B	157	PRO	C-N-CA	5.45	127.58	120.28
1	A	69	ARG	NE-CZ-NH2	-5.41	114.33	119.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	185	GLU	Peptide
1	A	186[A]	PRO	Mainchain
1	A	186[B]	PRO	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	1917	14	0
1	B	1930	0	1923	10	0
2	A	63	0	0	2	0
2	B	82	0	0	1	0
All	All	4001	0	3840	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:209:GLY:O	2:B:301:HOH:O	1.62	1.15
1:A:158:TYR:OH	1:B:72:GLU:OE2	2.02	0.77
1:B:144:ASP:OD1	1:B:146:GLN:NE2	2.28	0.66
1:A:163:HIS:HD2	2:A:325:HOH:O	1.76	0.66
1:A:230:ASP:HB3	1:A:232:THR:HG22	1.84	0.58
1:B:176:GLN:NE2	1:B:176:GLN:HA	2.22	0.55
1:A:69:ARG:HD3	1:B:66:HIS:NE2	2.23	0.53
1:A:10:GLN:C	1:A:10:GLN:HE21	2.21	0.48
1:A:116:LEU:C	1:A:116:LEU:HD23	2.39	0.47
1:B:58:LYS:HA	1:B:131:GLY:HA3	1.95	0.47
1:A:65:LYS:HA	1:A:117:SER:OG	2.16	0.46
1:B:65:LYS:HA	1:B:117:SER:OG	2.16	0.46
1:A:69:ARG:HD3	1:B:66:HIS:CE1	2.52	0.45
1:A:143:VAL:HB	1:A:220:ASN:HB3	1.99	0.44
1:A:188:ASN:O	1:A:189:SER:C	2.62	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:ILE:HG22	1:B:182:ALA:HB2	2.01	0.43
1:A:48:LYS:HA	1:A:48:LYS:HD3	1.93	0.41
1:A:139:ILE:O	1:A:241:LYS:HE3	2.20	0.41
1:B:116:LEU:HD23	1:B:116:LEU:C	2.46	0.41
1:A:69:ARG:HA	1:A:107:GLU:CD	2.46	0.41
1:A:83:GLY:HA2	2:A:337:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/259 (95%)	235 (95%)	12 (5%)	0	100	100
1	B	247/259 (95%)	235 (95%)	11 (4%)	1 (0%)	30	14
All	All	494/518 (95%)	470 (95%)	23 (5%)	1 (0%)	43	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/222 (96%)	206 (97%)	7 (3%)	33	11
1	B	213/222 (96%)	206 (97%)	7 (3%)	33	11
All	All	426/444 (96%)	412 (97%)	14 (3%)	35	11

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	10	GLN
1	A	21	SER
1	A	69	ARG
1	A	167	LYS
1	A	222	GLN
1	A	245	LYS
1	B	2	ILE
1	B	7	GLU
1	B	11	LYS
1	B	69[A]	ARG
1	B	69[B]	ARG
1	B	178	PHE
1	B	221	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	163	HIS
1	A	210	GLN
1	B	176	GLN
1	B	247	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/259 (95%)	0.93	31 (12%) 8 8	13, 35, 63, 97	1 (0%)
1	B	248/259 (95%)	0.77	34 (13%) 6 6	12, 30, 63, 88	1 (0%)
All	All	496/518 (95%)	0.85	65 (13%) 7 7	12, 33, 63, 97	2 (0%)

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ILE	7.2
1	B	177	THR	5.5
1	B	173	PHE	5.1
1	A	2	ILE	5.0
1	B	9	ILE	4.4
1	B	213	LEU	4.3
1	A	1	MET	4.2
1	B	4	TYR	4.2
1	A	158	TYR	4.1
1	B	23	SER	3.7
1	B	186	PRO	3.7
1	B	211	TYR	3.7
1	A	233	VAL	3.6
1	B	176	GLN	3.6
1	B	209	GLY	3.5
1	B	178	PHE	3.5
1	B	175	ASN	3.3
1	B	208	GLY	3.1
1	A	5	THR	3.1
1	A	229	THR	3.0
1	A	230	ASP	3.0
1	B	210	GLN	3.0
1	A	232	THR	3.0
1	A	126	TYR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	11	LYS	2.9
1	B	1	MET	2.9
1	A	196	LEU	2.9
1	A	30	GLN	2.9
1	B	16	PHE	2.9
1	B	18	THR	2.9
1	B	231	LYS	2.7
1	B	144	ASP	2.7
1	B	15	PHE	2.6
1	B	158	TYR	2.6
1	A	234	THR	2.6
1	B	5	THR	2.5
1	B	7	GLU	2.5
1	A	240	LEU	2.5
1	A	134	VAL	2.5
1	B	148	GLY	2.4
1	B	143	VAL	2.4
1	A	246	THR	2.3
1	B	172	VAL	2.3
1	A	11	LYS	2.3
1	B	212	GLY	2.3
1	A	6	ASP	2.2
1	A	9	ILE	2.2
1	B	147	LYS	2.2
1	B	6	ASP	2.2
1	A	142	ASN	2.2
1	A	238	ILE	2.2
1	A	141	ASP	2.1
1	A	231	LYS	2.1
1	B	22	ASP	2.1
1	A	166	TYR	2.1
1	A	7	GLU	2.1
1	B	174	ASN	2.1
1	A	144	ASP	2.1
1	B	69[A]	ARG	2.1
1	A	139	ILE	2.1
1	A	197	HIS	2.0
1	A	76	ASN	2.0
1	A	4	TYR	2.0
1	A	23	SER	2.0
1	A	226	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.