



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:12 AM UTC

PDB ID : 6DBD / pdb\_00006dbd  
Title : Crystal Structure of VHH R326  
Authors : Brooks, C.L.; Toride King, M.; Huh, I.  
Deposited on : 2018-05-03  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	 5% 82% 5% 14%
1	B	129	 6% 82% 5% 12%
1	C	129	 3% 81% 5% 14%
1	D	129	 5% 81% 6% 13%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6911 atoms, of which 3135 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called nanobody VHH R326.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	111	Total 1578	C 502	H 765	N 144	O 164	S 3	0	1	0
1	B	114	Total 1646	C 517	H 803	N 155	O 168	S 3	0	0	0
1	C	111	Total 1578	C 501	H 763	N 146	O 165	S 3	0	0	0
1	D	112	Total 1638	C 513	H 801	N 153	O 168	S 3	0	0	0

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Na 1	0	0

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	107	Total	O	0	0
			107	107		
4	B	129	Total	O	0	0
			129	129		
4	C	114	Total	O	0	0
			114	114		
4	D	113	Total	O	0	0
			113	113		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.70Å 97.70Å 243.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.85 – 1.75 48.85 – 1.75	Depositor EDS
% Data completeness (in resolution range)	97.9 (48.85-1.75) 97.9 (48.85-1.75)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	12.61 (at 1.75Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.160 , 0.182 0.163 , 0.185	Depositor DCC
$R_{free}$ test set	3454 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.1	Xtrriage
Anisotropy	0.167	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 55.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6911	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.47	0/827	0.60	0/1123
1	B	0.64	1/854 (0.1%)	0.70	1/1158 (0.1%)
1	C	0.53	1/826 (0.1%)	0.63	0/1122
1	D	0.54	0/848	0.70	0/1149
All	All	0.55	2/3355 (0.1%)	0.66	1/4552 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	81	MET	SD-CE	-10.01	1.54	1.79
1	C	81	MET	SD-CE	-5.22	1.66	1.79

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	81	MET	CG-SD-CE	-7.88	83.57	100.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	813	765	765	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	843	803	802	8	0
1	C	815	763	762	4	0
1	D	837	801	800	4	0
2	B	1	0	0	0	0
3	D	4	3	3	1	0
4	A	107	0	0	2	1
4	B	129	0	0	4	1
4	C	114	0	0	1	1
4	D	113	0	0	2	3
All	All	3776	3135	3132	19	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:ARG:N	4:A:201:HOH:O	1.95	0.99
1:C:58:ARG:NH1	4:C:201:HOH:O	1.90	0.99
1:B:85:ARG:NH1	4:B:301:HOH:O	2.04	0.90
1:D:39:GLN:O	4:D:302:HOH:O	1.97	0.81
1:B:108:GLN:HB3	4:B:308:HOH:O	1.82	0.78
1:B:40:GLY:C	4:B:303:HOH:O	2.35	0.68
1:D:102:ASN:N	4:D:301:HOH:O	1.79	0.68
1:B:53:ARG:HH22	3:D:201:ACT:H2	1.70	0.55
1:C:4:LEU:HD22	1:C:24:THR:HG22	1.96	0.47
1:B:85:ARG:NE	4:B:306:HOH:O	2.48	0.46
1:A:24:THR:HG23	4:A:214:HOH:O	2.17	0.44
1:A:101:VAL:O	1:A:103:PRO:HD3	2.17	0.44
1:D:4:LEU:CD2	1:D:24:THR:HG22	2.49	0.42
1:B:81:MET:HB2	1:B:81:MET:HE3	1.96	0.41
1:B:33:LEU:HD23	1:B:33:LEU:C	2.44	0.41
1:C:9:GLY:HA3	1:C:110:THR:CG2	2.51	0.41
1:C:4:LEU:CD2	1:C:24:THR:HG22	2.51	0.41
1:D:9:GLY:HA3	1:D:110:THR:CG2	2.51	0.41
1:B:4:LEU:CD2	1:B:24:THR:HG22	2.51	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:226:HOH:O	4:D:376:HOH:O[6_554]	1.96	0.24
4:B:418:HOH:O	4:D:402:HOH:O[6_554]	2.00	0.20
4:A:277:HOH:O	4:D:323:HOH:O[10_665]	2.04	0.16

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/129 (84%)	106 (98%)	2 (2%)	0	100	100
1	B	110/129 (85%)	110 (100%)	0	0	100	100
1	C	107/129 (83%)	105 (98%)	2 (2%)	0	100	100
1	D	108/129 (84%)	107 (99%)	1 (1%)	0	100	100
All	All	433/516 (84%)	428 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/105 (79%)	82 (99%)	1 (1%)	63	49
1	B	87/105 (83%)	87 (100%)	0	100	100
1	C	83/105 (79%)	82 (99%)	1 (1%)	63	49
1	D	88/105 (84%)	86 (98%)	2 (2%)	44	24
All	All	341/420 (81%)	337 (99%)	4 (1%)	63	49

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	SER
1	C	99	SER
1	D	75	ASN
1	D	99	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	13	GLN
1	D	75	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	D	201	-	3,3,3	0.93	0	3,3,3	1.36	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	201	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	111/129 (86%)	0.16	7 (6%) 26 30	19, 36, 70, 93	1 (0%)
1	B	114/129 (88%)	-0.18	8 (7%) 22 26	18, 29, 62, 85	0
1	C	111/129 (86%)	-0.11	4 (3%) 46 52	18, 32, 56, 81	0
1	D	112/129 (86%)	0.04	6 (5%) 31 35	21, 33, 62, 83	0
All	All	448/516 (86%)	-0.02	25 (5%) 30 34	18, 33, 65, 93	1 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	101	VAL	5.4
1	A	102	ASN	5.0
1	A	101	VAL	4.8
1	A	116	SER	4.2
1	B	40	GLY	3.3
1	C	116	SER	3.3
1	D	101	VAL	3.3
1	A	1	GLN	3.3
1	D	1	GLN	3.2
1	B	1	GLN	3.2
1	C	1	GLN	2.8
1	D	108	GLN	2.7
1	B	101	VAL	2.7
1	D	117	GLU	2.7
1	A	38	ARG	2.6
1	D	63	ALA	2.5
1	B	116	SER	2.5
1	B	43	LYS	2.3
1	C	39	GLN	2.2
1	B	102	ASN	2.2
1	D	28	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	44	GLN	2.1
1	A	39	GLN	2.1
1	B	39	GLN	2.1
1	A	100	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
3	ACT	D	201	4/4	0.79	0.29	23,51,61,67	0
2	NA	B	201	1/1	0.97	0.07	28,28,28,28	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.