



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:30 PM UTC

PDB ID : 6DBM / pdb_00006dbm
Title : Tyk2 with compound 23
Authors : Vajdos, F.F.
Deposited on : 2018-05-03
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2387 atoms, of which 21 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-receptor tyrosine-protein kinase TYK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	274	2238	1441	383	400	1	13	0	2	0

There are 29 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	865	MET	-	expression tag	UNP P29597
A	866	ALA	-	expression tag	UNP P29597
A	867	HIS	-	expression tag	UNP P29597
A	868	HIS	-	expression tag	UNP P29597
A	869	HIS	-	expression tag	UNP P29597
A	870	HIS	-	expression tag	UNP P29597
A	871	HIS	-	expression tag	UNP P29597
A	872	HIS	-	expression tag	UNP P29597
A	873	HIS	-	expression tag	UNP P29597
A	874	HIS	-	expression tag	UNP P29597
A	875	HIS	-	expression tag	UNP P29597
A	876	HIS	-	expression tag	UNP P29597
A	877	GLY	-	expression tag	UNP P29597
A	878	ALA	-	expression tag	UNP P29597
A	879	LEU	-	expression tag	UNP P29597
A	880	GLU	-	expression tag	UNP P29597
A	881	VAL	-	expression tag	UNP P29597
A	882	LEU	-	expression tag	UNP P29597
A	883	PHE	-	expression tag	UNP P29597
A	884	GLN	-	expression tag	UNP P29597
A	885	GLY	-	expression tag	UNP P29597
A	886	PRO	-	expression tag	UNP P29597
A	887	GLY	-	expression tag	UNP P29597
A	936	ALA	CYS	engineered mutation	UNP P29597
A	969	ALA	GLN	engineered mutation	UNP P29597
A	971	ALA	GLU	engineered mutation	UNP P29597
A	972	ALA	LYS	engineered mutation	UNP P29597

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1016	SER	ALA	conflict	UNP P29597
A	1142	ALA	CYS	engineered mutation	UNP P29597

- Molecule 2 is [(1S)-2,2-difluorocyclopropyl][(1R,5S)-3-{2-[(1-methyl-1H-pyrazol-4-yl)amino]pyrimidin-4-yl}-3,8-diazabicyclo[3.2.1]octan-8-yl]methanone (CCD ID: G4J) (formula: C₁₈H₂₁F₂N₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	H	N	O	21	0
			49	18	2	21	7	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	100	Total	O	0	0
			100	100		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.24Å 73.32Å 101.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.61 – 2.37 50.61 – 2.37	Depositor EDS
% Data completeness (in resolution range)	92.6 (50.61-2.37) 92.6 (50.61-2.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.37Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.201 , 0.264 0.203 , 0.263	Depositor DCC
R_{free} test set	544 reflections (1.08%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.951	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2387	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, G4J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2283	0.67	0/3083

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2238	0	2209	76	0
2	A	28	21	0	1	0
3	A	100	0	0	8	2
All	All	2366	21	2209	76	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1116:GLN:NE2	3:A:4101:HOH:O	2.08	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1124:LEU:HA	1:A:1127:LEU:HD23	1.69	0.74
1:A:1166:ILE:HB	1:A:1167:PRO:HD3	1.69	0.72
1:A:1096:THR:HG22	1:A:1137:PRO:HB3	1.72	0.72
1:A:1034:ASP:O	1:A:1035:ARG:HD2	1.90	0.71
1:A:894:LYS:HA	1:A:897:LEU:HD12	1.73	0.71
1:A:1104:PRO:HB2	1:A:1105:PRO:HD3	1.74	0.70
1:A:947:GLU:OE2	3:A:4102:HOH:O	2.10	0.68
1:A:964:GLY:O	1:A:977[B]:VAL:HG22	1.94	0.67
1:A:892:PHE:CE1	1:A:977[B]:VAL:HG21	2.30	0.66
1:A:1009:GLU:OE2	3:A:4103:HOH:O	2.14	0.65
1:A:949:ASP:O	1:A:953:THR:HG23	1.97	0.65
1:A:1125:THR:O	1:A:1129:GLU:HG3	1.98	0.64
1:A:1126:GLU:O	1:A:1130:ARG:HG3	1.98	0.63
1:A:951:LEU:HD23	1:A:976:LEU:HD21	1.81	0.61
1:A:968:ASP:HB2	1:A:975:GLN:HG3	1.82	0.61
1:A:894:LYS:HG3	1:A:897:LEU:HD12	1.83	0.61
1:A:1163:GLU:HG2	3:A:4188:HOH:O	2.01	0.60
1:A:894:LYS:HD3	1:A:967:GLU:O	2.03	0.59
1:A:1124:LEU:HA	1:A:1127:LEU:CD2	2.32	0.59
1:A:1034:ASP:OD1	1:A:1035:ARG:HD3	2.02	0.59
1:A:1117:GLY:N	3:A:4105:HOH:O	2.36	0.57
1:A:908:PHE:O	1:A:933:LYS:HG3	2.04	0.56
1:A:932:LEU:HD22	1:A:944:TRP:HB2	1.88	0.56
1:A:1090:THR:O	1:A:1094:LEU:HD13	2.05	0.56
1:A:964:GLY:C	1:A:977[B]:VAL:HG22	2.30	0.56
1:A:957:GLU:HG3	1:A:958:HIS:CD2	2.42	0.54
1:A:1106:THR:O	1:A:1110:GLU:HG3	2.07	0.54
1:A:1104:PRO:O	1:A:1108:PHE:HB2	2.10	0.52
1:A:999:GLN:NE2	1:A:1034:ASP:HB2	2.25	0.51
1:A:948:ILE:HG23	1:A:976:LEU:HD13	1.93	0.51
1:A:1177:GLN:O	1:A:1177:GLN:HG2	2.10	0.51
1:A:1092:TYR:O	1:A:1096:THR:HG23	2.11	0.50
1:A:987:ARG:HH21	1:A:1027[B]:ARG:HD3	1.78	0.49
1:A:1138:ASP:O	1:A:1139:LYS:HB2	2.13	0.49
1:A:929:VAL:HG22	1:A:977[A]:VAL:HG22	1.93	0.48
1:A:1118:GLN:NE2	3:A:4107:HOH:O	2.37	0.48
1:A:1170:LYS:O	1:A:1174:GLU:HG3	2.12	0.48
1:A:997:LEU:O	1:A:1001:LEU:HG	2.13	0.48
1:A:1002:LEU:O	1:A:1006:GLN:HG3	2.14	0.47
1:A:1176:TYR:C	1:A:1178:GLY:H	2.22	0.47
1:A:1091:LEU:HD23	1:A:1148:MET:HG3	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:932:LEU:HD22	1:A:944:TRP:CB	2.45	0.47
1:A:1034:ASP:C	1:A:1035:ARG:HD2	2.39	0.47
1:A:1021:HIS:O	1:A:1022:ARG:HB2	2.15	0.46
1:A:1023:ASP:OD1	1:A:1027[B]:ARG:NH1	2.47	0.46
1:A:995:ILE:CG2	1:A:999:GLN:HB3	2.46	0.46
1:A:898:LYS:HB3	1:A:915:CYS:HB3	1.98	0.46
1:A:975:GLN:NE2	3:A:4112:HOH:O	2.49	0.46
1:A:994:SER:HB2	3:A:4138:HOH:O	2.17	0.45
1:A:981:VAL:HG12	1:A:1031:LEU:O	2.16	0.45
1:A:1124:LEU:O	1:A:1128:LEU:HG	2.17	0.45
1:A:1161:THR:OG1	1:A:1163:GLU:HB2	2.17	0.45
1:A:1150:ASN:OD1	1:A:1160:PRO:HD3	2.17	0.45
1:A:941:ARG:NH2	1:A:972:ALA:HB3	2.32	0.45
1:A:893:HIS:C	1:A:895:ARG:H	2.25	0.44
1:A:1104:PRO:CB	1:A:1105:PRO:HD3	2.46	0.44
1:A:997:LEU:HD21	1:A:1141:PRO:HG2	2.00	0.44
1:A:892:PHE:HE1	1:A:963:LYS:O	2.01	0.44
1:A:944:TRP:CE2	1:A:948:ILE:HD11	2.52	0.44
1:A:1124:LEU:CA	1:A:1127:LEU:HD23	2.43	0.44
1:A:951:LEU:CD2	1:A:976:LEU:HD21	2.46	0.43
1:A:905:GLU:HA	2:A:4000:G4J:F2	2.08	0.43
1:A:999:GLN:HE22	1:A:1034:ASP:HB2	1.83	0.43
1:A:894:LYS:HA	1:A:897:LEU:CD1	2.46	0.43
1:A:980:TYR:CE2	1:A:982:PRO:HA	2.54	0.42
1:A:1145:TYR:CD1	1:A:1148:MET:HE2	2.54	0.42
1:A:1172:VAL:O	1:A:1176:TYR:HD2	2.02	0.42
1:A:950:ILE:O	1:A:954:LEU:HG	2.20	0.42
1:A:1031:LEU:HD22	1:A:1033:ASN:O	2.19	0.42
1:A:1079:TYR:O	1:A:1082:SER:HB2	2.20	0.42
1:A:1175:LYS:HD3	1:A:1176:TYR:CE2	2.55	0.42
1:A:983:LEU:HB2	1:A:1031:LEU:HB3	2.02	0.41
1:A:1162:PHE:O	1:A:1166:ILE:HG13	2.21	0.41
1:A:944:TRP:NE1	1:A:948:ILE:HD11	2.36	0.40
1:A:1013:TYR:O	1:A:1017:GLN:HG2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:4108:HOH:O	3:A:4130:HOH:O[3_555]	1.88	0.32
3:A:4184:HOH:O	3:A:4186:HOH:O[3_555]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/318 (84%)	258 (97%)	8 (3%)	1 (0%)	30	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	971	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/271 (88%)	227 (96%)	10 (4%)	26	35

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	927	VAL
1	A	942	SER
1	A	967	GLU
1	A	1031	LEU
1	A	1036	LEU
1	A	1037	VAL
1	A	1056	ARG
1	A	1122	LEU
1	A	1127	LEU
1	A	1147	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	1054	1	15,16,17	1.33	1 (6%)	17,22,24	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1054	1	-	0/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1054	PTR	OH-CZ	-4.42	1.30	1.40

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	G4J	A	4000	-	30,32,32	1.64	6 (20%)	40,49,49	1.34	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G4J	A	4000	-	-	0/16/45/45	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	4000	G4J	C2-N1	4.64	1.39	1.34
2	A	4000	G4J	C15-N7	3.60	1.43	1.35
2	A	4000	G4J	C8-N5	3.54	1.39	1.34
2	A	4000	G4J	C5-N4	3.26	1.39	1.34
2	A	4000	G4J	C4-N2	2.56	1.36	1.32
2	A	4000	G4J	C8-N6	-2.26	1.32	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	4000	G4J	C1-N1-C2	-3.57	124.36	127.94
2	A	4000	G4J	C1-N1-N2	3.11	122.52	120.03
2	A	4000	G4J	C18-C17-C16	-2.46	58.87	60.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	4000	G4J	C7-C8-N5	-2.31	119.83	123.62
2	A	4000	G4J	C4-N2-N1	2.26	105.65	104.41
2	A	4000	G4J	C12-C13-C14	-2.02	109.80	111.70

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4000	G4J	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	273/318 (85%)	0.02	10 (3%)	45 51	9, 25, 59, 77	2 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	889	PRO	3.9
1	A	969	ALA	3.6
1	A	890	THR	3.4
1	A	896	TYR	3.2
1	A	971	ALA	2.9
1	A	955	TYR	2.6
1	A	973	SER	2.4
1	A	1061	GLY	2.4
1	A	1178	GLY	2.2
1	A	970	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
1	PTR	A	1054	16/17	0.90	0.09	27,44,69,70	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	G4J	A	4000	28/28	0.96	0.06	11,18,25,26	21

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.