



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 10:59 AM UTC

PDB ID : 2DCQ / pdb_00002dcq
Title : Fully automated NMR structure determination of the rhodanese homology domain At4g01050(175-295) from Arabidopsis thaliana
Authors : Lopez-Mendez, B.; Guntert, P.
Deposited on : 2006-01-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	134	78% 10% 11%

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:7-A:125 (119)	0.52	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20
2	4, 6
Single-model clusters	19

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2002 atoms, of which 1001 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Putative protein At4g01050.

Mol	Chain	Residues	Atoms					Trace
1	A	134	Total	C	H	N	O	0
			2002	631	1001	170	200	

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP Q9M158
A	2	SER	-	cloning artifact	UNP Q9M158
A	3	SER	-	cloning artifact	UNP Q9M158
A	4	GLY	-	cloning artifact	UNP Q9M158
A	5	SER	-	cloning artifact	UNP Q9M158
A	6	SER	-	cloning artifact	UNP Q9M158
A	7	GLY	-	cloning artifact	UNP Q9M158
A	129	SER	-	cloning artifact	UNP Q9M158
A	130	GLY	-	cloning artifact	UNP Q9M158
A	131	PRO	-	cloning artifact	UNP Q9M158
A	132	SER	-	cloning artifact	UNP Q9M158
A	133	SER	-	cloning artifact	UNP Q9M158
A	134	GLY	-	cloning artifact	UNP Q9M158

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Putative protein At4g01050




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Putative protein At4g01050




4.2.2 Score per residue for model 2

- Molecule 1: Putative protein At4g01050




4.2.3 Score per residue for model 3

- Molecule 1: Putative protein At4g01050




4.2.4 Score per residue for model 4

- Molecule 1: Putative protein At4g01050




4.2.5 Score per residue for model 5

- Molecule 1: Putative protein At4g01050



4.2.6 Score per residue for model 6


- Molecule 1: Putative protein At4g01050



4.2.7 Score per residue for model 7



- Molecule 1: Putative protein At4g01050





4.2.8 Score per residue for model 8

- Molecule 1: Putative protein At4g01050



4.2.9 Score per residue for model 9

- Molecule 1: Putative protein At4g01050



4.2.10 Score per residue for model 10

- Molecule 1: Putative protein At4g01050




4.2.11 Score per residue for model 11

- Molecule 1: Putative protein At4g01050




4.2.12 Score per residue for model 12

- Molecule 1: Putative protein At4g01050




4.2.13 Score per residue for model 13

- Molecule 1: Putative protein At4g01050




4.2.14 Score per residue for model 14

- Molecule 1: Putative protein At4g01050




4.2.15 Score per residue for model 15

- Molecule 1: Putative protein At4g01050




4.2.16 Score per residue for model 16

- Molecule 1: Putative protein At4g01050




4.2.17 Score per residue for model 17

- Molecule 1: Putative protein At4g01050




4.2.18 Score per residue for model 18

- Molecule 1: Putative protein At4g01050




4.2.19 Score per residue for model 19

- Molecule 1: Putative protein At4g01050



4.2.20 Score per residue for model 20

- Molecule 1: Putative protein At4g01050



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Fully automated NMR spectrum analysis and structure calculation without human intervention. CHEMICAL SHIFT ASSIGNMENTS AND CONFORMATIONAL RESTRAINTS HAVE NOT BEEN VERIFIED MANUALLY..*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
FLYA	structure solution	1.0
CYANA	structure solution	2.1
OPALp	refinement	1.2

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.61±0.01	0±0/930 (0.0± 0.0%)	1.33±0.02	2±1/1258 (0.2± 0.1%)
All	All	0.61	0/18600 (0.0%)	1.33	43/25160 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.9±0.9
All	All	0	18

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	26	ASP	CA-CB-CG	13.24	125.84	112.60	18	7
1	A	107	ASP	CA-CB-CG	9.73	122.33	112.60	18	2
1	A	54	ASN	CA-CB-CG	6.91	119.51	112.60	5	1
1	A	57	ASP	CA-CB-CG	6.44	119.04	112.60	17	3
1	A	19	ASP	CA-C-N	6.36	133.68	121.54	10	2
1	A	19	ASP	C-N-CA	6.36	133.68	121.54	10	2
1	A	81	ASP	CA-C-N	5.98	128.22	120.44	11	2
1	A	81	ASP	C-N-CA	5.98	128.22	120.44	11	2
1	A	55	GLY	CA-C-N	5.60	128.05	120.38	12	1
1	A	55	GLY	C-N-CA	5.60	128.05	120.38	12	1
1	A	84	ASP	CA-CB-CG	5.49	118.09	112.60	4	1
1	A	114	GLY	CA-C-N	5.38	128.27	120.79	19	1
1	A	114	GLY	C-N-CA	5.38	128.27	120.79	19	1
1	A	30	THR	CA-C-N	5.37	128.01	120.28	8	1
1	A	30	THR	C-N-CA	5.37	128.01	120.28	8	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	71	ASP	CA-CB-CG	5.27	117.87	112.60	16	1
1	A	18	THR	CB-CA-C	5.26	116.28	109.28	7	2
1	A	38	SER	CA-C-N	5.24	126.68	120.23	19	3
1	A	38	SER	C-N-CA	5.24	126.68	120.23	19	3
1	A	18	THR	N-CA-C	-5.24	108.65	114.62	7	1
1	A	74	ASN	CA-CB-CG	5.16	117.76	112.60	14	1
1	A	111	GLY	CA-C-N	5.12	124.84	119.82	20	1
1	A	111	GLY	C-N-CA	5.12	124.84	119.82	20	1
1	A	116	LEU	N-CA-C	-5.08	103.77	110.33	17	1
1	A	124	GLU	CB-CA-C	5.07	120.17	110.17	14	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	28	ARG	Sidechain	7
1	A	113	ARG	Sidechain	4
1	A	78	TYR	Sidechain	3
1	A	53	TYR	Sidechain	2
1	A	69	PHE	Sidechain	1
1	A	103	TYR	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	911	914	914	2±1
All	All	18220	18280	18280	30

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:28:ARG:HH11	1:A:109:ALA:HB3	0.70	1.46	16	1
1:A:100:LYS:HA	1:A:100:LYS:HE3	0.53	1.80	14	3
1:A:13:TYR:CE1	1:A:44:LEU:HD11	0.49	2.43	4	3
1:A:47:LYS:HE2	1:A:123:ILE:CD1	0.46	2.40	19	2
1:A:100:LYS:HA	1:A:100:LYS:HE2	0.46	1.87	6	1
1:A:13:TYR:CD2	1:A:120:LEU:HD22	0.46	2.46	17	2
1:A:44:LEU:HD11	1:A:46:LYS:HE2	0.45	1.88	1	1
1:A:15:LYS:HE2	1:A:15:LYS:HA	0.45	1.89	13	1
1:A:60:GLY:O	1:A:63:LYS:HE3	0.45	2.11	18	1
1:A:35:GLN:HG3	1:A:36:VAL:HG23	0.44	1.90	4	1
1:A:60:GLY:O	1:A:63:LYS:HE2	0.44	2.13	7	3
1:A:81:ASP:CG	1:A:82:LYS:N	0.44	2.76	17	3
1:A:53:TYR:CE2	1:A:89:LEU:HD22	0.43	2.49	5	1
1:A:62:LEU:HA	1:A:65:LEU:HD13	0.43	1.91	13	1
1:A:13:TYR:CE2	1:A:44:LEU:HD11	0.42	2.50	15	1
1:A:47:LYS:CE	1:A:123:ILE:CD1	0.41	2.98	11	1
1:A:16:LEU:HB3	1:A:44:LEU:HD13	0.41	1.93	11	1
1:A:100:LYS:HA	1:A:100:LYS:CE	0.40	2.46	9	1
1:A:67:LEU:H	1:A:67:LEU:HD23	0.40	1.74	18	1
1:A:69:PHE:CZ	1:A:77:LEU:HD11	0.40	2.52	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/134 (89%)	102±3 (86±3%)	15±3 (13±2%)	1±1 (1±1%)	14	63
All	All	2380/2680 (89%)	2048 (86%)	305 (13%)	27 (1%)	14	63

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	125	PRO	6
1	A	20	ASP	4
1	A	85	GLY	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	100	LYS	3
1	A	119	SER	2
1	A	84	ASP	2
1	A	86	ASN	1
1	A	117	ASN	1
1	A	124	GLU	1
1	A	73	GLU	1
1	A	45	GLY	1
1	A	98	GLY	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/107 (90%)	79±3 (82±3%)	17±3 (18±3%)	4	37
All	All	1920/2140 (90%)	1578 (82%)	342 (18%)	4	37

All 53 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	24	LEU	20
1	A	25	LEU	20
1	A	46	LYS	20
1	A	105	ILE	20
1	A	63	LYS	19
1	A	80	LEU	18
1	A	47	LYS	17
1	A	77	LEU	16
1	A	38	SER	15
1	A	42	LYS	13
1	A	18	THR	10
1	A	81	ASP	10
1	A	100	LYS	10
1	A	75	THR	9
1	A	32	ASP	8
1	A	67	LEU	8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	26	ASP	7
1	A	124	GLU	6
1	A	89	LEU	6
1	A	66	SER	6
1	A	70	LYS	6
1	A	16	LEU	5
1	A	10	LYS	5
1	A	82	LYS	5
1	A	11	ASN	4
1	A	44	LEU	4
1	A	52	VAL	4
1	A	73	GLU	4
1	A	8	SER	3
1	A	23	GLN	3
1	A	87	SER	3
1	A	92	GLU	3
1	A	74	ASN	3
1	A	106	LYS	3
1	A	84	ASP	3
1	A	83	PHE	3
1	A	118	SER	2
1	A	50	SER	2
1	A	54	ASN	2
1	A	113	ARG	2
1	A	41	ILE	2
1	A	119	SER	2
1	A	86	ASN	1
1	A	101	SER	1
1	A	125	PRO	1
1	A	71	ASP	1
1	A	19	ASP	1
1	A	20	ASP	1
1	A	57	ASP	1
1	A	117	ASN	1
1	A	68	LYS	1
1	A	64	LYS	1
1	A	110	GLU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided