



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:06 AM UTC

PDB ID : 3DC1 / pdb_00003dc1
Title : Crystal structure of kynurenine aminotransferase II complex with alpha-ketoglutarate
Authors : Han, Q.; Cai, T.; Tagle, D.A.; Robinson, H.; Li, J.
Deposited on : 2008-06-03
Resolution : 2.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	 2% 56% 35% 6% •
1	B	425	 2% 55% 36% 6% ••
1	C	425	 4% 56% 37% 7%
1	D	425	 2% 55% 35% 7% ••

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 14067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine/alpha-aminoadipate aminotransferase mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	415	Total 3265	C 2097	N 544	O 606	P 1	S 17	0	0	0
1	B	417	Total 3285	C 2109	N 549	O 609	P 1	S 17	0	0	0
1	C	425	Total 3347	C 2147	N 560	O 621	P 1	S 18	0	0	0
1	D	418	Total 3296	C 2115	N 553	O 610	P 1	S 17	0	0	0

- Molecule 2 is 2-OXOGLUTARIC ACID (CCD ID: AKG) (formula: C₅H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 10	C 5	O 5	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	225	Total O 225 225	0	0
4	B	226	Total O 226 226	0	0
4	C	202	Total O 202 202	0	0
4	D	175	Total O 175 175	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial




- Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial






● Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial



● Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.93Å 72.09Å 109.08Å 89.97° 100.65° 93.88°	Depositor
Resolution (Å)	29.69 – 2.50 29.69 – 2.50	Depositor EDS
% Data completeness (in resolution range)	88.7 (29.69-2.50) 90.1 (29.69-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.48Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.244 0.215 , 0.242	Depositor DCC
R_{free} test set	2890 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtrriage
Anisotropy	0.278	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14067	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AKG, LLP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.12	2/3321 (0.1%)	1.14	5/4507 (0.1%)
1	B	1.12	3/3341 (0.1%)	1.25	17/4532 (0.4%)
1	C	1.21	13/3404 (0.4%)	1.27	22/4620 (0.5%)
1	D	1.12	10/3352 (0.3%)	1.23	22/4546 (0.5%)
All	All	1.14	28/13418 (0.2%)	1.22	66/18205 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	6
1	C	0	1
1	D	0	5
All	All	0	13

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	290	VAL	CA-CB	8.38	1.63	1.54
1	C	50	THR	CA-CB	6.42	1.65	1.53
1	D	166	ILE	CA-CB	6.27	1.60	1.54
1	C	75	SER	CA-C	6.23	1.60	1.52
1	A	324	ASP	CA-C	-6.12	1.45	1.52

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	40	LEU	CA-C-N	-9.11	110.73	120.66
1	D	40	LEU	C-N-CA	-9.11	110.73	120.66
1	C	349	VAL	CA-C-N	7.90	128.38	119.93
1	C	349	VAL	C-N-CA	7.90	128.38	119.93
1	B	338	ASP	N-CA-C	-7.67	102.93	111.82

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	31	LYS	Peptide
1	B	18	PRO	Peptide
1	B	19	ILE	Peptide
1	B	30	PRO	Peptide
1	B	31	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3265	0	3265	189	0
1	B	3285	0	3289	151	0
1	C	3347	0	3356	147	0
1	D	3296	0	3302	165	0
2	A	10	0	4	0	0
3	A	18	0	24	3	0
3	C	12	0	16	5	0
3	D	6	0	8	0	0
4	A	225	0	0	34	0
4	B	226	0	0	19	0
4	C	202	0	0	24	0
4	D	175	0	0	29	0
All	All	14067	0	13264	623	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 623 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:LYS:HD3	1:A:31:LYS:C	1.35	1.38
1:B:31:LYS:NZ	1:B:31:LYS:HB3	1.46	1.23
1:A:31:LYS:C	1:A:31:LYS:CD	2.26	1.07
1:B:31:LYS:HB3	1:B:31:LYS:HZ3	0.99	1.06
1:B:31:LYS:HG2	1:B:32:SER:H	1.17	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/425 (96%)	376 (92%)	32 (8%)	2 (0%)	24	43
1	B	412/425 (97%)	382 (93%)	29 (7%)	1 (0%)	43	63
1	C	422/425 (99%)	395 (94%)	27 (6%)	0	100	100
1	D	413/425 (97%)	375 (91%)	34 (8%)	4 (1%)	12	24
All	All	1657/1700 (98%)	1528 (92%)	122 (7%)	7 (0%)	30	49

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	200	GLY
1	B	203	PRO
1	D	203	PRO
1	A	201	ASN
1	D	163	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/369 (97%)	332 (92%)	27 (8%)	12	26
1	B	361/369 (98%)	321 (89%)	40 (11%)	6	12
1	C	369/369 (100%)	336 (91%)	33 (9%)	9	20
1	D	362/369 (98%)	328 (91%)	34 (9%)	8	18
All	All	1451/1476 (98%)	1317 (91%)	134 (9%)	8	19

5 of 134 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	248	SER
1	D	293	LEU
1	D	376	LYS
1	B	216	ILE
1	B	192	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	348	HIS
1	D	201	ASN
1	C	185	ASN
1	D	155	ASN
1	D	308	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	C	263	1	23,24,25	1.83	3 (13%)	25,32,34	1.72	4 (16%)
1	LLP	D	263	1	23,24,25	1.91	4 (17%)	25,32,34	1.53	4 (16%)
1	LLP	A	263	1	23,24,25	1.88	4 (17%)	25,32,34	1.75	3 (12%)
1	LLP	B	263	1	23,24,25	1.73	3 (13%)	25,32,34	1.73	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	C	263	1	-	6/16/17/19	0/1/1/1
1	LLP	D	263	1	-	11/16/17/19	0/1/1/1
1	LLP	A	263	1	-	9/16/17/19	0/1/1/1
1	LLP	B	263	1	-	7/16/17/19	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	263	LLP	O3-C3	-6.35	1.22	1.36
1	D	263	LLP	O3-C3	-6.32	1.22	1.36
1	C	263	LLP	O3-C3	-6.19	1.22	1.36
1	B	263	LLP	O3-C3	-5.89	1.23	1.36
1	C	263	LLP	P-OP3	-2.90	1.44	1.54

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	263	LLP	OP4-C5'-C5	6.71	121.93	109.36
1	A	263	LLP	OP4-C5'-C5	6.48	121.50	109.36
1	B	263	LLP	OP4-C5'-C5	5.68	120.00	109.36
1	D	263	LLP	OP4-C5'-C5	4.21	117.24	109.36
1	B	263	LLP	C4-C4'-NZ	-3.16	109.48	124.04

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	263	LLP	C5'-OP4-P-OP1
1	A	263	LLP	C5'-OP4-P-OP3
1	C	263	LLP	C5'-OP4-P-OP1
1	D	263	LLP	C4-C4'-NZ-CE
1	D	263	LLP	C5'-OP4-P-OP2

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	263	LLP	2	0
1	D	263	LLP	3	0
1	A	263	LLP	4	0
1	B	263	LLP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	D	426	-	5,5,5	0.32	0	5,5,5	0.99	0
3	GOL	C	426	-	5,5,5	0.66	0	5,5,5	1.72	2 (40%)
3	GOL	A	427	-	5,5,5	0.53	0	5,5,5	0.56	0
3	GOL	A	428	-	5,5,5	0.88	0	5,5,5	1.28	0
2	AKG	A	426	-	9,9,9	2.20	4 (44%)	11,11,11	2.29	4 (36%)
3	GOL	C	427	-	5,5,5	0.52	0	5,5,5	0.69	0
3	GOL	A	429	-	5,5,5	0.74	0	5,5,5	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	426	-	-	4/4/4/4	-
3	GOL	C	426	-	-	2/4/4/4	-
3	GOL	A	427	-	-	1/4/4/4	-
3	GOL	A	428	-	-	4/4/4/4	-
2	AKG	A	426	-	-	3/9/9/9	-
3	GOL	C	427	-	-	2/4/4/4	-
3	GOL	A	429	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	426	AKG	C2-C1	-3.91	1.47	1.53
2	A	426	AKG	O1-C1	2.82	1.29	1.22
2	A	426	AKG	O3-C5	2.81	1.31	1.22
2	A	426	AKG	C4-C5	2.74	1.56	1.50

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	426	AKG	O1-C1-C2	-5.06	115.51	121.81
2	A	426	AKG	O5-C2-C1	-3.08	115.27	119.67
2	A	426	AKG	O2-C1-C2	2.67	120.99	113.59
3	C	426	GOL	O2-C2-C1	-2.52	98.74	109.18
3	C	426	GOL	O2-C2-C3	-2.30	99.66	109.18

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	426	AKG	O1-C1-C2-C3
2	A	426	AKG	O2-C1-C2-C3
3	A	428	GOL	O1-C1-C2-C3
3	C	427	GOL	C1-C2-C3-O3
3	D	426	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	426	GOL	2	0
3	A	427	GOL	1	0
3	A	428	GOL	2	0
3	C	427	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/425 (97%)	0.26	8 (1%) 66 62	18, 32, 47, 84	0
1	B	416/425 (97%)	0.22	7 (1%) 69 65	19, 32, 50, 76	0
1	C	424/425 (99%)	0.27	16 (3%) 44 39	16, 31, 47, 88	0
1	D	417/425 (98%)	0.23	9 (2%) 62 58	17, 33, 51, 73	0
All	All	1671/1700 (98%)	0.25	40 (2%) 59 55	16, 32, 49, 88	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	29	GLY	6.2
1	C	21	THR	5.5
1	C	23	THR	5.4
1	C	22	MET	5.4
1	A	30	PRO	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	263	24/25	0.92	0.10	23,30,33,34	0
1	LLP	B	263	24/25	0.93	0.09	13,29,34,35	0
1	LLP	D	263	24/25	0.93	0.09	21,27,31,31	0
1	LLP	C	263	24/25	0.95	0.09	17,29,33,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	429	6/6	0.66	0.16	45,47,49,49	0
3	GOL	A	428	6/6	0.68	0.16	40,42,44,46	0
3	GOL	C	426	6/6	0.69	0.19	24,26,27,29	0
3	GOL	C	427	6/6	0.77	0.12	38,42,47,48	0
2	AKG	A	426	10/10	0.84	0.11	33,41,47,48	0
3	GOL	A	427	6/6	0.84	0.15	39,42,43,46	0
3	GOL	D	426	6/6	0.91	0.09	29,36,37,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.