



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:47 AM UTC

PDB ID : 5DCD / pdb_00005dcd
Title : Neisseria meningitidis 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase regulated (Tyrosine)
Authors : Heyes, L.C.; Parker, E.
Deposited on : 2015-08-24
Resolution : 2.31 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	90% 7% ..
1	B	351	6% 86% 12% .
1	C	351	% 90% 8% ..
1	D	351	2% 91% 6% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	402	-	-	X	-
4	TMO	A	403	-	-	X	-
4	TMO	C	402	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10756 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phospho-2-dehydro-3-deoxyheptonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	346	2638	1652	478	494	14	0	1	0
1	B	346	2540	1592	450	484	14	0	0	0
1	C	346	2608	1632	467	495	14	0	1	0
1	D	346	2599	1630	465	490	14	0	0	0

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		
2	B	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		
2	D	1	Total	Mn	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0

- Molecule 4 is trimethylamine oxide (CCD ID: TMO) (formula: C₃H₉NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	5	3	1	1	0	0
4	B	1	5	3	1	1	0	0
4	C	1	5	3	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 5 is TYROSINE (CCD ID: TYR) (formula: C₉H₁₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			13	9	1	3		
5	B	1	Total	C	N	O	0	0
			13	9	1	3		
5	C	1	Total	C	N	O	0	0
			13	9	1	3		
5	D	1	Total	C	N	O	0	0
			13	9	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	93	Total	O	0	0
			93	93		
6	B	46	Total	O	0	0
			46	46		
6	C	78	Total	O	0	0
			78	78		
6	D	73	Total	O	0	0
			73	73		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase




- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase



- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase



- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.94Å 141.29Å 74.67Å 90.00° 96.28° 90.00°	Depositor
Resolution (Å)	74.23 – 2.31 74.23 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.5 (74.23-2.31) 99.5 (74.23-2.31)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.75 (at 2.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.194 , 0.239 0.199 , 0.239	Depositor DCC
R_{free} test set	3299 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtrriage
Anisotropy	0.177	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.020 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10756	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMO, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	2/2690 (0.1%)	0.88	2/3647 (0.1%)
1	B	0.74	2/2591 (0.1%)	0.85	1/3528 (0.0%)
1	C	0.81	1/2659 (0.0%)	0.88	1/3612 (0.0%)
1	D	0.78	2/2650 (0.1%)	0.88	3/3595 (0.1%)
All	All	0.78	7/10590 (0.1%)	0.87	7/14382 (0.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	213	SER	C-O	-8.87	1.17	1.24
1	A	213	SER	C-O	-6.85	1.17	1.24
1	B	214	VAL	C-O	-6.70	1.17	1.24
1	A	214	VAL	C-O	-6.54	1.17	1.24
1	D	323	THR	CA-C	5.99	1.57	1.53
1	D	213	SER	C-O	-5.14	1.17	1.24
1	C	214	VAL	C-O	-5.05	1.19	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	222	ILE	CB-CA-C	-6.20	103.45	110.96
1	B	202	ILE	CB-CA-C	-6.09	104.17	111.97
1	A	202	ILE	CB-CA-C	-6.03	104.25	111.97
1	D	222	ILE	CB-CA-C	-5.99	101.91	110.83
1	D	150	ILE	N-CA-C	5.93	116.66	111.56
1	A	222	ILE	CB-CA-C	-5.69	103.10	110.84
1	D	111	ASN	N-CA-C	5.26	116.70	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2638	0	2585	29	0
1	B	2540	0	2396	23	0
1	C	2608	0	2510	32	0
1	D	2599	0	2518	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	3	0
4	A	5	0	9	7	0
4	B	5	0	9	2	0
4	C	5	0	9	4	0
4	D	5	0	9	1	0
5	A	13	0	8	2	0
5	B	13	0	8	0	0
5	C	13	0	8	2	0
5	D	13	0	8	0	0
6	A	93	0	0	2	0
6	B	46	0	0	1	0
6	C	78	0	0	1	0
6	D	73	0	0	3	0
All	All	10756	0	10077	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:167:ARG:HG3	6:D:562:HOH:O	1.48	1.12
4:A:403:TMO:HABB	6:A:589:HOH:O	1.68	0.94
1:C:18:LEU:CD1	1:D:222:ILE:HD11	2.16	0.75
1:C:18:LEU:CD1	1:D:222:ILE:CD1	2.65	0.74
1:B:74:TYR:OH	1:B:305:SER:OG	2.10	0.69
1:A:18:LEU:HD11	1:B:212:LEU:HD22	1.77	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:236:ARG:HE	4:D:402:TMO:HABA	1.62	0.64
1:C:18:LEU:CD1	1:D:222:ILE:HG13	2.27	0.64
1:A:236:ARG:HE	4:A:403:TMO:HAD	1.63	0.63
1:B:166:ALA:H	4:B:402:TMO:HAA	1.63	0.62
1:A:222:ILE:CG1	1:B:18:LEU:HD13	2.30	0.62
1:C:111:ASN:O	1:C:119:PHE:HA	2.01	0.61
1:B:237:GLY:N	1:B:242:ASN:OD1	2.34	0.60
1:A:222:ILE:HG13	1:B:18:LEU:HD13	1.83	0.60
1:C:18:LEU:HD11	1:D:222:ILE:HD11	1.82	0.60
1:A:247:HIS:NE2	3:A:402:SO4:O4	2.35	0.60
1:C:18:LEU:HD11	1:D:222:ILE:CG1	2.32	0.59
1:C:224:HIS:HD2	6:D:563:HOH:O	1.87	0.58
1:A:214:VAL:CG1	1:A:218:GLY:HA2	2.35	0.56
1:C:18:LEU:HD11	1:D:222:ILE:CD1	2.36	0.56
1:C:18:LEU:HD11	1:D:222:ILE:HG13	1.87	0.56
1:C:188:LYS:HZ2	4:C:402:TMO:HAA	1.70	0.56
1:C:18:LEU:CD1	1:D:222:ILE:CG1	2.84	0.55
1:C:188:LYS:NZ	4:C:402:TMO:HAA	2.22	0.55
1:A:111:ASN:O	1:A:119:PHE:HA	2.08	0.54
1:A:247:HIS:NE2	3:A:402:SO4:O3	2.41	0.54
1:B:111:ASN:O	1:B:119:PHE:HA	2.08	0.54
1:A:213:SER:OG	5:A:404:TYR:OH	2.18	0.53
1:A:214:VAL:HG13	1:A:218:GLY:C	2.33	0.53
1:C:18:LEU:HD13	1:D:222:ILE:HG13	1.91	0.53
1:A:188:LYS:NZ	4:A:403:TMO:HADB	2.25	0.52
1:A:236:ARG:HE	4:A:403:TMO:CAD	2.23	0.52
1:C:211:PHE:CZ	1:C:223:VAL:HG11	2.45	0.51
1:A:247:HIS:NE2	3:A:402:SO4:S	2.65	0.51
1:D:151:THR:N	1:D:152:PRO:CD	2.74	0.51
1:C:17:GLU:N	6:C:503:HOH:O	2.44	0.51
1:C:210:HIS:HA	1:C:223:VAL:O	2.11	0.50
1:C:211:PHE:CE2	1:C:223:VAL:HG11	2.46	0.50
1:D:167:ARG:HD3	6:D:572:HOH:O	2.12	0.50
1:C:18:LEU:HD12	1:D:222:ILE:HD11	1.92	0.49
1:D:202:ILE:HD11	1:D:233:VAL:HG23	1.95	0.49
1:B:99:LYS:HD3	1:B:100:PRO:O	2.11	0.49
1:B:176:GLU:OE1	6:B:501:HOH:O	2.19	0.48
1:A:213:SER:CB	5:A:404:TYR:OH	2.61	0.48
1:B:219:HIS:NE2	1:C:219:HIS:CD2	2.82	0.48
1:D:267:ASP:HA	1:D:302:MET:HB2	1.95	0.48
1:B:267:ASP:HA	1:B:302:MET:HB3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:210:HIS:HA	1:D:223:VAL:O	2.14	0.47
1:C:82:ARG:HA	1:C:91:ILE:HD12	1.96	0.47
1:B:268:CYS:HA	1:B:283:VAL:HG11	1.97	0.46
1:B:270:HIS:CD2	4:B:402:TMO:HAD	2.50	0.46
1:A:106:TRP:CZ2	1:A:108:GLY:C	2.94	0.46
1:A:151:THR:N	1:A:152:PRO:CD	2.78	0.46
1:D:150:ILE:C	1:D:152:PRO:HD2	2.41	0.46
1:A:135:LEU:O	1:A:138:MET:HG2	2.16	0.46
1:B:77:ARG:HB3	1:B:329:TRP:CZ2	2.51	0.45
1:D:214:VAL:HA	1:D:219:HIS:O	2.17	0.45
1:C:295:GLY:O	1:C:346:ARG:NH1	2.50	0.45
1:C:213:SER:CB	5:C:403:TYR:OH	2.65	0.44
1:A:236:ARG:NE	4:A:403:TMO:HAD	2.31	0.44
1:B:208:SER:OG	1:B:225:THR:O	2.25	0.44
1:D:77:ARG:HB3	1:D:329:TRP:CZ2	2.52	0.44
1:C:151:THR:N	1:C:152:PRO:CD	2.80	0.44
1:C:270:HIS:CG	4:C:402:TMO:HAAA	2.52	0.44
1:A:170:GLU:HG2	1:B:167:ARG:HG2	1.99	0.44
1:A:202:ILE:HD11	1:A:233:VAL:HG13	2.00	0.44
1:B:59:ILE:HD13	1:B:92:VAL:HB	2.00	0.44
1:A:194:ASN:C	1:A:194:ASN:OD1	2.61	0.43
1:A:18:LEU:HG	1:B:222:ILE:HG13	2.00	0.43
1:B:210:HIS:HA	1:B:223:VAL:O	2.18	0.43
1:C:194:ASN:ND2	1:D:192:ASP:OD2	2.50	0.43
4:A:403:TMO:HAAA	6:A:587:HOH:O	2.18	0.42
1:B:146:PHE:CD1	1:B:152:PRO:HG3	2.54	0.42
1:A:148:ASP:OD1	1:A:148:ASP:C	2.63	0.42
1:A:59:ILE:HD13	1:A:92:VAL:HB	2.02	0.42
1:C:135:LEU:O	1:C:138:MET:HG2	2.19	0.42
1:D:268:CYS:HA	1:D:283:VAL:HG11	2.01	0.42
1:C:18:LEU:HD12	1:C:18:LEU:HA	1.76	0.41
1:C:114:HIS:C	1:C:115:LEU:HD12	2.45	0.41
1:C:148:ASP:C	1:C:148:ASP:OD1	2.63	0.41
1:A:214:VAL:HG12	1:A:215:THR:O	2.21	0.41
1:C:213:SER:OG	5:C:403:TYR:OH	2.21	0.41
1:C:270:HIS:CD2	4:C:402:TMO:HAAA	2.55	0.41
1:B:309:GLU:HB3	1:B:328:GLY:N	2.36	0.41
1:A:59:ILE:HG22	1:A:94:ARG:HG3	2.03	0.40
1:B:109:LEU:HD21	1:B:127:GLN:OE1	2.21	0.40
1:A:292:GLU:O	1:A:349:ARG:NH1	2.55	0.40
1:C:268:CYS:HA	1:C:283:VAL:HG11	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:LYS:HZ2	4:A:403:TMO:HAAA	1.86	0.40
1:A:219[B]:HIS:NE2	1:D:219:HIS:HB2	2.36	0.40
1:B:111:ASN:HA	1:B:121:ILE:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/351 (98%)	337 (98%)	7 (2%)	1 (0%)	36	45
1	B	344/351 (98%)	334 (97%)	9 (3%)	1 (0%)	36	45
1	C	345/351 (98%)	338 (98%)	7 (2%)	0	100	100
1	D	344/351 (98%)	336 (98%)	7 (2%)	1 (0%)	36	45
All	All	1378/1404 (98%)	1345 (98%)	30 (2%)	3 (0%)	43	53

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	294	ASP
1	B	294	ASP
1	D	294	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/289 (94%)	269 (98%)	4 (2%)	57	73
1	B	251/289 (87%)	245 (98%)	6 (2%)	43	60
1	C	265/289 (92%)	259 (98%)	6 (2%)	44	62
1	D	265/289 (92%)	262 (99%)	3 (1%)	65	80
All	All	1054/1156 (91%)	1035 (98%)	19 (2%)	51	69

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	ARG
1	A	115	LEU
1	A	189	ASN
1	A	236	ARG
1	B	17	GLU
1	B	33	GLU
1	B	138	MET
1	B	305	SER
1	B	306	HIS
1	B	317	VAL
1	C	17	GLU
1	C	18	LEU
1	C	83	LYS
1	C	94	ARG
1	C	115	LEU
1	C	269	SER
1	D	236	ARG
1	D	269	SER
1	D	306	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	207	HIS
1	B	43	GLN
1	B	210	HIS
1	B	219	HIS
1	B	293	GLN
1	C	207	HIS
1	C	297	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	43	GLN
1	D	297	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	402	-	4,4,4	0.71	0	6,6,6	0.21	0
5	TYR	B	403	-	12,13,13	0.79	0	13,17,17	0.93	0
4	TMO	B	402	-	4,4,4	5.68	1 (25%)	6,6,6	0.43	0
5	TYR	D	403	-	12,13,13	0.88	1 (8%)	13,17,17	0.70	0
5	TYR	A	404	-	12,13,13	0.86	1 (8%)	13,17,17	0.89	1 (7%)
4	TMO	D	402	-	4,4,4	5.36	1 (25%)	6,6,6	0.74	0
4	TMO	C	402	-	4,4,4	6.10	1 (25%)	6,6,6	0.72	0
4	TMO	A	403	-	4,4,4	4.91	1 (25%)	6,6,6	0.85	0
5	TYR	C	403	-	12,13,13	0.86	1 (8%)	13,17,17	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TYR	B	403	-	-	0/8/8/8	0/1/1/1
5	TYR	A	404	-	-	0/8/8/8	0/1/1/1
5	TYR	C	403	-	-	0/8/8/8	0/1/1/1
5	TYR	D	403	-	-	0/8/8/8	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	402	TMO	OAE-NAC	-12.11	1.26	1.42
4	B	402	TMO	OAE-NAC	-11.25	1.27	1.42
4	D	402	TMO	OAE-NAC	-10.67	1.28	1.42
4	A	403	TMO	OAE-NAC	-9.55	1.29	1.42
5	C	403	TYR	OXT-C	-2.45	1.22	1.30
5	D	403	TYR	OXT-C	-2.24	1.23	1.30
5	A	404	TYR	OXT-C	-2.20	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	404	TYR	CG-CB-CA	-2.05	109.93	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	SO4	3	0
4	B	402	TMO	2	0
5	A	404	TYR	2	0
4	D	402	TMO	1	0
4	C	402	TMO	4	0
4	A	403	TMO	7	0
5	C	403	TYR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/351 (98%)	-0.03	1 (0%) 90 90	19, 36, 54, 76	1 (0%)
1	B	346/351 (98%)	0.45	21 (6%) 27 30	28, 46, 79, 100	0
1	C	346/351 (98%)	0.01	5 (1%) 73 75	23, 38, 60, 84	1 (0%)
1	D	346/351 (98%)	0.17	6 (1%) 69 71	28, 39, 57, 70	0
All	All	1384/1404 (98%)	0.15	33 (2%) 59 62	19, 39, 65, 100	2 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	VAL	4.7
1	B	325	ALA	4.0
1	B	103	THR	3.8
1	B	276	ASP	3.8
1	B	313	ASP	3.7
1	B	275	LYS	3.4
1	B	101	ARG	3.4
1	C	103	THR	3.2
1	B	104	VAL	3.2
1	B	315	PRO	3.1
1	C	5	TYR	2.8
1	B	51	GLY	2.8
1	B	102	THR	2.7
1	C	104	VAL	2.7
1	D	102	THR	2.6
1	B	100	PRO	2.5
1	D	103	THR	2.4
1	D	101	ARG	2.4
1	B	319	GLY	2.4
1	A	104	VAL	2.4
1	B	349	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	277	TYR	2.3
1	B	314	LYS	2.3
1	B	22	ILE	2.3
1	D	100	PRO	2.3
1	B	320	LYS	2.2
1	B	119	PHE	2.2
1	B	318	TYR	2.1
1	C	192[A]	ASP	2.1
1	B	308	VAL	2.1
1	D	278	THR	2.1
1	B	322	ILE	2.0
1	C	102	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TMO	C	402	5/5	0.71	0.23	39,42,48,50	0
4	TMO	D	402	5/5	0.78	0.23	42,42,54,55	0
4	TMO	A	403	5/5	0.80	0.21	35,40,47,49	0
4	TMO	B	402	5/5	0.81	0.18	41,42,46,47	0
3	SO4	A	402	5/5	0.85	0.16	52,58,65,65	0
5	TYR	C	403	13/13	0.91	0.10	33,36,37,37	0
5	TYR	D	403	13/13	0.91	0.10	34,38,40,41	0
5	TYR	B	403	13/13	0.92	0.08	35,37,38,40	0
5	TYR	A	404	13/13	0.94	0.08	36,39,42,47	0
2	MN	B	401	1/1	0.96	0.06	62,62,62,62	0
2	MN	D	401	1/1	0.99	0.05	48,48,48,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MN	C	401	1/1	0.99	0.03	40,40,40,40	0
2	MN	A	401	1/1	1.00	0.04	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.