



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 01:19 AM UTC

PDB ID : 6DCX / pdb\_00006dcx  
Title : iASPP-PP-1c structure and targeting of p53  
Authors : Glover, J.N.M.; Zhou, Y.; Edwards, R.A.  
Deposited on : 2018-05-08  
Resolution : 3.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	81% 7% 12%
1	B	339	79% 10% 11%
2	C	228	75% 14% 12%
2	D	228	78% 12% 10%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 7972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase PP1-alpha catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	300	2411	1543	406	444	18	0	0	0
1	B	303	2434	1557	413	446	18	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	expression tag	UNP P62136
A	-7	PRO	-	expression tag	UNP P62136
A	-6	GLY	-	expression tag	UNP P62136
A	-5	TYR	-	expression tag	UNP P62136
A	-4	GLN	-	expression tag	UNP P62136
A	-3	ASP	-	expression tag	UNP P62136
A	-2	PRO	-	expression tag	UNP P62136
A	-1	ASN	-	expression tag	UNP P62136
A	0	SER	-	expression tag	UNP P62136
B	-8	GLY	-	expression tag	UNP P62136
B	-7	PRO	-	expression tag	UNP P62136
B	-6	GLY	-	expression tag	UNP P62136
B	-5	TYR	-	expression tag	UNP P62136
B	-4	GLN	-	expression tag	UNP P62136
B	-3	ASP	-	expression tag	UNP P62136
B	-2	PRO	-	expression tag	UNP P62136
B	-1	ASN	-	expression tag	UNP P62136
B	0	SER	-	expression tag	UNP P62136

- Molecule 2 is a protein called RelA-associated inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	201	1542	970	259	303	10	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	206	1585	996	270	309	10	0	0	0


There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	601	GLY	-	expression tag	UNP Q8WUF5
C	602	PRO	-	expression tag	UNP Q8WUF5
C	603	GLY	-	expression tag	UNP Q8WUF5
C	604	TYR	-	expression tag	UNP Q8WUF5
C	605	GLN	-	expression tag	UNP Q8WUF5
C	606	ASP	-	expression tag	UNP Q8WUF5
C	607	PRO	-	expression tag	UNP Q8WUF5
D	601	GLY	-	expression tag	UNP Q8WUF5
D	602	PRO	-	expression tag	UNP Q8WUF5
D	603	GLY	-	expression tag	UNP Q8WUF5
D	604	TYR	-	expression tag	UNP Q8WUF5
D	605	GLN	-	expression tag	UNP Q8WUF5
D	606	ASP	-	expression tag	UNP Q8WUF5
D	607	PRO	-	expression tag	UNP Q8WUF5

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain A:  81% 7% 12%





- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit

Chain B:  79% 10% 11%




- Molecule 2: RelA-associated inhibitor

Chain C:  75% 14% 12%



- Molecule 2: RelA-associated inhibitor

Chain D:  78% 12% 10%



Y813	Y814	L817	K822	PRO	GLN	ARG	SER	LYS	VAL
------	------	------	------	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.46Å 135.46Å 165.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.87 – 3.41 48.87 – 3.41	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.87-3.41) 99.5 (48.87-3.41)	Depositor EDS
$R_{merge}$	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 3.40Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.187 , 0.241 0.190 , 0.245	Depositor DCC
$R_{free}$ test set	1115 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	92.0	Xtrriage
Anisotropy	0.277	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 77.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7972	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	103.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.18	0/2467	0.41	0/3334
1	B	0.17	0/2490	0.40	0/3364
2	C	0.16	0/1580	0.39	0/2157
2	D	0.16	0/1622	0.39	0/2211
All	All	0.17	0/8159	0.40	0/11066

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2411	0	2369	14	0
1	B	2434	0	2400	17	0
2	C	1542	0	1447	20	0
2	D	1585	0	1497	18	0
All	All	7972	0	7713	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:687:ASN:HD21	2:D:718:ALA:H	1.30	0.79
2:D:755:MET:HG2	2:D:817:LEU:HD13	1.70	0.73
1:A:116:GLU:HB3	2:D:609:SER:HB2	1.74	0.68
1:A:270:PRO:HB3	1:A:298:PRO:HD3	1.75	0.68
2:C:640:LEU:HG	2:C:644:GLN:HE21	1.63	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/339 (87%)	280 (95%)	16 (5%)	0	100	100
1	B	299/339 (88%)	282 (94%)	17 (6%)	0	100	100
2	C	199/228 (87%)	193 (97%)	5 (2%)	1 (0%)	24	54
2	D	202/228 (89%)	194 (96%)	7 (4%)	1 (0%)	24	54
All	All	996/1134 (88%)	949 (95%)	45 (4%)	2 (0%)	43	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	726	ASP
2	D	726	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/295 (90%)	262 (99%)	3 (1%)	65	74
1	B	267/295 (90%)	263 (98%)	4 (2%)	57	69
2	C	160/182 (88%)	160 (100%)	0	100	100
2	D	165/182 (91%)	162 (98%)	3 (2%)	51	67
All	All	857/954 (90%)	847 (99%)	10 (1%)	63	72

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	609	SER
2	D	612	ARG
2	D	658	GLU
1	B	133	ILE
1	B	198	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	687	ASN
2	D	759	ASN
1	B	117	ASN
2	C	644	GLN
2	C	687	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	300/339 (88%)	-0.55	0 100 100	51, 82, 139, 172	0
1	B	303/339 (89%)	-0.57	0 100 100	53, 85, 131, 172	0
2	C	201/228 (88%)	-0.40	0 100 100	77, 115, 166, 195	0
2	D	206/228 (90%)	-0.39	0 100 100	81, 124, 174, 223	0
All	All	1010/1134 (89%)	-0.49	0 100 100	51, 100, 160, 223	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.