



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 01:43 PM UTC

PDB ID : 1DDO / pdb\_00001ddo  
Title : REDUCED D-AMINO ACID OXIDASE FROM PIG KIDNEY IN COMPLEX WITH IMINO-TRP  
Authors : Todone, F.; Mattevi, A.  
Deposited on : 1997-01-16  
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	
1	C	347	
1	D	347	
1	E	347	
1	F	347	
1	G	347	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	H	347	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DTR	E	350	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 22441 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-AMINO ACID OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	339	Total 2720	C 1749	N 473	O 489	S 9	161	0	0
1	B	339	Total 2720	C 1749	N 473	O 489	S 9	190	0	0
1	C	339	Total 2720	C 1749	N 473	O 489	S 9	173	0	0
1	D	339	Total 2720	C 1749	N 473	O 489	S 9	131	0	0
1	E	339	Total 2720	C 1749	N 473	O 489	S 9	164	0	0
1	F	339	Total 2720	C 1749	N 473	O 489	S 9	154	0	0
1	G	339	Total 2720	C 1749	N 473	O 489	S 9	210	0	0
1	H	339	Total 2720	C 1749	N 473	O 489	S 9	166	0	0


- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			15	11	2	2		
3	B	1	Total	C	N	O	0	0
			15	11	2	2		
3	C	1	Total	C	N	O	0	0
			15	11	2	2		
3	D	1	Total	C	N	O	0	0
			15	11	2	2		
3	E	1	Total	C	N	O	0	0
			15	11	2	2		
3	F	1	Total	C	N	O	0	0
			15	11	2	2		
3	G	1	Total	C	N	O	0	0
			15	11	2	2		
3	H	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 4 is D-TRYPTOPHAN (CCD ID: DTR) (formula: C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	E	1	Total	C	N	O	0	0
			15	11	2	2		
4	F	1	Total	C	N	O	0	0
			15	11	2	2		


- Molecule 5 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	17	Total	O	0	0
			17	17		
5	B	7	Total	O	0	0
			7	7		
5	C	12	Total	O	0	0
			12	12		
5	D	21	Total	O	0	0
			21	21		
5	E	16	Total	O	0	0
			16	16		
5	F	21	Total	O	0	0
			21	21		
5	G	3	Total	O	0	0
			3	3		
5	H	10	Total	O	0	0
			10	10		

### 3 Residue-property plots [i](#)




These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: D-AMINO ACID OXIDASE










● Molecule 1: D-AMINO ACID OXIDASE




● Molecule 1: D-AMINO ACID OXIDASE




● Molecule 1: D-AMINO ACID OXIDASE




● Molecule 1: D-AMINO ACID OXIDASE




● Molecule 1: D-AMINO ACID OXIDASE





● Molecule 1: D-AMINO ACID OXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	325.20Å 137.10Å 196.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10 20.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-3.10) 97.8 (20.00-3.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.18 (at 3.13Å)	Xtrriage
Refinement program	TNT 5E	Depositor
R, $R_{free}$	0.236 , 0.250 0.235 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.4	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.53 , 172.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	22441	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DTR, ITR, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.16	8/2796 (0.3%)	1.79	69/3808 (1.8%)
1	B	1.12	4/2796 (0.1%)	1.75	59/3808 (1.5%)
1	C	1.12	5/2796 (0.2%)	1.72	61/3808 (1.6%)
1	D	1.21	14/2796 (0.5%)	1.78	64/3808 (1.7%)
1	E	1.13	3/2796 (0.1%)	1.71	60/3808 (1.6%)
1	F	1.15	4/2796 (0.1%)	1.76	68/3808 (1.8%)
1	G	1.09	4/2796 (0.1%)	1.72	61/3808 (1.6%)
1	H	1.13	4/2796 (0.1%)	1.74	61/3808 (1.6%)
All	All	1.14	46/22368 (0.2%)	1.75	503/30464 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	1	0
1	E	0	2
1	F	0	1
All	All	1	4

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	177	VAL	CA-CB	-6.38	1.45	1.54
1	D	288	GLN	C-O	6.33	1.31	1.23
1	D	178	ILE	CA-CB	-6.25	1.46	1.54
1	A	225	ASN	C-O	6.08	1.31	1.24
1	D	124	MET	SD-CE	-6.03	1.64	1.79

The worst 5 of 503 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	TYR	N-CA-C	16.25	133.38	113.23
1	B	223	ILE	N-CA-C	14.72	127.19	111.58
1	B	23	TYR	N-CA-C	13.61	130.69	113.55
1	D	23	TYR	N-CA-C	11.40	127.40	113.38
1	D	42	PHE	N-CA-C	11.37	127.26	113.16

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	23	TYR	CA

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	54	PRO	Mainchain
1	E	23	TYR	Sidechain
1	E	58	GLU	Sidechain
1	F	23	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2675	193	1
1	B	2720	0	2675	208	0
1	C	2720	0	2675	188	3
1	D	2720	0	2675	233	1
1	E	2720	0	2675	202	0
1	F	2720	0	2675	198	0
1	G	2720	0	2675	215	0
1	H	2720	0	2675	202	3
2	A	53	0	31	5	0
2	B	53	0	31	3	0
2	C	53	0	31	4	0
2	D	53	0	31	4	0
2	E	53	0	31	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	53	0	31	4	0
2	G	53	0	31	5	0
2	H	53	0	31	5	0
3	A	15	0	8	0	0
3	B	15	0	8	0	0
3	C	15	0	8	0	0
3	D	15	0	8	0	0
3	E	15	0	8	0	0
3	F	15	0	8	1	0
3	G	15	0	8	0	0
3	H	15	0	8	0	0
4	E	15	0	11	7	0
4	F	15	0	11	3	0
5	A	17	0	0	0	0
5	B	7	0	0	3	0
5	C	12	0	0	2	0
5	D	21	0	0	5	0
5	E	16	0	0	0	0
5	F	21	0	0	3	0
5	G	3	0	0	1	0
5	H	10	0	0	0	0
All	All	22441	0	21734	1590	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 1590 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:223:ILE:HG13	1:D:224:TYR:CE1	1.52	1.42
1:D:223:ILE:HG13	1:D:224:TYR:CD1	1.69	1.27
1:A:253:ILE:HG12	1:F:42:PHE:CD1	1.73	1.23
1:E:22:ARG:HH11	1:E:22:ARG:HG3	1.09	1.14
1:F:223:ILE:HG12	1:F:224:TYR:CD1	1.85	1.11

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:123:ASP:OD1	1:H:162:ARG:NH2[4_565]	2.01	0.19

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:332:LYS:NZ	1:D:126:PRO:CG[7_545]	2.11	0.09
1:C:123:ASP:OD2	1:H:35:TYR:OH[4_565]	2.15	0.05
1:C:162:ARG:NH2	1:H:123:ASP:OD1[3_655]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/347 (97%)	310 (92%)	25 (7%)	2 (1%)	21	52
1	B	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	14	44
1	C	337/347 (97%)	312 (93%)	23 (7%)	2 (1%)	21	52
1	D	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	14	44
1	E	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	14	44
1	F	337/347 (97%)	311 (92%)	24 (7%)	2 (1%)	21	52
1	G	337/347 (97%)	306 (91%)	27 (8%)	4 (1%)	10	37
1	H	337/347 (97%)	310 (92%)	23 (7%)	4 (1%)	10	37
All	All	2696/2776 (97%)	2476 (92%)	197 (7%)	23 (1%)	14	44

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	28	GLN
1	E	28	GLN
1	A	28	GLN
1	B	24	HIS
1	G	222	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/298 (97%)	259 (89%)	31 (11%)	6	25
1	B	290/298 (97%)	260 (90%)	30 (10%)	7	27
1	C	290/298 (97%)	260 (90%)	30 (10%)	7	27
1	D	290/298 (97%)	261 (90%)	29 (10%)	7	29
1	E	290/298 (97%)	258 (89%)	32 (11%)	6	24
1	F	290/298 (97%)	257 (89%)	33 (11%)	5	23
1	G	290/298 (97%)	256 (88%)	34 (12%)	5	22
1	H	290/298 (97%)	262 (90%)	28 (10%)	8	30
All	All	2320/2384 (97%)	2073 (89%)	247 (11%)	6	26

5 of 247 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	256	HIS
1	H	28	GLN
1	E	241	THR
1	H	14	SER
1	H	224	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	288	GLN
1	F	63	GLN
1	H	307	HIS
1	H	70	GLN
1	D	307	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ITR	G	349	-	15,16,16	1.02	1 (6%)	17,22,22	0.67	0
2	FAD	C	348	-	58,58,58	1.06	5 (8%)	85,89,89	1.35	11 (12%)
3	ITR	D	349	-	15,16,16	1.02	2 (13%)	17,22,22	0.67	0
2	FAD	E	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.36	11 (12%)
2	FAD	F	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.35	11 (12%)
2	FAD	D	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.35	11 (12%)
2	FAD	G	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.35	11 (12%)
2	FAD	H	348	-	58,58,58	1.08	5 (8%)	85,89,89	1.35	11 (12%)
3	ITR	F	349	-	15,16,16	1.02	1 (6%)	17,22,22	0.66	0
3	ITR	C	349	-	15,16,16	1.01	1 (6%)	17,22,22	0.66	0
3	ITR	E	349	-	15,16,16	1.01	1 (6%)	17,22,22	0.67	0
2	FAD	A	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.35	11 (12%)
3	ITR	B	349	-	15,16,16	1.01	1 (6%)	17,22,22	0.67	0
2	FAD	B	348	-	58,58,58	1.07	5 (8%)	85,89,89	1.35	11 (12%)
4	DTR	E	350	-	15,16,16	1.13	1 (6%)	18,22,22	0.79	1 (5%)
3	ITR	A	349	-	15,16,16	1.01	2 (13%)	17,22,22	0.67	0
3	ITR	H	349	-	15,16,16	1.01	2 (13%)	17,22,22	0.67	0
4	DTR	F	350	-	15,16,16	1.34	2 (13%)	18,22,22	1.15	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ITR	G	349	-	-	0/5/8/8	0/2/2/2
2	FAD	C	348	-	-	5/34/50/50	0/6/6/6
3	ITR	D	349	-	-	0/5/8/8	0/2/2/2
2	FAD	E	348	-	-	5/34/50/50	0/6/6/6
2	FAD	F	348	-	-	5/34/50/50	0/6/6/6
2	FAD	D	348	-	-	5/34/50/50	0/6/6/6
2	FAD	G	348	-	-	5/34/50/50	0/6/6/6
2	FAD	H	348	-	-	5/34/50/50	0/6/6/6
3	ITR	F	349	-	-	0/5/8/8	0/2/2/2
3	ITR	C	349	-	-	0/5/8/8	0/2/2/2
3	ITR	E	349	-	-	0/5/8/8	0/2/2/2
2	FAD	A	348	-	-	5/34/50/50	0/6/6/6
3	ITR	B	349	-	-	0/5/8/8	0/2/2/2
2	FAD	B	348	-	-	5/34/50/50	0/6/6/6
4	DTR	E	350	-	-	1/8/8/8	0/2/2/2
3	ITR	A	349	-	-	0/5/8/8	0/2/2/2
3	ITR	H	349	-	-	0/5/8/8	0/2/2/2
4	DTR	F	350	-	-	2/8/8/8	0/2/2/2

The worst 5 of 54 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	348	FAD	C5X-N5	-3.44	1.33	1.39
2	B	348	FAD	C5X-N5	-3.43	1.33	1.39
2	A	348	FAD	C5X-N5	-3.41	1.33	1.39
2	E	348	FAD	C5X-N5	-3.41	1.33	1.39
2	H	348	FAD	C5X-N5	-3.40	1.33	1.39

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	348	FAD	C2B-C1B-N9A	4.32	124.03	113.30
2	C	348	FAD	C2B-C1B-N9A	4.31	124.02	113.30
2	G	348	FAD	C2B-C1B-N9A	4.31	124.02	113.30
2	E	348	FAD	C2B-C1B-N9A	4.31	124.01	113.30
2	B	348	FAD	C2B-C1B-N9A	4.31	124.01	113.30

There are no chirality outliers.

5 of 43 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	348	FAD	C5B-O5B-PA-O2A
2	A	348	FAD	C5B-O5B-PA-O3P
2	B	348	FAD	C5B-O5B-PA-O2A
2	B	348	FAD	C5B-O5B-PA-O3P
2	C	348	FAD	C5B-O5B-PA-O2A


There are no ring outliers.


11 monomers are involved in 43 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	348	FAD	4	0
2	E	348	FAD	2	0
2	F	348	FAD	4	0
2	D	348	FAD	4	0
2	G	348	FAD	5	0
2	H	348	FAD	5	0
3	F	349	ITR	1	0
2	A	348	FAD	5	0
2	B	348	FAD	3	0
4	E	350	DTR	7	0
4	F	350	DTR	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	330/347 (95%)	0.48	17 (5%)	33 17	12, 34, 67, 100	23 (6%)
1	B	328/347 (94%)	0.52	22 (6%)	24 13	13, 35, 70, 100	24 (7%)
1	C	331/347 (95%)	0.45	17 (5%)	33 18	13, 35, 71, 100	26 (7%)
1	D	332/347 (95%)	0.49	24 (7%)	21 11	12, 35, 67, 100	19 (5%)
1	E	331/347 (95%)	0.41	19 (5%)	29 16	13, 35, 69, 100	26 (7%)
1	F	329/347 (94%)	0.38	13 (3%)	42 23	13, 35, 68, 100	19 (5%)
1	G	327/347 (94%)	0.67	24 (7%)	21 11	8, 35, 79, 100	26 (7%)
1	H	329/347 (94%)	0.45	19 (5%)	29 16	13, 35, 69, 100	22 (6%)
All	All	2637/2776 (94%)	0.48	155 (5%)	28 15	8, 35, 70, 100	185 (7%)

The worst 5 of 155 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	303	THR	6.5
1	D	299	GLY	6.4
1	H	221	ARG	5.7
1	G	60	SER	5.4
1	C	224	TYR	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ITR	E	349	15/15	0.77	0.21	45,52,58,72	0
3	ITR	C	349	15/15	0.82	0.23	45,52,58,72	0
3	ITR	H	349	15/15	0.82	0.23	45,52,58,72	0
3	ITR	G	349	15/15	0.83	0.20	45,52,58,72	0
3	ITR	D	349	15/15	0.84	0.20	45,52,58,72	0
3	ITR	F	349	15/15	0.85	0.20	45,52,58,72	0
3	ITR	B	349	15/15	0.89	0.16	45,52,58,72	0
4	DTR	E	350	15/15	0.89	0.15	9,42,100,100	0
2	FAD	G	348	53/53	0.91	0.14	20,27,42,43	0
3	ITR	A	349	15/15	0.91	0.16	45,52,58,72	0
2	FAD	B	348	53/53	0.93	0.12	20,27,42,43	0
4	DTR	F	350	15/15	0.93	0.17	1,49,100,100	0
2	FAD	F	348	53/53	0.94	0.12	20,27,42,43	0
2	FAD	C	348	53/53	0.94	0.12	20,27,42,43	0
2	FAD	E	348	53/53	0.94	0.11	20,27,42,43	0
2	FAD	A	348	53/53	0.95	0.10	20,27,42,43	0
2	FAD	D	348	53/53	0.96	0.10	20,27,42,43	0
2	FAD	H	348	53/53	0.96	0.09	20,27,42,43	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FAD G 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD B 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD F 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD C 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD E 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD A 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD D 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD H 348:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.