



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:59 AM UTC

PDB ID : 6DDK / pdb_00006ddk
Title : Crystal structure of the double mutant (D52N/R367Q) of the full-length NT5C2 in the basal state
Authors : Forouhar, F.; Dieck, C.L.; Tzoneva, G.; Carpenter, Z.; Ambesi-Impiombato, A.; Sanchez-Martin, M.; Kirschner-Schwabe, R.; Lew, S.; Seetharaman, J.; Ferrando, A.A.; Tong, L.
Deposited on : 2018-05-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	579	4% 75% 10% 14%
1	B	579	4% 71% 10% 17%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytosolic purine 5'-nucleotidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	500	Total	C	N	O	S	0	0	0
			4071	2627	679	744	21			
1	B	482	Total	C	N	O	S	0	0	0
			3931	2539	655	717	20			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	GLY	-	expression tag	UNP P49902
A	-16	SER	-	expression tag	UNP P49902
A	-15	SER	-	expression tag	UNP P49902
A	-14	HIS	-	expression tag	UNP P49902
A	-13	HIS	-	expression tag	UNP P49902
A	-12	HIS	-	expression tag	UNP P49902
A	-11	HIS	-	expression tag	UNP P49902
A	-10	HIS	-	expression tag	UNP P49902
A	-9	HIS	-	expression tag	UNP P49902
A	-8	SER	-	expression tag	UNP P49902
A	-7	SER	-	expression tag	UNP P49902
A	-6	GLY	-	expression tag	UNP P49902
A	-5	LEU	-	expression tag	UNP P49902
A	-4	VAL	-	expression tag	UNP P49902
A	-3	PRO	-	expression tag	UNP P49902
A	-2	ARG	-	expression tag	UNP P49902
A	-1	GLY	-	expression tag	UNP P49902
A	0	SER	-	expression tag	UNP P49902
A	52	ASN	ASP	engineered mutation	UNP P49902
A	367	GLN	ARG	engineered mutation	UNP P49902
B	-17	GLY	-	expression tag	UNP P49902
B	-16	SER	-	expression tag	UNP P49902
B	-15	SER	-	expression tag	UNP P49902
B	-14	HIS	-	expression tag	UNP P49902
B	-13	HIS	-	expression tag	UNP P49902

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	expression tag	UNP P49902
B	-11	HIS	-	expression tag	UNP P49902
B	-10	HIS	-	expression tag	UNP P49902
B	-9	HIS	-	expression tag	UNP P49902
B	-8	SER	-	expression tag	UNP P49902
B	-7	SER	-	expression tag	UNP P49902
B	-6	GLY	-	expression tag	UNP P49902
B	-5	LEU	-	expression tag	UNP P49902
B	-4	VAL	-	expression tag	UNP P49902
B	-3	PRO	-	expression tag	UNP P49902
B	-2	ARG	-	expression tag	UNP P49902
B	-1	GLY	-	expression tag	UNP P49902
B	0	SER	-	expression tag	UNP P49902
B	52	ASN	ASP	engineered mutation	UNP P49902
B	367	GLN	ARG	engineered mutation	UNP P49902

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	290	Total 290	O 290	0	0
3	B	224	Total 224	O 224	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytosolic purine 5'-nucleotidase



• Molecule 1: Cytosolic purine 5'-nucleotidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	121.85Å 171.40Å 122.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.81 – 2.50 49.81 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.81-2.50) 100.0 (49.81-2.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.25 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.162 , 0.223 0.168 , 0.226	Depositor DCC
R_{free} test set	4459 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8526	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/4177	0.53	0/5645
1	B	0.29	0/4032	0.50	0/5445
All	All	0.31	0/8209	0.52	0/11090

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	51	PHE	Peptide
1	B	51	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4071	0	3997	37	0
1	B	3931	0	3876	38	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	290	0	0	5	0
3	B	224	0	0	0	0
All	All	8526	0	7873	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:506:THR:HG22	1:B:509:ARG:H	1.52	0.74
1:A:100:LEU:HD11	1:A:437:MET:HG2	1.75	0.68
1:A:302:LEU:HD23	1:B:190:LEU:HD11	1.76	0.67
1:B:399:LEU:HD13	1:B:419:ILE:HD13	1.79	0.64
1:B:148:GLU:OE2	1:B:195:ARG:NH1	2.31	0.63
1:A:399:LEU:HD13	1:A:419:ILE:HD12	1.83	0.60
1:A:295:PHE:HZ	1:A:335:ILE:HD12	1.69	0.58
1:A:287:LEU:HD13	1:A:335:ILE:HD11	1.86	0.57
1:B:76:ARG:O	1:B:80:ILE:HG13	2.04	0.57
1:A:306:ASP:O	1:A:310:GLY:N	2.28	0.55
1:A:76:ARG:NE	1:A:80:ILE:HD11	2.23	0.54
1:B:232:LEU:HB3	1:B:233:PRO:HD3	1.90	0.54
1:B:375:LEU:HB3	1:B:463:ALA:HB2	1.90	0.52
1:A:195:ARG:NH2	1:A:199:GLN:OE1	2.43	0.52
1:A:190:LEU:HD21	1:B:302:LEU:HD23	1.91	0.52
1:A:25:LYS:H	1:A:25:LYS:HE2	1.75	0.52
1:A:303:ARG:HG3	1:A:312:LEU:HD22	1.92	0.51
1:A:429:ASP:OD2	3:A:701:HOH:O	2.20	0.50
1:B:296:PHE:HB3	1:B:358:LEU:HD13	1.93	0.50
1:B:290:ALA:O	1:B:295:PHE:HB2	2.13	0.49
1:A:395:LEU:HD22	1:A:419:ILE:HG12	1.95	0.49
1:A:422:ARG:NH1	1:A:425:LYS:HZ2	2.11	0.49
1:A:311:LYS:HD2	1:A:317:TYR:OH	2.13	0.48
1:B:288:VAL:O	1:B:328:SER:HA	2.13	0.48
1:A:500:MET:HE2	3:A:810:HOH:O	2.14	0.47
1:A:157:PHE:O	1:A:160:PRO:HD2	2.14	0.47
1:A:288:VAL:O	1:A:328:SER:HA	2.14	0.46
1:A:466:ILE:HA	1:A:466:ILE:HD12	1.83	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:466:ILE:HD12	1:B:466:ILE:HA	1.81	0.46
1:B:51:PHE:O	1:B:247:LEU:HD12	2.16	0.46
1:B:309:THR:HB	1:B:311:LYS:HG2	1.98	0.46
1:A:303:ARG:HB3	1:A:314:ILE:HA	1.97	0.45
1:B:157:PHE:O	1:B:160:PRO:HD2	2.17	0.45
1:B:423:ILE:O	1:B:427:THR:HG23	2.17	0.44
1:A:232:LEU:HB3	1:A:233:PRO:HD3	1.98	0.44
1:A:61:LYS:HG2	1:A:227:VAL:HG23	2.00	0.44
1:A:76:ARG:O	1:A:80:ILE:HG13	2.17	0.44
1:A:130:GLY:O	1:A:133:THR:HB	2.19	0.43
1:A:268:HIS:HB3	3:A:848:HOH:O	2.17	0.43
1:A:161:GLU:OE2	1:A:202:ARG:NH1	2.52	0.43
1:B:338:LEU:HD23	1:B:338:LEU:HA	1.86	0.43
1:B:304:GLN:NE2	1:B:316:THR:HG23	2.34	0.43
1:B:308:LYS:H	1:B:308:LYS:HG3	1.58	0.43
1:A:86:LEU:HG	1:A:89:PHE:CE2	2.54	0.42
1:B:86:LEU:HD21	1:B:168:LEU:HD21	2.02	0.42
1:B:219:VAL:HG13	1:B:258:LYS:HG2	2.01	0.42
1:A:132:GLU:H	1:A:132:GLU:HG3	1.71	0.42
1:B:313:LYS:HD3	1:B:317:TYR:HA	2.00	0.42
1:B:486:HIS:O	1:B:490:VAL:HG13	2.20	0.42
1:A:230:GLY:O	1:A:233:PRO:HD2	2.20	0.42
1:B:207:TRP:O	1:B:211:LYS:HB2	2.20	0.42
1:B:175:CYS:HB3	1:B:178:TYR:HD2	1.85	0.41
1:B:76:ARG:HD3	1:B:207:TRP:CE3	2.55	0.41
1:A:161:GLU:OE1	1:A:202:ARG:NH1	2.53	0.41
1:B:295:PHE:CZ	1:B:335:ILE:HD12	2.56	0.41
1:A:359:LYS:HB3	1:A:359:LYS:HE3	1.84	0.41
1:B:80:ILE:HG21	1:B:80:ILE:HD13	1.89	0.41
1:B:207:TRP:CD1	1:B:211:LYS:HE3	2.55	0.41
1:B:130:GLY:O	1:B:133:THR:HB	2.20	0.41
1:A:72:LEU:HD23	1:A:72:LEU:HA	1.87	0.41
1:A:175:CYS:SG	1:A:177:ARG:HG3	2.61	0.41
1:B:202:ARG:HH11	1:B:202:ARG:HG3	1.86	0.41
1:A:302:LEU:HD21	1:A:338:LEU:HD11	2.02	0.41
1:B:76:ARG:NE	1:B:80:ILE:HD11	2.36	0.41
1:B:211:LYS:HD3	1:B:211:LYS:HA	1.87	0.41
1:B:145:ASP:O	1:B:147:THR:N	2.54	0.40
1:A:235:LEU:HB2	1:A:469:LEU:HD13	2.04	0.40
1:B:76:ARG:HD3	1:B:207:TRP:CD2	2.57	0.40
1:B:294:LEU:HD13	1:B:294:LEU:HA	1.89	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:372:ILE:O	1:B:463:ALA:HA	2.21	0.40
1:A:39:ARG:HD3	3:A:775:HOH:O	2.21	0.40
1:A:425:LYS:HD2	3:A:701:HOH:O	2.22	0.40
1:B:222:LEU:HD13	1:B:262:TYR:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	496/579 (86%)	485 (98%)	11 (2%)	0	100	100
1	B	478/579 (83%)	466 (98%)	12 (2%)	0	100	100
All	All	974/1158 (84%)	951 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/520 (86%)	419 (94%)	26 (6%)	18	38
1	B	430/520 (83%)	401 (93%)	29 (7%)	15	31
All	All	875/1040 (84%)	820 (94%)	55 (6%)	16	34

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LYS
1	A	144	ARG
1	A	147	THR
1	A	190	LEU
1	A	195	ARG
1	A	202	ARG
1	A	213	SER
1	A	242	VAL
1	A	254	LYS
1	A	271	LYS
1	A	287	LEU
1	A	294	LEU
1	A	303	ARG
1	A	311	LYS
1	A	335	ILE
1	A	357	ILE
1	A	358	LEU
1	A	363	ARG
1	A	377	GLN
1	A	417	SER
1	A	419	ILE
1	A	422	ARG
1	A	425	LYS
1	A	446	ARG
1	A	455	MET
1	A	544	ILE
1	B	25	LYS
1	B	32	TYR
1	B	79	SER
1	B	80	ILE
1	B	147	THR
1	B	161	GLU
1	B	195	ARG
1	B	211	LYS
1	B	213	SER
1	B	254	LYS
1	B	258	LYS
1	B	294	LEU
1	B	303	ARG
1	B	308	LYS
1	B	311	LYS
1	B	322	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	361	LYS
1	B	375	LEU
1	B	377	GLN
1	B	387	SER
1	B	406	LEU
1	B	417	SER
1	B	419	ILE
1	B	446	ARG
1	B	455	MET
1	B	466	ILE
1	B	469	LEU
1	B	490	VAL
1	B	508	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	420	GLN
1	A	481	HIS
1	A	548	HIS
1	B	33	HIS
1	B	52	ASN
1	B	352	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	601	-	4,4,4	0.97	0	6,6,6	0.72	0
2	PO4	B	601	-	4,4,4	1.04	0	6,6,6	0.90	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	500/579 (86%)	-0.38	21 (4%) 40 36	5, 28, 69, 99	2 (0%)
1	B	482/579 (83%)	-0.19	21 (4%) 39 34	10, 35, 68, 124	3 (0%)
All	All	982/1158 (84%)	-0.28	42 (4%) 40 35	5, 32, 69, 124	5 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	495	VAL	6.2
1	B	320	PRO	5.2
1	A	358	LEU	5.1
1	B	298	GLU	4.7
1	B	318	THR	4.5
1	A	550	GLU	4.2
1	B	380	HIS	4.1
1	A	189	ASP	4.1
1	A	549	ASP	4.0
1	A	497	ILE	3.8
1	B	357	ILE	3.8
1	A	357	ILE	3.7
1	B	317	TYR	3.5
1	B	321	LEU	3.5
1	B	189	ASP	3.5
1	A	344	LYS	3.4
1	A	24	LEU	3.3
1	B	24	LEU	3.2
1	A	319	GLY	3.1
1	A	318	THR	3.0
1	A	210	TYR	3.0
1	B	358	LEU	3.0
1	A	539	LEU	2.9
1	B	324	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	421	ARG	2.9
1	A	500	MET	2.9
1	B	319	GLY	2.8
1	A	537	PHE	2.7
1	B	500	MET	2.7
1	A	320	PRO	2.5
1	A	298	GLU	2.5
1	A	356	ASP	2.5
1	B	80	ILE	2.4
1	B	323	HIS	2.3
1	B	360	SER	2.3
1	B	494	HIS	2.2
1	A	540	ALA	2.1
1	A	364	GLN	2.1
1	A	543	GLU	2.1
1	A	538	PRO	2.1
1	B	322	GLN	2.0
1	B	258	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	601	5/5	0.99	0.04	10,11,16,17	0
2	PO4	B	601	5/5	0.99	0.04	21,25,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.