



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:14 PM UTC

PDB ID : 7DDW / pdb\_00007ddw  
Title : Crystal structure of a mutant Staphylococcus equorum manganese superoxide dismutase S126C  
Authors : Retnoningrum, D.S.; Yoshida, H.; Razani, M.D.; Meidianto, V.F.; Hartanto, A.; Artarini, A.; Ismaya, W.T.  
Deposited on : 2020-10-30  
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	 86% 10% ..
1	B	205	 86% 9% ..
1	C	205	 89% 6% ..
1	D	205	 87% 9% .
1	E	205	 82% 12% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	205	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into segments: a small red segment at the start, followed by a large green segment labeled '80%', then a yellow segment labeled '15%', and finally a small grey segment at the end. Above the bar is a '%' symbol, and below the bar are two dots '••'.</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10584 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1599	1019	270	307	3	0	1	0
1	B	197	1590	1013	268	306	3	0	1	0
1	C	198	1599	1019	270	307	3	0	1	0
1	D	198	1599	1019	270	307	3	0	1	0
1	E	198	1599	1019	270	307	3	0	1	0
1	F	198	1599	1019	270	307	3	0	1	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85
A	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
A	200	HIS	-	expression tag	UNP A0A1E5TT85
A	201	HIS	-	expression tag	UNP A0A1E5TT85
A	202	HIS	-	expression tag	UNP A0A1E5TT85
A	203	HIS	-	expression tag	UNP A0A1E5TT85
A	204	HIS	-	expression tag	UNP A0A1E5TT85
A	205	HIS	-	expression tag	UNP A0A1E5TT85
B	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85
B	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
B	200	HIS	-	expression tag	UNP A0A1E5TT85
B	201	HIS	-	expression tag	UNP A0A1E5TT85
B	202	HIS	-	expression tag	UNP A0A1E5TT85
B	203	HIS	-	expression tag	UNP A0A1E5TT85
B	204	HIS	-	expression tag	UNP A0A1E5TT85
B	205	HIS	-	expression tag	UNP A0A1E5TT85
C	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
C	200	HIS	-	expression tag	UNP A0A1E5TT85
C	201	HIS	-	expression tag	UNP A0A1E5TT85
C	202	HIS	-	expression tag	UNP A0A1E5TT85
C	203	HIS	-	expression tag	UNP A0A1E5TT85
C	204	HIS	-	expression tag	UNP A0A1E5TT85
C	205	HIS	-	expression tag	UNP A0A1E5TT85
D	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85
D	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
D	200	HIS	-	expression tag	UNP A0A1E5TT85
D	201	HIS	-	expression tag	UNP A0A1E5TT85
D	202	HIS	-	expression tag	UNP A0A1E5TT85
D	203	HIS	-	expression tag	UNP A0A1E5TT85
D	204	HIS	-	expression tag	UNP A0A1E5TT85
D	205	HIS	-	expression tag	UNP A0A1E5TT85
E	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85
E	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
E	200	HIS	-	expression tag	UNP A0A1E5TT85
E	201	HIS	-	expression tag	UNP A0A1E5TT85
E	202	HIS	-	expression tag	UNP A0A1E5TT85
E	203	HIS	-	expression tag	UNP A0A1E5TT85
E	204	HIS	-	expression tag	UNP A0A1E5TT85
E	205	HIS	-	expression tag	UNP A0A1E5TT85
F	13	ARG	ASP	engineered mutation	UNP A0A1E5TT85
F	126	CYS	SER	engineered mutation	UNP A0A1E5TT85
F	200	HIS	-	expression tag	UNP A0A1E5TT85
F	201	HIS	-	expression tag	UNP A0A1E5TT85
F	202	HIS	-	expression tag	UNP A0A1E5TT85
F	203	HIS	-	expression tag	UNP A0A1E5TT85
F	204	HIS	-	expression tag	UNP A0A1E5TT85
F	205	HIS	-	expression tag	UNP A0A1E5TT85

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mn 1 1	0	0
2	B	1	Total Mn 1 1	0	0
2	C	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	1	Total 1	Mn 1	0	0
2	F	1	Total 1	Mn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	178	Total 178	O 178	0	0
3	B	146	Total 146	O 146	0	0
3	C	195	Total 195	O 195	0	0
3	D	204	Total 204	O 204	0	0
3	E	130	Total 130	O 130	0	0
3	F	140	Total 140	O 140	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Superoxide dismutase

Chain A:  86% 10% ..



- Molecule 1: Superoxide dismutase

Chain B:  86% 9% ..




- Molecule 1: Superoxide dismutase

Chain C:  89% 6% ..





- Molecule 1: Superoxide dismutase

Chain D:  87% 9% .





- Molecule 1: Superoxide dismutase

Chain E:  82% 12% ..



## ● Molecule 1: Superoxide dismutase

Chain F:  %

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.29Å 131.73Å 80.19Å 90.00° 110.35° 90.00°	Depositor
Resolution (Å)	42.44 – 1.88 42.44 – 1.88	Depositor EDS
% Data completeness (in resolution range)	97.1 (42.44-1.88) 97.1 (42.44-1.88)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.41 (at 1.88Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.157 , 0.222 0.166 , 0.228	Depositor DCC
$R_{free}$ test set	5222 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.7	Xtrriage
Anisotropy	1.413	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10584	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.98	0/1646	1.33	0/2244
1	B	0.96	0/1637	1.36	2/2233 (0.1%)
1	C	0.96	0/1646	1.33	0/2244
1	D	0.98	0/1646	1.32	0/2244
1	E	0.95	0/1646	1.32	0/2244
1	F	0.95	0/1646	1.35	2/2244 (0.1%)
All	All	0.96	0/9867	1.33	4/13453 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	ASP	CA-C-N	5.65	128.17	120.54
1	B	20	ASP	C-N-CA	5.65	128.17	120.54
1	F	20	ASP	CA-C-N	5.16	127.50	120.54
1	F	20	ASP	C-N-CA	5.16	127.50	120.54

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1599	0	1520	14	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1590	0	1507	14	0
1	C	1599	0	1520	13	0
1	D	1599	0	1520	12	0
1	E	1599	0	1520	22	0
1	F	1599	0	1520	28	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	178	0	0	2	0
3	B	146	0	0	2	0
3	C	195	0	0	3	0
3	D	204	0	0	1	0
3	E	130	0	0	2	0
3	F	140	0	0	2	0
All	All	10584	0	9107	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:17:PRO:O	1:E:170:LYS:HE2	1.46	1.15
1:C:118:ASN:HB2	3:C:454:HOH:O	1.59	1.02
1:B:47:ASP:CG	1:F:118:ASN:HD22	1.77	0.93
1:F:17:PRO:O	1:F:170:LYS:HE2	1.68	0.92
1:B:17:PRO:O	1:B:170:LYS:HE2	1.73	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/205 (96%)	192 (98%)	5 (2%)	0	100	100
1	B	196/205 (96%)	191 (97%)	5 (3%)	0	100	100
1	C	197/205 (96%)	192 (98%)	5 (2%)	0	100	100
1	D	197/205 (96%)	193 (98%)	4 (2%)	0	100	100
1	E	197/205 (96%)	191 (97%)	6 (3%)	0	100	100
1	F	197/205 (96%)	189 (96%)	8 (4%)	0	100	100
All	All	1181/1230 (96%)	1148 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/177 (97%)	166 (97%)	5 (3%)	37	20
1	B	170/177 (96%)	167 (98%)	3 (2%)	51	38
1	C	171/177 (97%)	168 (98%)	3 (2%)	51	38
1	D	171/177 (97%)	166 (97%)	5 (3%)	37	20
1	E	171/177 (97%)	163 (95%)	8 (5%)	23	8
1	F	171/177 (97%)	166 (97%)	5 (3%)	37	20
All	All	1025/1062 (96%)	996 (97%)	29 (3%)	38	22

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	137	LYS
1	F	101	LYS
1	E	61	ASP
1	F	13	ARG
1	E	55	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	33	ASN
1	D	119	GLN
1	F	118	ASN
1	F	21	GLN
1	F	74	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/205 (96%)	-0.39	1 (0%) 87 90	14, 22, 49, 70	1 (0%)
1	B	197/205 (96%)	-0.22	2 (1%) 79 83	16, 29, 52, 70	1 (0%)
1	C	198/205 (96%)	-0.39	1 (0%) 87 90	14, 21, 45, 73	1 (0%)
1	D	198/205 (96%)	-0.44	1 (0%) 87 90	12, 21, 43, 64	1 (0%)
1	E	198/205 (96%)	-0.23	4 (2%) 65 71	17, 29, 51, 72	1 (0%)
1	F	198/205 (96%)	-0.25	3 (1%) 72 77	18, 31, 54, 75	1 (0%)
All	All	1187/1230 (96%)	-0.32	12 (1%) 79 83	12, 26, 51, 75	6 (0%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	3.0
1	B	45	GLY	2.8
1	A	199	LYS	2.6
1	D	199	LYS	2.5
1	C	199	LYS	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	A	301	1/1	0.99	0.01	16,16,16,16	0
2	MN	D	301	1/1	0.99	0.02	13,13,13,13	0
2	MN	C	301	1/1	1.00	0.03	20,20,20,20	0
2	MN	B	301	1/1	1.00	0.01	19,19,19,19	0
2	MN	E	301	1/1	1.00	0.01	22,22,22,22	0
2	MN	F	301	1/1	1.00	0.01	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.