



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:36 PM UTC

PDB ID : 4DEL / pdb_00004del
Title : Active site loop dynamics of a class IIa fructose 1,6-bisphosphate aldolase from M. tuberculosis
Authors : Pegan, S.D.; Mesecar, A.D.
Deposited on : 2012-01-20
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	13% 87% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	405	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	7	21	0
			2738	1741	465	522	10			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	345	HIS	-	expression tag	UNP P67475
A	346	HIS	-	expression tag	UNP P67475
A	347	HIS	-	expression tag	UNP P67475
A	348	HIS	-	expression tag	UNP P67475
A	349	HIS	-	expression tag	UNP P67475


- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is PHOSPHOGLYCOLOHYDROXAMIC ACID (CCD ID: PGH) (formula: C₂H₆NO₆P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			10	2	1	6	1		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is HEXAETHYLENE GLYCOL (CCD ID: P6G) (formula: $C_{12}H_{26}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			19	12	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	513	Total	O	0	0
			513	513		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructose-bisphosphate aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	61.80Å 119.61Å 164.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.74 – 1.58 49.74 – 1.58	Depositor EDS
% Data completeness (in resolution range)	93.9 (49.74-1.58) 93.9 (49.74-1.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.20 (at 1.58Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.189 , 0.205 0.186 , 0.202	Depositor DCC
R_{free} test set	3910 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.999	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 70.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3287	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ACT, PGH, ZN, P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/2817	0.77	0/3817

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2738	0	2718	35	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	10	0	3	0	0
5	A	4	0	3	4	0
6	A	19	0	26	4	0
7	A	513	0	0	8	0
All	All	3287	0	2750	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:405:ACT:H1	7:A:866:HOH:O	1.67	0.93
1:A:87[A]:PRO:O	1:A:343:THR:HG21	1.70	0.92
1:A:344[A]:HIS:CE1	7:A:992:HOH:O	2.23	0.89
1:A:231:GLN:HE22	1:A:247:PHE:H	1.26	0.83
1:A:263:GLU:OE2	1:A:336:HIS:HE1	1.75	0.70
1:A:92:LEU:H	1:A:126:GLN:NE2	1.90	0.70
6:A:406:P6G:H81	7:A:960:HOH:O	1.94	0.68
1:A:14:GLY:HA2	5:A:405:ACT:H3	1.76	0.67
1:A:92:LEU:H	1:A:126:GLN:HE21	1.44	0.65
1:A:290:HIS:HD2	1:A:319:LYS:NZ	1.95	0.63
1:A:231:GLN:HE21	1:A:231:GLN:HA	1.66	0.60
1:A:272:LYS:NZ	1:A:274:ASN:HD21	2.01	0.57
1:A:211[B]:VAL:HG11	1:A:220:VAL:HG22	1.87	0.57
1:A:10:ALA:HB1	6:A:406:P6G:H61	1.86	0.57
5:A:405:ACT:H2	6:A:406:P6G:H121	1.89	0.55
1:A:26:ILE:HD13	1:A:275[A]:VAL:HG13	1.88	0.55
1:A:290:HIS:HD2	1:A:319:LYS:HZ2	1.55	0.54
1:A:164:VAL:O	1:A:207[B]:THR:HG21	2.09	0.52
1:A:166:GLY:C	1:A:211[B]:VAL:HG23	2.35	0.52
1:A:290:HIS:HE1	1:A:310:VAL:O	1.92	0.52
1:A:211[B]:VAL:CG1	1:A:220:VAL:HG21	2.39	0.52
1:A:211[B]:VAL:HG11	1:A:220:VAL:CG2	2.39	0.52
1:A:26:ILE:HD13	1:A:275[A]:VAL:CG1	2.40	0.50
1:A:211[B]:VAL:CG1	1:A:220:VAL:CG2	2.90	0.50
1:A:263:GLU:OE2	1:A:336:HIS:CE1	2.63	0.49
1:A:26:ILE:CD1	1:A:275[A]:VAL:HG13	2.43	0.48
1:A:38:ILE:HG13	1:A:50[A]:ILE:HD11	1.96	0.48
1:A:80:HIS:HE1	7:A:587:HOH:O	1.98	0.47
1:A:319:LYS:NZ	7:A:949:HOH:O	2.47	0.46
1:A:15:GLN:HG2	7:A:997:HOH:O	2.16	0.46
1:A:181:TYR:HD1	1:A:226[B]:ILE:CD1	2.30	0.45
1:A:272:LYS:HZ3	1:A:274:ASN:HD21	1.63	0.44
1:A:14:GLY:HA2	5:A:405:ACT:CH3	2.46	0.43
1:A:290:HIS:CG	1:A:316:TYR:HB3	2.55	0.42
1:A:24[B]:PRO:HD3	1:A:337:CYS:SG	2.59	0.42
1:A:313:PRO:HA	1:A:316:TYR:CE2	2.55	0.42
6:A:406:P6G:H31	6:A:406:P6G:H62	1.81	0.41
1:A:219:ASN:ND2	7:A:1012:HOH:O	2.54	0.41
1:A:309[A]:LYS:HG3	7:A:862:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	364/349 (104%)	357 (98%)	6 (2%)	1 (0%)	36 20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	96	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/271 (106%)	282 (98%)	6 (2%)	47 17

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8[A]	VAL
1	A	8[B]	VAL
1	A	170	ASP
1	A	231	GLN
1	A	240	LEU
1	A	259	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	GLN
1	A	219	ASN
1	A	231	GLN
1	A	274	ASN
1	A	280	GLN
1	A	290	HIS
1	A	294	ASN
1	A	336	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	P6G	A	406	-	18,18,18	0.43	0	17,17,17	0.41	0
5	ACT	A	405	-	3,3,3	0.82	0	3,3,3	0.87	0
4	PGH	A	404	3,2	9,9,9	2.61	1 (11%)	10,12,12	0.95	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	P6G	A	406	-	-	6/16/16/16	-
4	PGH	A	404	3,2	-	1/8/8/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	PGH	O1-C1	7.53	1.38	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	PGH	C2-C1-N2	2.04	119.92	116.41

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	406	P6G	C6-C5-O4-C3
6	A	406	P6G	C8-C9-O10-C11
6	A	406	P6G	O4-C5-C6-O7
4	A	404	PGH	O1-C1-C2-O1P
6	A	406	P6G	C9-C8-O7-C6
6	A	406	P6G	O1-C2-C3-O4
6	A	406	P6G	C5-C6-O7-C8


There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	406	P6G	4	0
5	A	405	ACT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	8[A]:VAL	C	9:TYR	N	1.04

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/349 (98%)	0.56	46 (13%) 7 12	6, 18, 36, 54	20 (5%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	GLY	9.4
1	A	172	VAL	6.8
1	A	173	ALA	6.4
1	A	176	ILE	6.4
1	A	178	GLU	5.8
1	A	170	ASP	5.8
1	A	345[A]	HIS	4.8
1	A	223	ARG	3.9
1	A	169	GLU	3.9
1	A	233	VAL	3.8
1	A	225	ASP	3.6
1	A	185[A]	GLU	3.5
1	A	238	LEU	3.4
1	A	193	ALA	3.4
1	A	167	GLY	3.3
1	A	174	ASN	3.3
1	A	218	GLY	3.2
1	A	217	PRO	3.2
1	A	240	LEU	3.2
1	A	191	ILE	2.9
1	A	184	PRO	2.9
1	A	236	ALA	2.9
1	A	229	GLN	2.9
1	A	242	ALA	2.8
1	A	180	LEU	2.7
1	A	239	GLY	2.7
1	A	235	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	346	HIS	2.5
1	A	219	ASN	2.5
1	A	181	TYR	2.5
1	A	241	PRO	2.4
1	A	175	GLU	2.4
1	A	183	SER	2.3
1	A	182	THR	2.3
1	A	177	ASN	2.3
1	A	117	VAL	2.2
1	A	164	VAL	2.2
1	A	119	LYS	2.2
1	A	186	ASP	2.2
1	A	134	ALA	2.1
1	A	137	ILE	2.1
1	A	118	SER	2.1
1	A	194	LEU	2.1
1	A	188[A]	GLU	2.1
1	A	179	LYS	2.1
1	A	168	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ACT	A	405	4/4	0.80	0.20	21,23,23,23	0
6	P6G	A	406	19/19	0.90	0.12	20,29,33,35	0
3	ZN	A	403	1/1	0.96	0.06	28,28,28,28	0
2	NA	A	401	1/1	0.98	0.04	15,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	402	1/1	0.99	0.03	16,16,16,16	0
4	PGH	A	404	10/10	0.99	0.04	12,15,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.