



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 07:15 AM UTC

PDB ID : 1DFK / pdb\_00001dfk  
Title : NUCLEOTIDE-FREE SCALLOP MYOSIN S1-NEAR RIGOR STATE  
Authors : Houdusse, A.; Szent-Gyorgyi, A.G.; Cohen, C.  
Deposited on : 1999-11-19  
Resolution : 4.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1019 (4.50-3.90)
Ramachandran outliers	187476	1020 (4.56-3.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	830	
2	Y	139	
3	Z	152	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	728	3597	2140	728	729	0	0	0

- Molecule 2 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	Y	139	688	409	139	140	0	0	0

- Molecule 3 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	Z	152	746	441	152	153	0	0	0

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Z	1	Total	Ca	0	0
			1	1		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: MYOSIN HEAD




- Molecule 2: MYOSIN HEAD



- Molecule 3: MYOSIN HEAD





## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.50Å 52.10Å 164.60Å 90.00° 100.30° 90.00°	Depositor
Resolution (Å)	20.00 – 4.20	Depositor
% Data completeness (in resolution range)	95.0 (20.00-4.20)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.415 , 0.410	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5032	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.47	10/3587 (0.3%)	1.92	68/4978 (1.4%)
2	Y	0.69	1/687 (0.1%)	1.38	12/954 (1.3%)
3	Z	1.01	1/745 (0.1%)	1.34	11/1031 (1.1%)
All	All	1.33	12/5019 (0.2%)	1.78	91/6963 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	268	GLU	C-N	-51.10	0.62	1.33
1	A	683	ASP	C-N	-34.01	0.91	1.33
1	A	705	LYS	C-N	28.13	1.74	1.33
1	A	706	GLY	C-N	24.14	1.79	1.33
3	Z	7	GLU	C-N	21.68	1.59	1.33

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	515	ASP	O-C-N	-47.20	59.82	122.59
1	A	268	GLU	O-C-N	-42.60	65.94	122.59
1	A	268	GLU	CA-C-N	-32.25	59.95	121.54
1	A	268	GLU	C-N-CA	-32.25	59.95	121.54
1	A	600	ASP	O-C-N	-20.13	98.17	121.32

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	268	GLU	Mainchain
1	A	445	THR	Mainchain
1	A	515	ASP	Peptide,Mainchain
1	A	600	ASP	Mainchain
1	A	601	PRO	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3597	0	1592	71	0
2	Y	688	0	307	21	0
3	Z	746	0	340	17	0
4	Z	1	0	0	0	0
All	All	5032	0	2239	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:705:LYS:C	1:A:706:GLY:N	1.74	1.44
1:A:706:GLY:C	1:A:707:PHE:N	1.80	1.40
1:A:584:ALA:HB1	1:A:693:CYS:HA	1.37	1.07
1:A:770:LEU:O	1:A:774:ARG:CB	2.10	0.99
1:A:507:TRP:O	1:A:508:GLU:C	2.07	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	708/830 (85%)	576 (81%)	95 (13%)	37 (5%)	1	16
2	Y	137/139 (99%)	96 (70%)	34 (25%)	7 (5%)	1	16
3	Z	150/152 (99%)	106 (71%)	33 (22%)	11 (7%)	1	11
All	All	995/1121 (89%)	778 (78%)	162 (16%)	55 (6%)	1	15

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	GLU
1	A	269	LYS
1	A	542	ALA
1	A	601	PRO
1	A	623	LYS

### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	6

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	706:GLY	C	707:PHE	N	1.80
1	A	705:LYS	C	706:GLY	N	1.74
1	A	489:ASN	C	490:HIS	N	1.20
1	A	445:THR	C	446:LEU	N	1.10
1	A	683:ASP	C	684:ALA	N	0.91

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.