



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 03:19 PM UTC

PDB ID : 2DFP / pdb\_00002dfp  
Title : X-RAY STRUCTURE OF AGED DI-ISOPROPYL-PHOSPHORO-FLUORIDE (DFP) BOUND TO ACETYLCHOLINESTERASE  
Authors : Kryger, G.; Millard, C.B.; Silman, I.; Sussman, J.L.  
Deposited on : 1998-12-07  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	
2	B	2	
2	C	2	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ACETYLCHOLINESTERASE).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	534	4270	2734	724	789	1	22	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0
2	C	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	12	6	1	4	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	376	Total 376	O 376	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (ACETYLCHOLINESTERASE)



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.64Å 112.64Å 136.88Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.7 (20.00-2.30) 93.6 (20.00-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.21Å)	Xtrriage
Refinement program	CNS 0.4	Depositor
R, $R_{free}$	0.186 , 0.228 0.190 , 0.230	Depositor DCC
$R_{free}$ test set	2015 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.0	Xtrriage
Anisotropy	0.563	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4728	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MIS, NAG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.05	3/4380 (0.1%)	1.25	32/5944 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	520	MET	SD-CE	7.98	1.99	1.79
1	A	353	MET	SD-CE	5.25	1.92	1.79
1	A	310	ASN	CA-C	-5.09	1.46	1.52

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	518	VAL	CB-CA-C	-9.20	100.20	111.97
1	A	443	GLU	N-CA-C	-8.38	103.08	113.38
1	A	504	ASP	N-CA-C	-8.27	98.28	110.52
1	A	133	LYS	N-CA-C	7.27	119.20	111.28
1	A	329	SER	N-CA-C	7.16	119.09	111.28
1	A	439	ILE	N-CA-C	7.07	119.80	109.63
1	A	142	VAL	N-CA-C	-6.78	99.94	108.89
1	A	234	ALA	N-CA-C	6.64	121.38	113.28
1	A	347	ILE	N-CA-C	6.46	117.16	107.80
1	A	5	GLU	N-CA-C	6.31	118.97	111.71

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	527	PHE	N-CA-C	6.29	118.02	111.03
1	A	401	ILE	N-CA-C	6.06	116.18	110.30
1	A	297	ASP	N-CA-C	6.03	120.79	113.38
1	A	157	ALA	N-CA-C	6.00	118.70	108.02
1	A	222	ALA	N-CA-C	5.79	118.20	109.23
1	A	101	VAL	N-CA-C	5.77	114.48	107.61
1	A	283	PRO	N-CA-C	5.73	121.51	114.35
1	A	346	LYS	N-CA-C	-5.69	101.42	109.96
1	A	22	VAL	CB-CA-C	-5.61	102.10	111.29
1	A	360	VAL	CA-C-N	5.56	125.92	119.47
1	A	360	VAL	C-N-CA	5.56	125.92	119.47
1	A	55	SER	N-CA-C	-5.51	102.03	110.14
1	A	270	LYS	N-CA-C	-5.50	102.28	110.20
1	A	310	ASN	N-CA-C	5.41	117.17	111.28
1	A	228	SER	CB-CA-C	-5.37	100.06	108.91
1	A	442	TYR	N-CA-C	5.26	118.83	112.93
1	A	167	ASN	N-CA-C	5.25	119.27	112.86
1	A	292	PHE	N-CA-C	5.18	117.38	107.75
1	A	308	MET	N-CA-C	-5.10	105.62	111.07
1	A	255	ASN	N-CA-C	-5.07	102.14	109.59
1	A	328	GLY	N-CA-C	5.05	121.94	115.32
1	A	278	GLU	N-CA-C	5.04	118.53	112.38

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	442	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4270	0	4111	74	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
3	A	14	0	13	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	12	0	13	0	0
5	A	376	0	0	9	0
All	All	4728	0	4187	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:VAL:HG13	1:A:133:LYS:HG3	1.63	0.79
1:A:468:ARG:HD3	5:A:2327:HOH:O	1.84	0.78
1:A:487:SER:OG	1:A:489:GLU:HG2	1.90	0.71
1:A:485:PRO:HA	5:A:2363:HOH:O	1.92	0.70
1:A:366:LEU:HD23	1:A:535:THR:HG21	1.77	0.65
1:A:223:ILE:HA	1:A:320:LEU:O	1.98	0.64
1:A:275:ILE:HA	1:A:278:GLU:OE1	1.99	0.63
1:A:345:SER:O	1:A:388:ARG:HG3	2.01	0.60
1:A:426:ARG:HD3	5:A:2085:HOH:O	2.02	0.60
1:A:5:GLU:OE2	1:A:104:PRO:HA	2.02	0.59
1:A:419:TYR:CZ	1:A:494:LEU:HD13	2.37	0.59
1:A:452:LEU:HD13	1:A:467:ARG:NH2	2.18	0.58
1:A:461:GLU:H	1:A:461:GLU:CD	2.10	0.58
1:A:515:ARG:O	1:A:518:VAL:HG22	2.05	0.57
1:A:197:PHE:CB	1:A:223:ILE:HB	2.34	0.57
1:A:451:PRO:HA	1:A:458:TYR:CD2	2.41	0.55
1:A:63:TYR:CD1	1:A:126:THR:HG22	2.43	0.54
1:A:366:LEU:CD2	1:A:535:THR:HG21	2.38	0.54
1:A:127:LEU:HD12	1:A:130:TYR:CE2	2.44	0.53
1:A:349:ARG:HD3	1:A:381:ASP:O	2.09	0.52
1:A:452:LEU:HD22	1:A:463:GLU:HG3	1.91	0.52
1:A:200:MIS:H33	1:A:233:TRP:CE2	2.45	0.52
1:A:23:LEU:HD11	1:A:452:LEU:HD12	1.93	0.50
1:A:204:ALA:O	1:A:208:MET:HG3	2.11	0.50
1:A:216:ARG:HB3	1:A:315:LYS:HB2	1.93	0.49
1:A:22:VAL:O	1:A:23:LEU:C	2.56	0.49
1:A:212:SER:O	1:A:216:ARG:HG2	2.13	0.49
1:A:405:MET:HA	1:A:408:VAL:HG12	1.94	0.48
1:A:132:GLY:HA3	1:A:143:LEU:HD22	1.94	0.48
1:A:244:ARG:HD3	1:A:291:SER:HB3	1.95	0.48
1:A:197:PHE:HB3	1:A:223:ILE:HB	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ARG:CD	5:A:2264:HOH:O	2.62	0.47
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.96	0.47
1:A:528:LEU:HB3	1:A:529:PRO:HD3	1.95	0.47
1:A:19:ARG:NH2	1:A:26:HIS:HB2	2.30	0.47
1:A:293:VAL:HB	1:A:294:PRO:HD2	1.96	0.47
1:A:454:LYS:HG3	1:A:455:GLU:N	2.30	0.47
1:A:308:MET:HE2	1:A:313:ASN:O	2.15	0.47
1:A:146:LEU:C	1:A:146:LEU:HD12	2.40	0.46
1:A:484:GLU:HB2	1:A:487:SER:HB3	1.97	0.46
1:A:24:SER:HA	5:A:2179:HOH:O	2.16	0.46
1:A:199:GLU:HA	1:A:225:GLN:O	2.16	0.45
1:A:211:LEU:HD23	1:A:314:PHE:HB3	1.99	0.45
1:A:452:LEU:HD13	1:A:467:ARG:CZ	2.45	0.45
1:A:465:LEU:HD13	1:A:506:ASN:O	2.17	0.45
1:A:88:ARG:HD2	5:A:2264:HOH:O	2.16	0.45
1:A:481:ASN:OD1	1:A:482:PRO:HD2	2.17	0.45
1:A:487:SER:OG	1:A:488:GLN:N	2.51	0.44
1:A:450:LEU:HD23	1:A:450:LEU:N	2.33	0.43
1:A:508:GLU:HB3	1:A:509:PRO:HD2	1.99	0.43
1:A:82:GLU:HG2	5:A:2204:HOH:O	2.18	0.43
1:A:265:CYS:O	1:A:268:GLU:HB2	2.18	0.43
1:A:527:PHE:O	1:A:528:LEU:C	2.61	0.43
1:A:135:LEU:HD23	1:A:143:LEU:CD1	2.49	0.43
1:A:357:LYS:HG2	1:A:368:LEU:HD21	2.01	0.43
1:A:228:SER:HB3	1:A:230:ASN:OD1	2.19	0.43
1:A:439:ILE:HG22	1:A:440:HIS:O	2.19	0.43
1:A:206:VAL:CG1	1:A:222:ALA:HB1	2.49	0.42
1:A:353:MET:HG3	5:A:2106:HOH:O	2.18	0.42
1:A:216:ARG:HG2	1:A:216:ARG:H	1.49	0.42
1:A:286:SER:OG	1:A:288:PHE:O	2.38	0.41
1:A:349:ARG:HA	1:A:349:ARG:HD2	1.89	0.41
1:A:350:GLU:OE1	1:A:350:GLU:HA	2.20	0.41
1:A:377:ASP:OD1	1:A:377:ASP:C	2.63	0.41
1:A:508:GLU:HB3	1:A:509:PRO:CD	2.51	0.41
1:A:19:ARG:CZ	1:A:26:HIS:HB2	2.50	0.41
3:A:3001:NAG:O6	5:A:2368:HOH:O	2.22	0.41
1:A:105:ARG:HA	1:A:106:PRO:HD3	1.92	0.41
1:A:502:PHE:CZ	1:A:513:HIS:HB2	2.55	0.41
1:A:221:ARG:HD3	1:A:318:GLN:OE1	2.21	0.41
1:A:383:ASN:OD1	1:A:383:ASN:C	2.63	0.41
1:A:484:GLU:HB2	1:A:487:SER:CB	2.51	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:LEU:HD23	1:A:143:LEU:HD11	2.03	0.40
1:A:207:GLY:HA3	1:A:229:PRO:HD3	2.03	0.40
1:A:461:GLU:CD	1:A:461:GLU:N	2.78	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	531/534 (99%)	508 (96%)	21 (4%)	2 (0%)	30 38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	485	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	466/466 (100%)	434 (93%)	32 (7%)	14 20

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	SER
1	A	22	VAL
1	A	38	PRO
1	A	44	ARG
1	A	89	GLU
1	A	129	VAL
1	A	140	GLU
1	A	143	LEU
1	A	145	SER
1	A	197	PHE
1	A	216	ARG
1	A	228	SER
1	A	246	VAL
1	A	281	VAL
1	A	287	ILE
1	A	295	VAL
1	A	299	GLU
1	A	330	PHE
1	A	337	PRO
1	A	356	VAL
1	A	357	LYS
1	A	358	LEU
1	A	365	ASP
1	A	368	LEU
1	A	430	LEU
1	A	450	LEU
1	A	454	LYS
1	A	461	GLU
1	A	499	GLU
1	A	516	LEU
1	A	518	VAL
1	A	531	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	ASN
1	A	68	GLN
1	A	519	GLN
1	A	526	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MIS	A	200	1	11,12,13	1.59	2 (18%)	11,16,18	1.13	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MIS	A	200	1	-	3/11/13/15	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	200	MIS	P-O3P	-3.89	1.47	1.59
1	A	200	MIS	P-OG	-2.81	1.48	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	200	MIS	P-O3P-C1	2.65	130.16	121.10

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	200	MIS	CB-OG-P-O2P
1	A	200	MIS	N-CA-CB-OG
1	A	200	MIS	C1-O3P-P-O1P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	200	MIS	1	0

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.75	0	17,19,21	0.87	0
2	NAG	B	2	2	14,14,15	1.44	1 (7%)	17,19,21	1.25	1 (5%)
2	NAG	C	1	1,2	14,14,15	1.48	3 (21%)	17,19,21	0.91	1 (5%)
2	NAG	C	2	2	14,14,15	1.20	2 (14%)	17,19,21	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	C1-C2	3.10	1.56	1.52
2	B	2	NAG	C4-C5	3.09	1.59	1.53
2	C	1	NAG	O5-C5	2.33	1.48	1.43
2	C	2	NAG	C1-C2	2.20	1.55	1.52
2	C	1	NAG	O5-C1	2.14	1.47	1.43
2	C	2	NAG	O5-C5	2.04	1.47	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	-4.47	116.91	122.90
2	C	1	NAG	C2-N2-C7	-2.47	119.59	122.90

There are no chirality outliers.


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	MES	A	4001	-	12,12,12	8.46	7 (58%)	15,16,16	2.31	7 (46%)
3	NAG	A	3001	1	14,14,15	1.15	1 (7%)	17,19,21	1.04	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	4001	-	-	3/6/14/14	0/1/1/1
3	NAG	A	3001	1	-	2/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	4001	MES	C8-S	-21.83	1.47	1.77
4	A	4001	MES	O1S-S	12.01	1.79	1.45
4	A	4001	MES	O2S-S	10.96	1.76	1.45
4	A	4001	MES	O3S-S	8.32	1.78	1.47
4	A	4001	MES	C7-C8	-4.32	1.41	1.52
4	A	4001	MES	C7-N4	-2.85	1.40	1.47
4	A	4001	MES	C3-C2	-2.57	1.40	1.50
3	A	3001	NAG	O5-C5	2.15	1.47	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	4001	MES	O3S-S-C8	5.11	116.01	106.00
4	A	4001	MES	O1S-S-C8	3.94	112.69	106.73
4	A	4001	MES	O3S-S-O2S	-2.96	103.99	111.40
3	A	3001	NAG	C2-N2-C7	-2.49	119.56	122.90
4	A	4001	MES	O1-C2-C3	-2.48	106.43	111.77
4	A	4001	MES	O3S-S-O1S	-2.34	105.55	111.40
4	A	4001	MES	C8-C7-N4	2.32	121.12	112.36
3	A	3001	NAG	O5-C1-C2	-2.31	107.71	111.29
4	A	4001	MES	O2S-S-C8	2.16	110.00	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	4001	MES	C7-C8-S-O2S
3	A	3001	NAG	O5-C5-C6-O6
3	A	3001	NAG	C4-C5-C6-O6
4	A	4001	MES	C7-C8-S-O3S
4	A	4001	MES	C7-C8-S-O1S

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3001	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	533/534 (99%)	-0.63	2 (0%) 88 89	25, 39, 60, 91	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ASP	3.2
1	A	3	HIS	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	MIS	A	200	13/14	0.99	0.04	26,28,33,34	0


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.70	0.13	67,73,75,76	0
2	NAG	B	2	14/15	0.78	0.12	61,69,73,75	0
2	NAG	C	1	14/15	0.81	0.11	70,72,73,73	0
2	NAG	B	1	14/15	0.91	0.08	45,51,58,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	3001	14/15	0.78	0.12	67,69,72,74	0
4	MES	A	4001	12/12	0.96	0.10	64,66,67,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.