



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 03:39 PM UTC

PDB ID : 3DGV / pdb_00003dgv
Title : Crystal structure of thrombin activatable fibrinolysis inhibitor (TAFI)
Authors : Anand, K.; Pallares, I.; Valnickova, Z.; Christensen, T.; Schreuder, H.; Enghild, J.
Deposited on : 2008-06-16
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	27% 70% 27% •
1	B	401	66% 31% •
1	C	401	27% 37% 50% 7% 5%
2	D	2	100%
2	I	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	2	 50% 50%
4	F	2	 50% 50%
4	J	2	 50% 50%
5	G	3	 33% 67%
6	H	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	NAG	C	309	X	-	-	-
5	NAG	G	1	X	-	-	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 11015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carboxypeptidase B2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	0	0
			3278	2097	565	600	16			
1	B	401	Total	C	N	O	S	0	0	0
			3275	2094	564	601	16			
1	C	379	Total	C	N	O	S	0	0	0
			3068	1958	525	569	16			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called beta-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			24	14	1	9			


- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a

cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	G	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	H	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		
7	B	1	Total	Zn	0	0
			1	1		
7	C	1	Total	Zn	0	0
			1	1		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	C	1	Total	O	S	0	0
			5	4	1		
8	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is MALONATE ION (CCD ID: MLI) (formula: $\text{C}_3\text{H}_2\text{O}_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			7	3	4		
9	B	1	Total	C	O	0	0
			7	3	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	C	1	Total	C	O	0	0
			7	3	4		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	1	Total	C	N	O	0	0
			14	8	1	5		
10	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	484	Total	O	0	0
			484	484		
11	B	417	Total	O	0	0
			417	417		
11	C	118	Total	O	0	0
			118	118		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carboxypeptidase B2




• Molecule 1: Carboxypeptidase B2




• Molecule 1: Carboxypeptidase B2





- Molecule 2: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%



- Molecule 2: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%



- Molecule 3: beta-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%


MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%

MAG1
MAG2

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  33% 67%

MAG1
MAG2
FUL3

- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  33% 67%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	146.47Å 146.47Å 231.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.86 – 2.50 19.86 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.86-2.50) 99.4 (19.86-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.50Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.205 , 0.243 0.208 , 0.244	Depositor DCC
R_{free} test set	4345 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 80.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11015	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FUL, NAG, BMA, SO4, MLI, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3367	0.74	1/4560 (0.0%)
1	B	0.42	0/3364	0.74	2/4557 (0.0%)
1	C	0.29	0/3147	0.73	2/4261 (0.0%)
All	All	0.39	0/9878	0.74	5/13378 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	ALA	N-CA-C	-7.06	103.69	112.72
1	C	222	VAL	N-CA-C	-5.89	108.11	113.71
1	B	58	ALA	CB-CA-C	-5.38	110.39	116.63
1	A	138	CYS	N-CA-C	5.09	117.61	109.81
1	C	61	ALA	N-CA-C	5.06	115.98	108.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	92(A)	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3278	0	3194	103	0
1	B	3275	0	3183	108	0
1	C	3068	0	2926	228	0
2	D	28	0	24	0	0
2	I	28	0	24	1	0
3	E	24	0	22	1	0
4	F	28	0	25	2	0
4	J	28	0	25	2	0
5	G	38	0	34	0	0
6	H	39	0	34	2	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
8	A	45	0	0	2	0
8	B	55	0	0	0	0
8	C	10	0	0	1	0
9	A	7	0	2	0	0
9	B	7	0	2	1	0
9	C	7	0	2	0	0
10	B	14	0	13	2	0
10	C	14	0	13	0	0
11	A	484	0	0	19	0
11	B	417	0	0	20	0
11	C	118	0	0	17	0
All	All	11015	0	9523	441	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 441 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75(A):ASN:ND2	1:A:78(A):ASP:H	1.55	1.03
1:C:281:LEU:HD12	1:C:282:PRO:HD2	1.54	0.89
1:C:201:LYS:H	1:C:270:GLU:HB2	1.39	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:ASN:HB2	1:C:160:SER:OG	1.77	0.84
1:C:75:SER:HB3	1:C:76:PRO:HD3	1.60	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/401 (100%)	386 (97%)	12 (3%)	1 (0%)	36	55
1	B	399/401 (100%)	373 (94%)	20 (5%)	6 (2%)	8	16
1	C	365/401 (91%)	304 (83%)	47 (13%)	14 (4%)	2	3
All	All	1163/1203 (97%)	1063 (91%)	79 (7%)	21 (2%)	6	12

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	SER
1	B	91(A)	PRO
1	B	199	SER
1	C	91(A)	PRO
1	C	131	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/358 (100%)	326 (91%)	32 (9%)	9	20
1	B	357/358 (100%)	340 (95%)	17 (5%)	23	46
1	C	332/358 (93%)	312 (94%)	20 (6%)	17	36
All	All	1047/1074 (98%)	978 (93%)	69 (7%)	15	32

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	74	ILE
1	C	136	ASN
1	C	264	LYS
1	A	212	ARG
1	A	210	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	16	ASN
1	C	216	HIS
1	B	240	HIS
1	C	303	HIS
1	C	102	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	D	1	2,1	14,14,15	0.87	1 (7%)	17,19,21	1.21	3 (17%)
2	NDG	D	2	2	14,14,15	0.79	1 (7%)	17,19,21	0.67	0
3	NAG	E	1	3,1	14,14,15	0.89	1 (7%)	17,19,21	0.74	0
3	FUL	E	2	3	10,10,11	0.55	0	14,14,16	0.42	0
4	NAG	F	1	4,1	14,14,15	0.60	0	17,19,21	0.78	1 (5%)
4	NAG	F	2	4	14,14,15	0.71	0	17,19,21	0.55	0
5	NAG	G	1	5,1	14,14,15	0.60	0	17,19,21	0.87	1 (5%)
5	NAG	G	2	5	14,14,15	0.61	0	17,19,21	0.68	1 (5%)
5	FUL	G	3	5	10,10,11	0.66	0	14,14,16	0.56	0
6	NAG	H	1	1,6	14,14,15	0.56	0	17,19,21	0.81	1 (5%)
6	NAG	H	2	6	14,14,15	0.73	0	17,19,21	1.01	0
6	BMA	H	3	6	11,11,12	0.82	0	15,15,17	0.93	1 (6%)
2	NAG	I	1	2,1	14,14,15	0.80	0	17,19,21	1.31	2 (11%)
2	NDG	I	2	2	14,14,15	0.76	0	17,19,21	0.98	2 (11%)
4	NAG	J	1	4,1	14,14,15	0.66	0	17,19,21	0.67	0
4	NAG	J	2	4	14,14,15	0.69	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NDG	D	2	2	-	2/6/23/26	0/1/1/1
3	NAG	E	1	3,1	-	5/6/23/26	0/1/1/1
3	FUL	E	2	3	-	-	0/1/1/1
4	NAG	F	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	F	2	4	-	5/6/23/26	0/1/1/1
5	NAG	G	1	5,1	1/1/5/7	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	4/6/23/26	0/1/1/1
5	FUL	G	3	5	-	-	0/1/1/1
6	NAG	H	1	1,6	-	5/6/23/26	0/1/1/1
6	NAG	H	2	6	-	4/6/23/26	0/1/1/1
6	BMA	H	3	6	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	I	1	2,1	-	4/6/23/26	0/1/1/1
2	NDG	I	2	2	-	4/6/23/26	0/1/1/1
4	NAG	J	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	J	2	4	-	5/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1	NAG	C1-C2	2.61	1.55	1.52
2	D	1	NAG	C1-C2	2.41	1.55	1.52
2	D	2	NDG	C1-C2	2.19	1.55	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	1	NAG	C4-C3-C2	3.13	115.60	111.02
2	I	1	NAG	C3-C4-C5	2.91	115.51	110.23
6	H	3	BMA	C1-O5-C5	2.51	115.55	112.19
2	I	2	NDG	O5-C1-C2	-2.41	107.56	111.29
2	I	2	NDG	C2-N2-C7	-2.30	119.82	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	G	1	NAG	C1

5 of 49 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	1	NAG	C8-C7-N2-C2
2	I	1	NAG	O7-C7-N2-C2
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
4	F	1	NAG	C8-C7-N2-C2

There are no ring outliers.

8 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	1	NAG	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	2	NAG	2	0
4	J	1	NAG	2	0
2	I	1	NAG	1	0
6	H	2	NAG	2	0
6	H	3	BMA	1	0
4	F	1	NAG	2	0
3	E	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 3 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	SO4	A	325	-	4,4,4	0.35	0	6,6,6	0.06	0
8	SO4	B	323	-	4,4,4	0.38	0	6,6,6	0.11	0
8	SO4	B	321	-	4,4,4	0.36	0	6,6,6	0.10	0
8	SO4	B	326	-	4,4,4	0.37	0	6,6,6	0.10	0
8	SO4	B	324	-	4,4,4	0.38	0	6,6,6	0.07	0
8	SO4	B	322	-	4,4,4	0.39	0	6,6,6	0.08	0
8	SO4	C	311	-	4,4,4	0.37	0	6,6,6	0.10	0
8	SO4	A	327	-	4,4,4	0.33	0	6,6,6	0.17	0
8	SO4	A	324	-	4,4,4	0.33	0	6,6,6	0.10	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	SO4	B	318	-	4,4,4	0.34	0	6,6,6	0.18	0
8	SO4	B	319	-	4,4,4	0.35	0	6,6,6	0.16	0
9	MLI	A	328	7	6,6,6	1.23	0	7,7,7	0.97	0
9	MLI	C	313	-	6,6,6	1.13	0	7,7,7	1.33	1 (14%)
10	NAG	C	309	1	14,14,15	0.51	0	17,19,21	0.73	1 (5%)
8	SO4	A	322	-	4,4,4	0.42	0	6,6,6	0.09	0
10	NAG	B	316	1	14,14,15	0.59	0	17,19,21	0.79	0
8	SO4	B	320	-	4,4,4	0.36	0	6,6,6	0.18	0
8	SO4	A	323	-	4,4,4	0.34	0	6,6,6	0.20	0
8	SO4	B	325	-	4,4,4	0.37	0	6,6,6	0.08	0
8	SO4	B	328	-	4,4,4	0.37	0	6,6,6	0.11	0
8	SO4	A	319	-	4,4,4	0.33	0	6,6,6	0.17	0
8	SO4	A	321	-	4,4,4	0.38	0	6,6,6	0.10	0
8	SO4	C	312	-	4,4,4	0.35	0	6,6,6	0.15	0
8	SO4	B	327	-	4,4,4	0.35	0	6,6,6	0.10	0
8	SO4	A	326	-	4,4,4	0.35	0	6,6,6	0.10	0
8	SO4	A	320	-	4,4,4	0.37	0	6,6,6	0.11	0
9	MLI	B	329	-	6,6,6	1.20	0	7,7,7	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	MLI	C	313	-	-	2/4/4/4	-
9	MLI	A	328	7	-	2/4/4/4	-
10	NAG	C	309	1	1/1/5/7	2/6/23/26	0/1/1/1
9	MLI	B	329	-	-	2/4/4/4	-
10	NAG	B	316	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	C	313	MLI	C3-C1-C2	-2.41	104.45	112.95
10	C	309	NAG	C2-N2-C7	-2.22	119.93	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
10	C	309	NAG	C1

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	316	NAG	C8-C7-N2-C2
10	B	316	NAG	O7-C7-N2-C2
10	C	309	NAG	C8-C7-N2-C2
10	C	309	NAG	O7-C7-N2-C2
10	B	316	NAG	C4-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	C	311	SO4	1	0
8	A	327	SO4	1	0
10	B	316	NAG	2	0
8	A	323	SO4	1	0
9	B	329	MLI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/401 (100%)	-0.39	4 (0%) 79 76	29, 44, 66, 87	0
1	B	401/401 (100%)	-0.29	3 (0%) 84 81	30, 49, 70, 93	0
1	C	379/401 (94%)	1.55	108 (28%) 1 1	72, 123, 156, 160	0
All	All	1181/1203 (98%)	0.27	115 (9%) 13 11	29, 56, 150, 160	0

The worst 5 of 115 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	279	PHE	5.4
1	C	228	PHE	4.7
1	C	37(A)	ALA	4.4
1	C	72(A)	LEU	4.4
1	C	19(A)	ILE	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	BMA	H	3	11/12	0.11	0.20	102,103,104,104	0
5	FUL	G	3	10/11	0.26	0.20	100,101,101,102	0
4	NAG	J	2	14/15	0.30	0.18	106,107,108,108	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NDG	I	2	14/15	0.33	0.19	96,98,99,99	0
3	FUL	E	2	10/11	0.37	0.17	100,101,102,102	0
3	NAG	E	1	14/15	0.39	0.15	88,92,94,98	0
4	NAG	F	2	14/15	0.41	0.17	103,104,105,105	0
5	NAG	G	2	14/15	0.51	0.20	102,103,104,105	0
4	NAG	F	1	14/15	0.52	0.18	95,97,99,101	0
2	NDG	D	2	14/15	0.60	0.14	82,84,85,86	0
4	NAG	J	1	14/15	0.61	0.18	97,99,101,104	0
6	NAG	H	2	14/15	0.64	0.17	90,94,95,99	0
2	NAG	I	1	14/15	0.65	0.14	84,86,89,93	0
5	NAG	G	1	14/15	0.76	0.14	90,93,99,99	0
2	NAG	D	1	14/15	0.84	0.12	69,71,75,79	0
6	NAG	H	1	14/15	0.89	0.11	72,74,79,85	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	NAG	B	316	14/15	0.52	0.16	85,88,89,90	0
10	NAG	C	309	14/15	0.52	0.19	151,152,152,152	0
8	SO4	C	312	5/5	0.64	0.17	136,136,136,136	0
8	SO4	A	327	5/5	0.69	0.18	98,99,99,99	0
8	SO4	B	322	5/5	0.80	0.15	127,127,127,127	0
8	SO4	A	325	5/5	0.82	0.12	114,114,114,114	0
8	SO4	B	328	5/5	0.84	0.16	140,140,140,140	0
8	SO4	B	324	5/5	0.84	0.13	103,103,104,104	0
8	SO4	C	311	5/5	0.86	0.12	110,110,110,110	0
9	MLI	B	329	7/7	0.88	0.17	66,67,68,68	0
8	SO4	B	326	5/5	0.88	0.11	129,129,129,129	0
9	MLI	A	328	7/7	0.88	0.19	72,73,74,74	0
8	SO4	B	327	5/5	0.89	0.15	94,94,95,95	0
8	SO4	A	326	5/5	0.89	0.10	147,147,147,147	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	SO4	B	321	5/5	0.91	0.12	100,100,100,100	0
8	SO4	A	323	5/5	0.91	0.13	92,92,93,93	0
8	SO4	B	323	5/5	0.92	0.10	105,105,105,105	0
8	SO4	A	322	5/5	0.93	0.13	86,86,86,87	0
9	MLI	C	313	7/7	0.93	0.10	109,109,109,109	0
8	SO4	B	320	5/5	0.93	0.10	73,73,74,74	0
8	SO4	B	325	5/5	0.93	0.10	118,118,118,118	0
8	SO4	A	324	5/5	0.94	0.11	80,80,81,81	0
8	SO4	A	321	5/5	0.94	0.11	80,81,81,81	0
8	SO4	A	320	5/5	0.95	0.08	71,72,72,72	0
8	SO4	B	319	5/5	0.95	0.13	73,74,74,75	0
8	SO4	B	318	5/5	0.97	0.06	58,58,59,60	0
7	ZN	C	310	1/1	0.97	0.08	110,110,110,110	0
8	SO4	A	319	5/5	0.98	0.05	51,52,52,53	0
7	ZN	A	318	1/1	1.00	0.05	39,39,39,39	0
7	ZN	B	317	1/1	1.00	0.05	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.