



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:11 PM UTC

PDB ID : 5DGR / pdb_00005dgr
Title : Crystal structure of GH9 exo-beta-D-glucosaminidase PBPRA0520, glucosamine complex
Authors : Suzuki, K.; Honda, Y.; Fushinobu, S.
Deposited on : 2015-08-28
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	586	% 87% 10% ..
1	B	586	% 83% 13% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative endoglucanase-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	573	Total	C	N	O	S	0	0	0
			4573	2888	788	864	33			
1	B	571	Total	C	N	O	S	0	0	0
			4557	2878	786	860	33			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	579	LEU	-	expression tag	UNP Q6LUT2
A	580	GLU	-	expression tag	UNP Q6LUT2
A	581	HIS	-	expression tag	UNP Q6LUT2
A	582	HIS	-	expression tag	UNP Q6LUT2
A	583	HIS	-	expression tag	UNP Q6LUT2
A	584	HIS	-	expression tag	UNP Q6LUT2
A	585	HIS	-	expression tag	UNP Q6LUT2
A	586	HIS	-	expression tag	UNP Q6LUT2
B	579	LEU	-	expression tag	UNP Q6LUT2
B	580	GLU	-	expression tag	UNP Q6LUT2
B	581	HIS	-	expression tag	UNP Q6LUT2
B	582	HIS	-	expression tag	UNP Q6LUT2
B	583	HIS	-	expression tag	UNP Q6LUT2
B	584	HIS	-	expression tag	UNP Q6LUT2
B	585	HIS	-	expression tag	UNP Q6LUT2
B	586	HIS	-	expression tag	UNP Q6LUT2

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is 2-amino-2-deoxy-beta-D-glucopyranose (CCD ID: GCS) (formula: $C_6H_{13}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			12	6	1	5		
3	B	1	Total	C	N	O	0	0
			12	6	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	390	Total	O	0	0
			390	390		
4	B	343	Total	O	0	0
			343	343		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative endoglucanase-related protein



• Molecule 1: Putative endoglucanase-related protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.29Å 102.59Å 89.80Å 90.00° 97.22° 90.00°	Depositor
Resolution (Å)	28.90 – 1.90 28.90 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (28.90-1.90) 99.4 (28.90-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.162 , 0.204 0.162 , 0.205	Depositor DCC
R_{free} test set	4417 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9889	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.44	16/4690 (0.3%)	1.23	8/6357 (0.1%)
1	B	1.42	18/4673 (0.4%)	1.23	15/6332 (0.2%)
All	All	1.43	34/9363 (0.4%)	1.23	23/12689 (0.2%)

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	161	GLN	CA-C	9.03	1.57	1.52
1	B	535	GLY	C-O	7.05	1.30	1.23
1	B	245	GLY	C-O	6.76	1.31	1.23
1	B	539	LYS	N-CA	6.74	1.52	1.46
1	A	14	GLY	N-CA	6.55	1.53	1.45
1	B	323	TYR	N-CA	6.49	1.54	1.46
1	A	575	ARG	CA-C	6.28	1.61	1.52
1	B	121	LYS	CA-CB	6.25	1.59	1.53
1	B	559	PRO	N-CA	6.24	1.55	1.47
1	B	378	ALA	CA-C	6.23	1.61	1.52
1	B	40	GLN	CA-C	6.19	1.60	1.52
1	A	216	PHE	C-O	5.86	1.31	1.24
1	B	82	THR	C-O	5.78	1.31	1.24
1	A	408	SER	N-CA	5.73	1.53	1.46
1	A	436	PHE	N-CA	5.70	1.52	1.45
1	B	16	LYS	CA-C	-5.70	1.45	1.52
1	B	399	LEU	N-CA	5.53	1.53	1.46
1	A	134	HIS	C-O	5.40	1.31	1.23
1	A	454	ILE	N-CA	5.25	1.52	1.46
1	B	156	TYR	N-CA	5.22	1.52	1.46
1	A	248	GLN	N-CA	5.21	1.52	1.46
1	B	118	ALA	CA-CB	5.21	1.61	1.53
1	A	167	TRP	CA-C	5.16	1.59	1.52
1	B	245	GLY	N-CA	5.11	1.51	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	125	PHE	C-O	5.10	1.29	1.23
1	B	11	GLU	CA-C	5.10	1.59	1.52
1	A	470	ASP	C-O	5.08	1.30	1.23
1	B	509	TYR	CA-C	5.06	1.55	1.52
1	B	563	TRP	N-CA	5.06	1.52	1.46
1	A	381	GLN	N-CA	5.05	1.52	1.46
1	A	22	ALA	N-CA	5.04	1.52	1.45
1	A	291	TYR	C-O	5.03	1.29	1.24
1	B	284	LEU	CA-C	5.01	1.59	1.52
1	A	377	ILE	CA-CB	5.00	1.59	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	329	ASN	N-CA-C	7.41	119.36	111.28
1	B	551	TRP	N-CA-C	-6.39	104.74	112.54
1	B	558	ILE	CA-C-N	-6.29	112.36	119.28
1	B	558	ILE	C-N-CA	-6.29	112.36	119.28
1	B	504	ARG	NE-CZ-NH1	-6.29	115.21	121.50
1	A	287	ALA	N-CA-C	-5.68	105.17	111.36
1	A	554	GLY	N-CA-C	5.67	122.74	115.32
1	B	504	ARG	NE-CZ-NH2	5.65	124.29	119.20
1	B	40	GLN	N-CA-C	5.62	117.72	109.24
1	B	77	ILE	N-CA-C	-5.62	100.39	108.53
1	A	154	GLY	N-CA-C	-5.54	107.01	115.66
1	A	471	ASN	N-CA-C	5.54	118.09	111.71
1	A	374	LEU	CA-C-N	-5.52	113.50	119.24
1	A	374	LEU	C-N-CA	-5.52	113.50	119.24
1	A	124	LEU	CA-CB-CG	5.48	135.47	116.30
1	B	122	VAL	CB-CA-C	-5.35	101.63	111.36
1	B	154	GLY	N-CA-C	-5.26	107.45	115.66
1	B	378	ALA	N-CA-C	-5.26	105.72	111.82
1	B	459	THR	N-CA-C	-5.18	105.53	111.07
1	A	451	ASN	N-CA-C	5.14	117.55	111.33
1	B	36	CYS	CA-C-N	5.08	128.44	120.82
1	B	36	CYS	C-N-CA	5.08	128.44	120.82
1	B	367	PHE	N-CA-C	-5.08	101.02	108.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4573	0	4322	19	0
1	B	4557	0	4306	42	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	13	0	0
3	B	12	0	13	0	0
4	A	390	0	0	0	0
4	B	343	0	0	9	0
All	All	9889	0	8654	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:MET:CE	1:B:341:LYS:HE3	1.67	1.25
1:B:337:MET:HE2	1:B:341:LYS:CE	1.67	1.23
1:B:337:MET:CE	1:B:341:LYS:CE	2.34	0.97
1:B:337:MET:HE1	1:B:341:LYS:HE3	1.51	0.90
1:B:413:VAL:HG12	4:B:837:HOH:O	1.73	0.89
1:B:279:LYS:HA	4:B:860:HOH:O	1.73	0.88
1:B:50:CYS:HB3	4:B:953:HOH:O	1.79	0.81
1:B:337:MET:HE2	1:B:341:LYS:HE2	1.62	0.79
1:B:337:MET:HE2	1:B:341:LYS:CD	2.18	0.73
1:B:116:GLU:HG3	4:B:753:HOH:O	1.90	0.72
1:B:214:THR:HG23	1:B:229:CYS:O	1.94	0.68
1:B:127:THR:HG23	1:B:129:GLU:H	1.59	0.67
1:B:413:VAL:CG1	4:B:837:HOH:O	2.39	0.59
1:B:337:MET:HE2	1:B:341:LYS:HD2	1.88	0.56
1:A:463:MET:HG3	1:A:570:MET:HE2	1.89	0.55
1:B:512:GLU:HB2	4:B:916:HOH:O	2.05	0.55
1:B:382:TYR:OH	1:B:386:GLU:OE1	2.27	0.52
1:B:463:MET:HG3	1:B:570:MET:HE2	1.92	0.52
1:A:524:ASN:HA	1:A:560:HIS:NE2	2.25	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:ASP:HA	1:B:122:VAL:HG22	1.92	0.51
1:B:413:VAL:HG11	4:B:1013:HOH:O	2.11	0.51
1:B:43:MET:HE2	1:B:45:LEU:HD21	1.93	0.50
1:B:367:PHE:CE1	1:B:438:PRO:HD2	2.48	0.49
1:B:462:TYR:CE2	1:B:571:GLN:HB2	2.48	0.49
1:A:404:ASN:OD1	1:A:473:ILE:HD12	2.12	0.49
1:B:122:VAL:HG12	1:B:227:GLU:HA	1.96	0.48
1:B:347:MET:HE1	1:B:360:ASP:O	2.13	0.47
1:B:400:LEU:HD11	1:B:473:ILE:HG13	1.97	0.47
1:B:557:TRP:CD2	1:B:559:PRO:HD2	2.50	0.47
1:B:8:LEU:HD12	1:B:8:LEU:N	2.29	0.47
1:A:114:ILE:HD12	1:B:273:PRO:HB2	1.97	0.47
1:A:216:PHE:CZ	1:A:218:LYS:HA	2.51	0.46
1:A:542:LYS:HA	1:A:542:LYS:HD2	1.73	0.46
1:B:542:LYS:HA	1:B:542:LYS:HD2	1.48	0.45
1:B:78:ARG:NH2	1:B:81:ASP:OD1	2.49	0.45
1:B:9:GLY:HA3	1:B:88:CYS:O	2.17	0.45
1:A:262:ARG:HG3	1:A:325:SER:OG	2.17	0.44
1:A:273:PRO:HB2	1:B:114:ILE:HD12	2.00	0.43
1:A:220:SER:O	1:A:221:LYS:HB2	2.19	0.43
1:B:413:VAL:HG13	4:B:940:HOH:O	2.17	0.43
1:B:213:MET:C	1:B:214:THR:HG22	2.44	0.43
1:B:374:LEU:HD23	1:B:374:LEU:HA	1.93	0.43
1:B:573:LYS:HG3	4:B:766:HOH:O	2.18	0.43
1:A:541:GLU:HA	1:A:544:LYS:HD3	2.00	0.43
1:B:96:GLN:HG3	1:B:97:ASN:OD1	2.18	0.42
1:B:173:TYR:CE1	1:B:191:VAL:HB	2.54	0.42
1:B:557:TRP:HB3	1:B:560:HIS:CD2	2.54	0.42
1:A:462:TYR:CE2	1:A:571:GLN:HB2	2.54	0.42
1:A:65:ASP:OD1	1:A:65:ASP:C	2.62	0.42
1:B:216:PHE:CZ	1:B:218:LYS:HA	2.55	0.41
1:A:458:ILE:CG1	1:A:484:LEU:HB3	2.51	0.41
1:A:96:GLN:HG2	1:A:97:ASN:OD1	2.20	0.41
1:A:558:ILE:N	1:A:559:PRO:CD	2.84	0.41
1:A:95:MET:HE2	1:A:95:MET:HB2	1.74	0.41
1:B:53:VAL:O	1:B:54:ALA:C	2.63	0.40
1:A:173:TYR:CE1	1:A:191:VAL:HB	2.57	0.40
1:A:295:LYS:HA	1:A:295:LYS:HD2	1.84	0.40
1:B:297:MET:O	1:B:298:ASN:C	2.65	0.40
1:A:45:LEU:HD22	1:A:65:ASP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	569/586 (97%)	547 (96%)	21 (4%)	1 (0%)	43	36
1	B	565/586 (96%)	548 (97%)	17 (3%)	0	100	100
All	All	1134/1172 (97%)	1095 (97%)	38 (3%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	TYR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	476/488 (98%)	466 (98%)	10 (2%)	47	44
1	B	474/488 (97%)	461 (97%)	13 (3%)	39	34
All	All	950/976 (97%)	927 (98%)	23 (2%)	43	38

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ILE
1	A	40	GLN
1	A	44	GLN
1	A	81	ASP
1	A	122	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	180	GLU
1	A	214	THR
1	A	218	LYS
1	A	375	PRO
1	A	542	LYS
1	B	1	MET
1	B	23	GLN
1	B	40	GLN
1	B	76	ARG
1	B	122	VAL
1	B	163	PRO
1	B	214	THR
1	B	266	LEU
1	B	328	GLU
1	B	329	ASN
1	B	390	GLN
1	B	493	PRO
1	B	542	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	396	GLN
1	A	476	GLN
1	B	2	GLN
1	B	30	HIS
1	B	57	HIS
1	B	225	GLN
1	B	274	GLN
1	B	330	ASN
1	B	354	HIS
1	B	471	ASN
1	B	476	GLN
1	B	577	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GCS	B	602	-	12,12,12	2.34	5 (41%)	16,17,17	1.89	6 (37%)
3	GCS	A	602	-	12,12,12	2.26	5 (41%)	16,17,17	2.02	5 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GCS	B	602	-	-	0/2/22/22	0/1/1/1
3	GCS	A	602	-	-	0/2/22/22	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	GCS	C3-C2	5.00	1.59	1.53
3	A	602	GCS	C3-C2	4.50	1.59	1.53
3	B	602	GCS	O5-C1	3.70	1.51	1.42
3	A	602	GCS	O5-C1	3.20	1.50	1.42
3	B	602	GCS	C2-N2	3.12	1.52	1.47
3	A	602	GCS	C2-N2	2.86	1.51	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	GCS	C4-C3	2.59	1.59	1.52
3	A	602	GCS	C4-C3	2.58	1.59	1.52
3	A	602	GCS	O5-C5	2.33	1.50	1.44
3	B	602	GCS	O5-C5	2.13	1.49	1.44

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	GCS	O1-C1-O5	-4.34	97.52	110.41
3	B	602	GCS	O1-C1-O5	-3.75	99.27	110.41
3	A	602	GCS	O5-C1-C2	-3.70	105.19	109.51
3	B	602	GCS	C1-C2-C3	-3.24	106.34	110.60
3	A	602	GCS	O1-C1-C2	-3.18	102.40	108.96
3	A	602	GCS	C1-O5-C5	-2.86	108.11	113.65
3	B	602	GCS	C4-C3-C2	2.78	115.60	110.99
3	B	602	GCS	O5-C5-C4	-2.53	105.14	109.70
3	B	602	GCS	O5-C1-C2	-2.51	106.58	109.51
3	A	602	GCS	O4-C4-C3	-2.32	104.90	110.38
3	B	602	GCS	O4-C4-C3	-2.17	105.27	110.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	573/586 (97%)	-0.38	7 (1%) 76 79	11, 21, 43, 87	0
1	B	571/586 (97%)	-0.31	4 (0%) 84 86	11, 22, 43, 77	0
All	All	1144/1172 (97%)	-0.35	11 (0%) 79 82	11, 21, 43, 87	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	578	VAL	3.6
1	A	578	VAL	3.4
1	B	276	GLY	2.6
1	A	29	HIS	2.5
1	B	577	HIS	2.4
1	A	42	ILE	2.2
1	A	38	SER	2.2
1	A	36	CYS	2.2
1	A	81	ASP	2.2
1	B	34	ILE	2.1
1	A	40	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GCS	A	602	12/12	0.96	0.05	16,19,21,24	0
2	NA	B	601	1/1	0.97	0.04	26,26,26,26	0
3	GCS	B	602	12/12	0.97	0.05	15,17,19,20	0
2	NA	A	601	1/1	0.99	0.09	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.