



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:18 PM UTC

PDB ID : 7DG2 / pdb_00007dg2
Title : Nse1-Nse3-Nse4 complex
Authors : Cho, Y.; Jo, A.
Deposited on : 2020-11-10
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	13% 89% 6%
2	C	217	8% 87% 8% 5%
3	D	79	16% 54% 14% 32%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural maintenance of chromosomes element 1 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1871	1174	332	346	19			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	expression tag	UNP Q6PAF4

- Molecule 2 is a protein called MAGE domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	207	Total	C	N	O	S	0	0	0
			1724	1108	295	315	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	44	MET	-	expression tag	UNP A0A1L8G3Z0

- Molecule 3 is a protein called Non-structural maintenance of chromosomes element 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	54	Total	C	N	O	S	0	0	0
			436	287	67	79	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	105	GLY	-	expression tag	UNP B1WBD6
D	106	SER	-	expression tag	UNP B1WBD6
D	107	HIS	-	expression tag	UNP B1WBD6

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	155	Total	O	0	0
			155	155		
7	C	132	Total	O	0	0
			132	132		
7	D	30	Total	O	0	0
			30	30		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-structural maintenance of chromosomes element 1 homolog



- Molecule 2: MAGE domain-containing protein



- Molecule 3: Non-structural maintenance of chromosomes element 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.49Å 66.16Å 168.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.59 – 1.70 41.59 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (41.59-1.70) 98.6 (41.59-1.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.190 , 0.218 0.191 , 0.216	Depositor DCC
R_{free} test set	3269 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4370	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1906	0.46	0/2565
2	C	0.27	0/1756	0.46	0/2358
3	D	0.25	0/450	0.46	0/607
All	All	0.27	0/4112	0.46	0/5530

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1871	0	1838	7	0
2	C	1724	0	1772	12	0
3	D	436	0	412	9	0
4	A	4	0	3	0	0
4	C	4	0	3	0	0
5	A	12	0	16	1	0
6	A	2	0	0	0	0
7	A	155	0	0	3	0
7	C	132	0	0	1	0
7	D	30	0	0	2	0
All	All	4370	0	4044	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:47:HIS:O	7:C:401:HOH:O	2.10	0.70
2:C:133:MET:HE1	3:D:170:PRO:HD3	1.75	0.69
2:C:164:ILE:HD12	2:C:214:PHE:CD2	2.31	0.66
2:C:80:ARG:HB3	2:C:80:ARG:HH21	1.62	0.64
2:C:172:ARG:HH12	3:D:149:TYR:HA	1.62	0.63
2:C:53:ASN:OD1	2:C:96:ARG:NH2	2.23	0.62
2:C:112:ILE:HG22	2:C:222:LYS:HG3	1.82	0.61
2:C:158:THR:OG1	2:C:215:ARG:NH1	2.34	0.60
2:C:113:ASP:HB3	2:C:118:ILE:HG12	1.89	0.54
1:A:130:LEU:HD12	7:A:434:HOH:O	2.07	0.54
1:A:154:ILE:HD11	7:A:406:HOH:O	2.06	0.54
3:D:121:LEU:HA	3:D:124:MET:HE3	1.95	0.48
1:A:175:ARG:NH1	1:A:189:ARG:O	2.47	0.46
1:A:211:GLY:O	1:A:215:HIS:HB2	2.15	0.46
3:D:107:HIS:N	7:D:202:HOH:O	2.50	0.45
3:D:120:LEU:HB3	3:D:124:MET:HE2	1.99	0.45
3:D:107:HIS:N	7:D:201:HOH:O	2.49	0.44
2:C:51:GLN:O	2:C:55:LYS:HG2	2.18	0.44
2:C:90:TYR:CD1	2:C:90:TYR:C	2.96	0.43
1:A:115:ASN:ND2	7:A:405:HOH:O	2.52	0.42
2:C:112:ILE:HB	2:C:218:GLN:HB3	2.01	0.42
3:D:180:LYS:HA	3:D:181:THR:HA	1.77	0.42
3:D:124:MET:HB3	3:D:154:ALA:HB1	2.02	0.41
3:D:179:PHE:O	3:D:181:THR:HA	2.20	0.41
1:A:186:ASN:HB3	1:A:205:LEU:HD21	2.04	0.40
1:A:151:LYS:HE3	5:A:303:GOL:O2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/247 (92%)	225 (99%)	2 (1%)	0	100	100
2	C	205/217 (94%)	198 (97%)	7 (3%)	0	100	100
3	D	50/79 (63%)	48 (96%)	2 (4%)	0	100	100
All	All	482/543 (89%)	471 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/222 (94%)	208 (100%)	0	100	100
2	C	192/200 (96%)	189 (98%)	3 (2%)	55	41
3	D	47/66 (71%)	46 (98%)	1 (2%)	47	30
All	All	447/488 (92%)	443 (99%)	4 (1%)	70	62

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	80	ARG
2	C	90	TYR
2	C	114	THR
3	D	126	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	208	GLN
2	C	196	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	A	301	-	3,3,3	1.54	1 (33%)	3,3,3	1.48	0
5	GOL	A	303	-	5,5,5	0.95	0	5,5,5	1.06	0
5	GOL	A	302	-	5,5,5	0.84	0	5,5,5	1.07	0
4	ACT	C	301	-	3,3,3	1.39	0	3,3,3	1.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	303	-	-	3/4/4/4	-
5	GOL	A	302	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	ACT	CH3-C	2.16	1.57	1.49

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	302	GOL	O1-C1-C2-C3
5	A	303	GOL	O1-C1-C2-C3
5	A	303	GOL	O1-C1-C2-O2
5	A	302	GOL	O1-C1-C2-O2
5	A	302	GOL	O2-C2-C3-O3
5	A	303	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	303	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/247 (93%)	0.72	32 (13%) 6 6	14, 25, 47, 64	0
2	C	207/217 (95%)	0.43	17 (8%) 17 19	12, 24, 45, 67	0
3	D	54/79 (68%)	1.13	13 (24%) 2 2	15, 29, 52, 72	0
All	All	492/543 (90%)	0.64	62 (12%) 8 7	12, 25, 47, 72	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	233	ILE	6.5
2	C	209	PRO	6.3
3	D	126	LEU	6.2
3	D	149	TYR	5.1
2	C	208	ASP	4.8
3	D	148	GLY	4.6
2	C	210	VAL	4.4
1	A	210	ALA	4.3
1	A	2	MET	4.0
1	A	218	ALA	4.0
1	A	3	ASP	3.8
1	A	214	PHE	3.7
1	A	131	GLN	3.7
3	D	180	LYS	3.7
1	A	213	TYR	3.5
3	D	181	THR	3.5
3	D	150	LEU	3.4
1	A	228	SER	3.3
1	A	211	GLY	3.3
3	D	110	VAL	3.2
1	A	215	HIS	3.1
1	A	157	ASN	3.1
3	D	152	THR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	206	HIS	3.1
2	C	49	GLN	3.0
1	A	229	TRP	3.0
1	A	176	ASN	2.9
1	A	226	ASN	2.9
2	C	253	GLU	2.9
2	C	207	ILE	2.7
1	A	209	CYS	2.7
3	D	107	HIS	2.7
1	A	217	LYS	2.7
1	A	136	LYS	2.7
1	A	208	GLN	2.6
2	C	178	MET	2.6
1	A	115	ASN	2.6
1	A	207	LEU	2.6
1	A	223	PRO	2.5
2	C	129	GLN	2.5
1	A	47	LEU	2.5
2	C	114	THR	2.4
2	C	158	THR	2.4
1	A	212	LYS	2.4
2	C	115	LYS	2.3
3	D	125	GLY	2.3
1	A	232	GLU	2.3
3	D	109	THR	2.3
2	C	52	VAL	2.3
2	C	116	SER	2.2
1	A	180	ASP	2.2
2	C	128	VAL	2.2
3	D	115	SER	2.2
2	C	48	SER	2.2
1	A	220	PRO	2.2
3	D	108	MET	2.1
1	A	72	GLU	2.1
1	A	199	GLU	2.1
1	A	4	ARG	2.1
1	A	187	VAL	2.1
2	C	244	TRP	2.1
1	A	156	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACT	C	301	4/4	0.67	0.18	41,43,47,49	0
5	GOL	A	303	6/6	0.74	0.17	39,41,47,48	0
4	ACT	A	301	4/4	0.78	0.17	31,32,35,35	0
5	GOL	A	302	6/6	0.82	0.14	32,37,39,39	0
6	ZN	A	304	1/1	0.98	0.05	31,31,31,31	0
6	ZN	A	305	1/1	0.98	0.04	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around ZN A 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 305:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.