



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:24 PM UTC

PDB ID : 5DHL / pdb_00005dhl
Title : Crystal structure of Toxin, mutant N197W
Authors : Parker, M.W.; Gorman, M.A.; Lawrence, S.L.
Deposited on : 2015-08-31
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	5% 88% 10% ..
1	B	472	4% 88% 10% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 14593 atoms, of which 7082 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Perfringolysin O.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	467	Total	C	H	N	O	S	0	2	0
			7176	2329	3487	620	736	4			
1	B	467	Total	C	H	N	O	S	0	1	0
			7250	2329	3561	620	736	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	71	ALA	THR	conflict	UNP P0C2E9
A	197	TRP	ASN	engineered mutation	UNP P0C2E9
A	459	ALA	CYS	conflict	UNP P0C2E9
B	71	ALA	THR	conflict	UNP P0C2E9
B	197	TRP	ASN	engineered mutation	UNP P0C2E9
B	459	ALA	CYS	conflict	UNP P0C2E9

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		
2	B	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	49	Total	O	0	0
			49	49		
3	B	54	Total	O	0	0
			54	54		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Perfringolysin O



• Molecule 1: Perfringolysin O



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.63Å 212.40Å 46.96Å 90.00° 97.17° 90.00°	Depositor
Resolution (Å)	45.25 – 2.67 45.25 – 2.67	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.25-2.67) 98.9 (45.25-2.67)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 2.65Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.191 , 0.239 0.205 , 0.255	Depositor DCC
R_{free} test set	2196 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	52.9	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14593	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	2/3775 (0.1%)	1.17	14/5126 (0.3%)
1	B	0.87	1/3767 (0.0%)	1.11	8/5115 (0.2%)
All	All	0.88	3/7542 (0.0%)	1.14	22/10241 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	222	MET	SD-CE	-6.54	1.63	1.79
1	A	222	MET	SD-CE	-6.26	1.64	1.79
1	A	59	LYS	CA-C	5.19	1.60	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	ASP	CA-C-N	14.59	145.54	120.58
1	A	339	ASP	C-N-CA	14.59	145.54	120.58
1	A	134	ASP	CA-CB-CG	8.21	120.81	112.60
1	A	58	ASP	CA-CB-CG	7.41	120.00	112.60
1	B	470	VAL	N-CA-C	-7.33	103.81	111.58
1	B	60	ILE	CA-C-N	6.92	134.15	121.70
1	B	60	ILE	C-N-CA	6.92	134.15	121.70
1	A	167	GLU	N-CA-C	-6.07	101.50	110.48
1	A	302	ASP	CA-CB-CG	6.06	118.66	112.60
1	B	134	ASP	CA-CB-CG	5.57	118.17	112.60
1	A	294	PHE	CA-CB-CG	5.47	119.27	113.80
1	B	91	ASP	CA-CB-CG	5.38	117.98	112.60
1	B	243	ASP	CA-CB-CG	5.35	117.95	112.60
1	A	64	VAL	N-CA-CB	5.32	115.92	110.23
1	A	60	ILE	CA-C-N	5.29	131.65	121.54
1	A	60	ILE	C-N-CA	5.29	131.65	121.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	326	ASP	CA-CB-CG	5.27	117.87	112.60
1	B	239	LYS	N-CA-C	-5.16	105.73	111.36
1	A	59	LYS	CA-C-N	5.16	130.98	121.70
1	A	59	LYS	C-N-CA	5.16	130.98	121.70
1	A	98	VAL	N-CA-CB	-5.09	105.33	111.94
1	A	236	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3689	3487	3628	13	0
1	B	3689	3561	3639	18	0
2	A	15	17	17	0	0
2	B	15	17	17	0	0
3	A	49	0	0	0	0
3	B	54	0	0	0	0
All	All	7511	7082	7301	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:ILE:HB	1:B:61:GLU:HA	1.68	0.74
1:B:98:VAL:O	1:B:98:VAL:HG12	2.01	0.60
1:B:60:ILE:HB	1:B:61:GLU:CA	2.32	0.58
1:B:297:LEU:HD13	1:B:313:TYR:CZ	2.40	0.56
1:B:165:TRP:CZ3	1:B:353:THR:HG21	2.42	0.55
1:B:287:SER:O	1:B:290:VAL:HG23	2.09	0.52
1:A:94:ILE:HD11	1:A:102:THR:HG21	1.94	0.50
1:B:222:MET:HE2	1:B:224:LEU:HD21	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:VAL:HG12	1:A:98:VAL:O	2.13	0.48
1:A:48:ASN:O	1:A:52:VAL:HG23	2.14	0.48
1:A:277:ILE:HD11	1:A:348:ASN:HB2	1.96	0.47
1:B:275:ARG:NH2	1:B:350:THR:O	2.47	0.47
1:B:90:VAL:O	1:B:363:THR:HG22	2.15	0.46
1:B:98:VAL:O	1:B:98:VAL:CG1	2.63	0.46
1:A:307:GLN:O	1:A:310:LYS:HG2	2.16	0.45
1:A:189:LYS:HG3	1:A:200:ALA:HB1	2.00	0.44
1:A:59:LYS:N	1:A:60:ILE:HA	2.32	0.44
1:A:400:GLY:O	1:A:437:ALA:HB2	2.17	0.43
1:A:167:GLU:O	1:A:168:LYS:CB	2.67	0.43
1:A:133:ILE:HD11	1:A:231:TYR:CD1	2.54	0.43
1:B:319:THR:HG23	1:B:334:VAL:HG22	2.00	0.43
1:B:283:THR:CG2	1:B:316:SER:OG	2.67	0.42
1:B:479:THR:HG21	1:B:499:TYR:HB3	2.01	0.42
1:A:464:TRP:HA	1:A:467:TRP:CE3	2.55	0.42
1:B:303:ILE:N	1:B:303:ILE:HD13	2.35	0.41
1:B:464:TRP:HA	1:B:467:TRP:CE3	2.55	0.41
1:B:133:ILE:HD11	1:B:231:TYR:CD1	2.55	0.41
1:A:325:GLY:HA3	1:A:358:TYR:CD2	2.56	0.41
1:B:297:LEU:HD13	1:B:313:TYR:CE1	2.56	0.41
1:A:437:ALA:HA	1:A:438:HIS:HA	1.88	0.40
1:B:270:ASN:HB2	1:B:363:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/472 (99%)	443 (95%)	22 (5%)	2 (0%)	30	51
1	B	466/472 (99%)	444 (95%)	22 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	933/944 (99%)	887 (95%)	44 (5%)	2 (0%)	43 65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	LYS
1	A	168	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	415/418 (99%)	392 (94%)	23 (6%)	19 41
1	B	414/418 (99%)	393 (95%)	21 (5%)	21 44
All	All	829/836 (99%)	785 (95%)	44 (5%)	20 43

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LYS
1	A	38	ILE
1	A	46	SER
1	A	55	SER
1	A	58	ASP
1	A	60	ILE
1	A	62	SER
1	A	97	SER
1	A	100	ASP
1	A	107	LEU
1	A	132	ASN
1	A	138	LEU
1	A	153	LYS
1	A	170	SER
1	A	182	SER
1	A	183	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	SER
1	A	350	THR
1	A	364	SER
1	A	365	VAL
1	A	385	THR
1	A	489	THR
1	A	498	THR
1	B	35	ASN
1	B	67	GLU
1	B	70	LYS
1	B	86	THR
1	B	107	LEU
1	B	132	ASN
1	B	174	THR
1	B	183	GLU
1	B	198	VAL
1	B	224	LEU
1	B	295	LYS
1	B	299	LYS
1	B	303	ILE
1	B	340	GLU
1	B	365	VAL
1	B	384	THR
1	B	387	THR
1	B	424	THR
1	B	436	THR
1	B	465	GLU
1	B	479	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	130	ASN
1	A	199	ASN
1	A	315	ASN
1	A	330	HIS
1	A	370	ASN
1	A	483	ASN
1	A	500	ASN
1	B	73	ASN
1	B	108	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	315	ASN
1	B	348	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EPE	A	601	-	15,15,15	1.14	1 (6%)	19,20,20	0.50	0
2	EPE	B	601	-	15,15,15	0.89	1 (6%)	19,20,20	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	A	601	-	-	3/9/19/19	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	B	601	-	-	5/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	EPE	C10-S	-4.32	1.71	1.77
2	B	601	EPE	C10-S	-3.29	1.73	1.77

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	EPE	C10-C9-N1-C2
2	B	601	EPE	C9-C10-S-O2S
2	A	601	EPE	C9-C10-S-O3S
2	B	601	EPE	C9-C10-S-O3S
2	B	601	EPE	N4-C7-C8-O8
2	A	601	EPE	C9-C10-S-O1S
2	A	601	EPE	C9-C10-S-O2S
2	B	601	EPE	C9-C10-S-O1S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	467/472 (98%)	0.36	23 (4%) 35 30	24, 56, 90, 109	1 (0%)
1	B	467/472 (98%)	0.39	20 (4%) 40 35	30, 57, 92, 121	1 (0%)
All	All	934/944 (98%)	0.38	43 (4%) 37 33	24, 56, 91, 121	2 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	60	ILE	7.8
1	A	139	LYS	5.6
1	B	304	LYS	4.7
1	A	138	LEU	4.1
1	A	303	ILE	4.0
1	A	58	ASP	3.9
1	A	34	LYS	3.9
1	B	34	LYS	3.9
1	B	491	LEU	3.6
1	B	173	HIS	3.3
1	A	137	GLY	3.3
1	A	171	SER	3.3
1	B	303	ILE	3.2
1	A	141	GLU	3.2
1	A	172	THR	3.1
1	A	61	GLU	3.1
1	B	358	TYR	2.9
1	A	59	LYS	2.9
1	B	35	ASN	2.8
1	A	492	TYR	2.7
1	A	383	GLU	2.6
1	A	60	ILE	2.6
1	A	197	TRP	2.5
1	A	324	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	63	PHE	2.5
1	A	491	LEU	2.4
1	B	384	THR	2.4
1	A	325	GLY	2.4
1	B	419	GLY	2.4
1	A	304	LYS	2.3
1	A	57	GLY	2.3
1	B	174	THR	2.3
1	B	64	VAL	2.3
1	A	460	THR	2.2
1	B	144	ILE	2.2
1	B	197	TRP	2.1
1	B	351	PHE	2.1
1	B	385	THR	2.1
1	A	44	SER	2.1
1	A	461	GLY	2.1
1	B	325	GLY	2.1
1	B	307	GLN	2.0
1	B	308	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EPE	A	601	15/15	0.88	0.17	81,83,93,94	0
2	EPE	B	601	15/15	0.89	0.13	69,75,94,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.