



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:30 PM UTC

PDB ID : 6DIG / pdb_00006dig
Title : Crystal structure of DQA1*01:02/DQB1*06:02 in complex with a hypocretin peptide
Authors : Birtley, J.R.; Stern, L.J.; Mellins, E.D.; Jiang, W.
Deposited on : 2018-05-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	4% 85% 12%
2	B	215	7% 73% 10% 16%
3	C	13	15% 92% 8%
4	D	2	100%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class II HLA-DQ-alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1458	938	237	278	5			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	197	GLY	-	expression tag	UNP Q30066
A	198	GLY	-	expression tag	UNP Q30066
A	199	GLY	-	expression tag	UNP Q30066
A	200	GLY	-	expression tag	UNP Q30066
A	201	SER	-	expression tag	UNP Q30066
A	202	LEU	-	expression tag	UNP Q30066
A	203	GLU	-	expression tag	UNP Q30066
A	204	VAL	-	expression tag	UNP Q30066
A	205	LEU	-	expression tag	UNP Q30066
A	206	PHE	-	expression tag	UNP Q30066
A	207	GLN	-	expression tag	UNP Q30066

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DQ beta 1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	180	Total	C	N	O	S	0	2	0
			1493	944	263	279	7			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP Q5SU54
B	-2	SER	-	expression tag	UNP Q5SU54
B	-1	GLY	-	expression tag	UNP Q5SU54
B	0	GLY	-	expression tag	UNP Q5SU54
B	1	GLY	-	expression tag	UNP Q5SU54

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	GLY	-	expression tag	UNP Q5SU54
B	199	GLY	-	expression tag	UNP Q5SU54
B	200	THR	-	expression tag	UNP Q5SU54
B	201	GLY	-	expression tag	UNP Q5SU54
B	202	GLY	-	expression tag	UNP Q5SU54
B	203	GLY	-	expression tag	UNP Q5SU54
B	204	GLY	-	expression tag	UNP Q5SU54
B	205	SER	-	expression tag	UNP Q5SU54
B	206	LEU	-	expression tag	UNP Q5SU54
B	207	GLU	-	expression tag	UNP Q5SU54
B	208	VAL	-	expression tag	UNP Q5SU54
B	209	LEU	-	expression tag	UNP Q5SU54
B	210	PHE	-	expression tag	UNP Q5SU54
B	211	GLN	-	expression tag	UNP Q5SU54

- Molecule 3 is a protein called 13-mer peptide: ALA-GLY-ASN-HIS-ALA-ALA-GLY-ILE-L
EU-THR-LEU-GLY-LYS.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	0	0
			85	53	17	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a
cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:
C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	162	Total 162	O 162	0	0
7	B	104	Total 104	O 104	0	0
7	C	6	Total 6	O 6	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MHC class II HLA-DQ-alpha chain



- Molecule 2: HLA class II histocompatibility antigen, DQ beta 1 chain



- Molecule 3: 13-mer peptide: ALA-GLY-ASN-HIS-ALA-ALA-GLY-ILE-LEU-THR-LEU-GLY-LYS



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	63.67Å 89.60Å 90.93Å 90.00° 99.52° 90.00°	Depositor
Resolution (Å)	44.84 – 2.00 44.84 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.84-2.00) 84.9 (44.84-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.82 (at 2.00Å)	Xtriage
Refinement program	phenix.refine 1.13_2998, PHENIX 1.13_2998	Depositor
R, R_{free}	0.216 , 0.255 0.216 , 0.255	Depositor DCC
R_{free} test set	1711 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3372	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1503	0.52	0/2050
2	B	0.40	0/1536	0.55	0/2088
3	C	0.40	0/85	0.56	0/113
All	All	0.38	0/3124	0.54	0/4251

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1458	0	1377	10	0
2	B	1493	0	1433	17	0
3	C	85	0	92	1	0
4	D	28	0	25	0	0
5	A	28	0	26	4	0
6	A	8	0	12	0	0
7	A	162	0	0	1	1
7	B	104	0	0	3	1
7	C	6	0	0	0	0
All	All	3372	0	2965	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ASN:HD21	5:A:1001:NAG:C1	1.27	1.44
1:A:81:ASN:ND2	5:A:1001:NAG:C1	1.86	1.35
1:A:81:ASN:CG	5:A:1001:NAG:C1	2.36	0.98
1:A:81:ASN:OD1	5:A:1001:NAG:C1	2.36	0.72
2:B:176:GLU:OE2	7:B:1101:HOH:O	2.13	0.66
1:A:79:ARG:HE	3:C:13:LYS:HD2	1.62	0.65
1:A:114:ASN:ND2	7:A:1102:HOH:O	2.30	0.59
1:A:99:PRO:HG2	2:B:100:THR:HG21	1.83	0.59
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.87	0.55
2:B:143:VAL:HB	2:B:160:MET:HE3	1.88	0.55
2:B:21:THR:HG22	7:B:1173:HOH:O	2.05	0.55
1:A:50:GLU:OE2	2:B:93:ARG:NH1	2.42	0.52
1:A:147:SER:HB3	2:B:34:ARG:HH11	1.76	0.51
2:B:40:PHE:HB2	2:B:47:TYR:CE2	2.45	0.51
2:B:22:GLU:HG2	2:B:23:ARG:HG2	1.94	0.48
2:B:133:ARG:NH1	2:B:169:ASP:OD2	2.46	0.48
2:B:116:VAL:HG22	2:B:160:MET:HG2	1.95	0.47
2:B:117:CYS:HB2	2:B:131:TRP:CZ2	2.50	0.46
2:B:177:HIS:CD2	2:B:178:PRO:HD2	2.52	0.45
2:B:22:GLU:HG2	2:B:23:ARG:CG	2.48	0.43
2:B:68:LEU:HD12	2:B:68:LEU:HA	1.89	0.43
1:A:51:PHE:CZ	2:B:153:TRP:CD2	3.07	0.42
2:B:100:THR:HG22	7:B:1134:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1239:HOH:O	7:B:1171:HOH:O[4_446]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/207 (87%)	177 (98%)	3 (2%)	0	100	100
2	B	178/215 (83%)	176 (99%)	2 (1%)	0	100	100
3	C	11/13 (85%)	11 (100%)	0	0	100	100
All	All	369/435 (85%)	364 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/182 (89%)	162 (100%)	0	100	100
2	B	165/186 (89%)	165 (100%)	0	100	100
3	C	7/7 (100%)	7 (100%)	0	100	100
All	All	334/375 (89%)	334 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
2	B	136	GLN
2	B	150	ASN
2	B	156	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	2,4	14,14,15	2.70	7 (50%)	17,19,21	2.75	4 (23%)
4	NAG	D	2	4	14,14,15	2.34	5 (35%)	17,19,21	1.57	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	NAG	O5-C1	5.62	1.53	1.43
4	D	2	NAG	O5-C1	5.07	1.52	1.43
4	D	1	NAG	O5-C5	4.13	1.51	1.43
4	D	1	NAG	C7-N2	4.08	1.47	1.34
4	D	2	NAG	C7-N2	3.88	1.46	1.34
4	D	2	NAG	O5-C5	3.51	1.50	1.43
4	D	1	NAG	O4-C4	2.70	1.49	1.43
4	D	2	NAG	C2-N2	2.67	1.50	1.46
4	D	2	NAG	C1-C2	2.63	1.55	1.52
4	D	1	NAG	C2-N2	2.60	1.50	1.46
4	D	1	NAG	C1-C2	2.52	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	NAG	O7-C7	-2.05	1.18	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	O5-C1-C2	-8.78	97.70	111.29
4	D	1	NAG	C1-O5-C5	4.54	118.27	112.19
4	D	2	NAG	C1-O5-C5	3.95	117.48	112.19
4	D	1	NAG	C3-C4-C5	-2.33	106.02	110.23
4	D	2	NAG	C8-C7-N2	2.28	119.89	116.12
4	D	1	NAG	C1-C2-N2	-2.24	106.90	110.43
4	D	2	NAG	C6-C5-C4	-2.12	107.82	113.02


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	1002	1	14,14,15	1.94	4 (28%)	17,19,21	1.58	3 (17%)
6	TRS	A	1003	-	7,7,7	0.58	0	9,9,9	0.75	0
5	NAG	A	1001	-	14,14,15	2.57	7 (50%)	17,19,21	2.00	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1002	1	-	1/6/23/26	0/1/1/1
6	TRS	A	1003	-	-	3/9/9/9	-
5	NAG	A	1001	-	-	2/6/23/26	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1001	NAG	O5-C1	4.87	1.51	1.43
5	A	1001	NAG	C7-N2	4.54	1.49	1.34
5	A	1002	NAG	O5-C1	4.49	1.51	1.43
5	A	1001	NAG	C1-C2	4.19	1.58	1.52
5	A	1002	NAG	C7-N2	3.62	1.46	1.34
5	A	1001	NAG	C2-N2	3.27	1.51	1.46
5	A	1001	NAG	C8-C7	2.62	1.56	1.50
5	A	1001	NAG	C3-C2	-2.31	1.47	1.52
5	A	1002	NAG	O5-C5	2.14	1.47	1.43
5	A	1001	NAG	O5-C5	2.10	1.47	1.43
5	A	1002	NAG	C2-N2	2.07	1.49	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1001	NAG	C1-C2-N2	5.86	119.67	110.43
5	A	1002	NAG	C1-C2-N2	-4.00	104.13	110.43
5	A	1001	NAG	C2-N2-C7	3.42	127.48	122.90
5	A	1001	NAG	C4-C3-C2	2.69	114.97	111.02
5	A	1002	NAG	C8-C7-N2	2.22	119.81	116.12
5	A	1002	NAG	O5-C1-C2	2.13	114.58	111.29

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1002	NAG	O5-C5-C6-O6
6	A	1003	TRS	C1-C-C3-O3
6	A	1003	TRS	N-C-C3-O3
5	A	1001	NAG	C1-C2-N2-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	1003	TRS	C2-C-C3-O3
5	A	1001	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1001	NAG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/207 (87%)	0.24	8 (4%) 39 38	19, 33, 58, 87	0
2	B	180/215 (83%)	0.55	16 (8%) 15 14	18, 39, 75, 101	2 (1%)
3	C	13/13 (100%)	0.59	2 (15%) 5 4	23, 27, 62, 70	0
All	All	375/435 (86%)	0.40	26 (6%) 23 21	18, 35, 68, 101	2 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	1	ALA	4.5
2	B	168	GLY	3.4
2	B	165	PRO	3.1
1	A	183	PRO	3.1
2	B	136	GLN	3.0
2	B	164	THR	2.9
1	A	81	ASN	2.9
2	B	21	THR	2.9
2	B	160	MET	2.9
2	B	104	SER	2.9
1	A	2	ASP	2.6
3	C	13	LYS	2.6
2	B	169	ASP	2.6
2	B	55	ARG	2.6
2	B	19	ASN	2.5
2	B	100	THR	2.4
2	B	190	ALA	2.3
1	A	3	ILE	2.3
2	B	166	GLN	2.3
2	B	113	ASN	2.2
1	A	132	THR	2.2
2	B	3	SER	2.1
1	A	133	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	23	ARG	2.1
1	A	161	ASP	2.0
1	A	175	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	D	2	14/15	0.58	0.18	97,100,102,102	0
4	NAG	D	1	14/15	0.63	0.20	75,80,89,93	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	1001	14/15	0.46	0.20	71,82,84,86	0
5	NAG	A	1002	14/15	0.71	0.20	66,72,80,83	0
6	TRS	A	1003	8/8	0.76	0.22	63,64,64,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.