



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 11:56 PM UTC

PDB ID : 1DJW / pdb_00001djw
Title : PHOSPHOINOSITIDE-SPECIFIC PHOSPHOLIPASE C-DELTA1 FROM
RAT COMPLEXED WITH INOSITOL-2-METHYLENE-1,2-CYCLIC-MON
OPHOSPHONATE
Authors : Essen, L.-O.; Perisic, O.; Williams, R.L.
Deposited on : 1996-08-24
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	624	 4% 45% 31% 5% 18%
1	B	624	 4% 50% 33% 6% 10%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOINOSITIDE-SPECIFIC PHOSPHOLIPASE C, ISOZYME DELTA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	513	Total	C	N	O	S	89	0	0
			4057	2565	709	761	22			
1	B	561	Total	C	N	O	S	78	0	0
			4465	2818	776	847	24			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		
2	B	3	Total	Ca	0	0
			3	3		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is INOSITOL-2-METHYLENE-1,2-CYCLIC-MONOPHOSPHATE (CCD ID: CIP) (formula: $C_7H_{13}O_7P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			15	7	7	1		
4	B	1	Total	C	O	P	0	0
			15	7	7	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	327	Total	O	0	0
			327	327		
5	B	425	Total	O	0	0
			425	425		



4 Data and refinement statistics

Property	Value	Source
Space group	F 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	397.03Å 397.03Å 397.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.45 10.00 – 2.45	Depositor EDS
% Data completeness (in resolution range)	96.0 (10.00-2.45) 96.4 (10.00-2.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.31Å)	Xtriage
Refinement program	TNT 5E	Depositor
R, R_{free}	0.217 , 0.270 0.208 , 0.252	Depositor DCC
R_{free} test set	4692 reflections (4.08%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 114.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9318	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACT, CIP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	4/4152 (0.1%)	1.14	25/5624 (0.4%)
1	B	0.89	0/4565	1.13	27/6174 (0.4%)
All	All	0.92	4/8717 (0.0%)	1.13	52/11798 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	277	MET	SD-CE	7.81	1.99	1.79
1	A	231	GLU	CD-OE1	-6.92	1.12	1.25
1	A	231	GLU	CA-CB	-5.07	1.45	1.53
1	A	702	PHE	CE2-CZ	-5.02	1.23	1.38

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	508	SER	N-CA-C	13.01	127.27	112.57
1	A	420	GLY	N-CA-C	-9.41	97.21	111.24
1	A	208	LEU	N-CA-C	-9.17	101.26	111.07
1	B	594	GLN	N-CA-C	-8.84	102.28	113.23
1	B	509	SER	N-CA-C	-8.74	101.50	113.57

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	238	GLN	Sidechain
1	A	580	GLN	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4057	0	3972	180	1
1	B	4465	0	4375	239	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	15	0	12	4	0
4	B	15	0	12	4	0
5	A	327	0	0	25	1
5	B	425	0	0	19	0
All	All	9318	0	8377	413	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 413 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:504:PHE:HB3	1:A:527:ARG:HH22	1.18	1.06
1:B:504:PHE:HB3	1:B:527:ARG:HH22	1.20	1.01
1:B:238:GLN:HG2	1:B:246:ALA:HB1	1.45	0.97
1:B:644:VAL:HG23	1:B:645:ASN:H	1.29	0.96
1:A:624:PRO:HD2	1:A:625:TRP:CE3	2.09	0.86

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:GLU:OE1	5:A:896:HOH:O[52_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	509/624 (82%)	472 (93%)	35 (7%)	2 (0%)	30	37
1	B	557/624 (89%)	511 (92%)	43 (8%)	3 (0%)	24	32
All	All	1066/1248 (85%)	983 (92%)	78 (7%)	5 (0%)	24	32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	173	ASN
1	B	645	ASN
1	B	647	ASN
1	A	421	VAL
1	A	744	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/545 (82%)	388 (87%)	56 (13%)	4	4
1	B	492/545 (90%)	438 (89%)	54 (11%)	6	5
All	All	936/1090 (86%)	826 (88%)	110 (12%)	5	4

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	160	MET
1	B	242	ARG
1	B	753	SER
1	B	646	LYS
1	B	169	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	573	GLN
1	B	728	GLN
1	B	734	HIS
1	B	718	GLN
1	A	734	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIP	A	1	2	14,16,16	1.85	3 (21%)	18,26,26	2.11	9 (50%)
3	ACT	A	5	-	3,3,3	1.01	0	3,3,3	0.65	0
3	ACT	B	5	-	3,3,3	0.95	0	3,3,3	0.85	0
4	CIP	B	1	2	14,16,16	2.02	3 (21%)	18,26,26	2.24	8 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIP	A	1	2	-	-	0/2/2/2
4	CIP	B	1	2	-	-	0/2/2/2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1	CIP	P-O1P	4.92	1.63	1.51
4	B	1	CIP	P-O1P	4.78	1.62	1.51
4	B	1	CIP	P-C7	-4.40	1.76	1.82
4	A	1	CIP	P-O2P	2.70	1.63	1.56
4	A	1	CIP	P-C7	-2.59	1.79	1.82

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	CIP	O4-C4-C3	-4.34	100.14	110.38
4	B	1	CIP	O1P-P-C7	4.10	118.86	109.29
4	B	1	CIP	O6-C6-C5	-3.33	102.53	110.38
4	B	1	CIP	C6-C5-C4	-3.29	105.06	110.83
4	A	1	CIP	O6-C6-C5	-3.28	102.65	110.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	CIP	4	0
4	B	1	CIP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	507/624 (81%)	-0.18	25 (4%) 35 33	14, 36, 81, 115	14 (2%)
1	B	558/624 (89%)	-0.18	28 (5%) 34 32	16, 37, 77, 108	16 (2%)
All	All	1065/1248 (85%)	-0.18	53 (4%) 34 32	14, 36, 78, 115	30 (2%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	202	GLU	6.6
1	B	648	LYS	6.2
1	B	514	GLY	5.8
1	A	203	THR	5.4
1	A	206	LYS	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CIP	A	1	15/15	0.72	0.22	42,49,55,56	15
4	CIP	B	1	15/15	0.74	0.22	43,49,55,55	15
2	CA	A	4	1/1	0.78	0.13	58,58,58,58	1
2	CA	B	3	1/1	0.86	0.09	54,54,54,54	1
2	CA	A	2	1/1	0.87	0.08	50,50,50,50	0
3	ACT	A	5	4/4	0.93	0.07	35,37,38,41	0
2	CA	A	3	1/1	0.96	0.12	56,56,56,56	1
2	CA	B	2	1/1	0.96	0.07	46,46,46,46	0
2	CA	B	4	1/1	0.97	0.11	59,59,59,59	1
3	ACT	B	5	4/4	0.98	0.05	23,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.