



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:23 PM UTC

PDB ID : 5DJC / pdb_00005djc
Title : Fc Heterodimer Design 8.1 L368V/Y407A + T366V/K409F
Authors : Atwell, S.; Leaver-Fay, A.; Froning, K.J.; Aldaz, H.; Pustilnik, A.; Lu, F.;
Huang, F.; Yuan, R.; Dhanani, S.H.; Chamberlain, A.K.; Fitchett, J.R.;
Gutierrez, B.; Hendle, J.; Secrist, E.; Demarest, S.J.; Kuhlman, B.
Deposited on : 2015-09-01
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	81% 10% 9%
1	D	227	81% 8% 9%
2	B	240	8% 82% 14%
2	E	240	8% 75% 11% 14%
3	C	13	92% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 100%
4	G	8	 75% 25%
4	H	8	 38% 62%
4	I	8	 50% 38% 12%
4	J	8	 62% 25% 12%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	0
			1637	1041	273	316	7			
1	D	207	Total	C	N	O	S	1	0	0
			1635	1042	270	316	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	GLU	ASP	variant	UNP P01857
A	358	MET	LEU	variant	UNP P01857
A	368	VAL	LEU	engineered mutation	UNP P01857
A	407	ALA	TYR	engineered mutation	UNP P01857
D	356	GLU	ASP	variant	UNP P01857
D	358	MET	LEU	variant	UNP P01857
D	368	VAL	LEU	engineered mutation	UNP P01857
D	407	ALA	TYR	engineered mutation	UNP P01857

- Molecule 2 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	206	Total	C	N	O	S	0	0	0
			1587	1020	261	300	6			
2	E	207	Total	C	N	O	S	0	0	0
			1580	1014	260	300	6			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	208	HIS	-	expression tag	UNP P01857
B	209	HIS	-	expression tag	UNP P01857
B	210	HIS	-	expression tag	UNP P01857
B	211	HIS	-	expression tag	UNP P01857

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	212	HIS	-	expression tag	UNP P01857
B	213	HIS	-	expression tag	UNP P01857
B	214	HIS	-	expression tag	UNP P01857
B	215	HIS	-	expression tag	UNP P01857
B	216	SER	-	expression tag	UNP P01857
B	217	GLY	-	expression tag	UNP P01857
B	218	SER	-	expression tag	UNP P01857
B	219	GLY	-	expression tag	UNP P01857
B	220	SER	-	expression tag	UNP P01857
B	252	GLU	MET	engineered mutation	UNP P01857
B	253	ALA	ILE	engineered mutation	UNP P01857
B	356	GLU	ASP	variant	UNP P01857
B	358	MET	LEU	variant	UNP P01857
B	366	VAL	THR	engineered mutation	UNP P01857
B	409	PHE	LYS	engineered mutation	UNP P01857
B	435	ALA	HIS	engineered mutation	UNP P01857
E	208	HIS	-	expression tag	UNP P01857
E	209	HIS	-	expression tag	UNP P01857
E	210	HIS	-	expression tag	UNP P01857
E	211	HIS	-	expression tag	UNP P01857
E	212	HIS	-	expression tag	UNP P01857
E	213	HIS	-	expression tag	UNP P01857
E	214	HIS	-	expression tag	UNP P01857
E	215	HIS	-	expression tag	UNP P01857
E	216	SER	-	expression tag	UNP P01857
E	217	GLY	-	expression tag	UNP P01857
E	218	SER	-	expression tag	UNP P01857
E	219	GLY	-	expression tag	UNP P01857
E	220	SER	-	expression tag	UNP P01857
E	252	GLU	MET	engineered mutation	UNP P01857
E	253	ALA	ILE	engineered mutation	UNP P01857
E	356	GLU	ASP	variant	UNP P01857
E	358	MET	LEU	variant	UNP P01857
E	366	VAL	THR	engineered mutation	UNP P01857
E	409	PHE	LYS	engineered mutation	UNP P01857
E	435	ALA	HIS	engineered mutation	UNP P01857

- Molecule 3 is a protein called Fc-III peptide.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	S	0	0	0
			107	69	17	19	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	13	Total	C	N	O	S	0	0	0
			107	69	17	19	2			

- Molecule 4 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	8	Total	C	N	O			
			96	54	3	39	0	0	0
4	H	8	Total	C	N	O			
			96	54	3	39	0	0	0
4	I	8	Total	C	N	O			
			96	54	3	39	0	0	0
4	J	8	Total	C	N	O			
			96	54	3	39	0	0	0

- Molecule 5 is IODIDE ION (CCD ID: IOD) (formula: I).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total I 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	64	Total O 64 64	0	0
6	B	25	Total O 25 25	0	0
6	C	1	Total O 1 1	0	0
6	D	67	Total O 67 67	0	0
6	E	32	Total O 32 32	0	0

- Molecule 1: Ig gamma-1 chain C region





- Molecule 3: Fc-III peptide

Chain C: 92% 8%



- Molecule 3: Fc-III peptide

Chain F: 100%

There are no outlier residues recorded for this chain.

- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 75% 25%



- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 38% 62%



- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 50% 38% 12%



- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.24Å 66.86Å 72.77Å 85.17° 80.71° 89.69°	Depositor
Resolution (Å)	30.00 – 2.10 30.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.3 (30.00-2.10) 96.3 (30.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0017	Depositor
R, R_{free}	0.229 , 0.285 (Not available) , 0.287	Depositor DCC
R_{free} test set	3179 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7227	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, IOD, FUC, BMA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/1682	0.76	1/2295 (0.0%)
1	D	0.71	1/1680 (0.1%)	0.83	1/2293 (0.0%)
2	B	0.64	0/1631	0.76	0/2230
2	E	0.61	0/1624	0.74	0/2224
3	C	1.02	0/111	0.66	0/151
3	F	1.05	0/111	0.57	0/151
All	All	0.68	1/6839 (0.0%)	0.77	2/9344 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	360	LYS	CG-CD	-8.26	1.27	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	360	LYS	CB-CG-CD	14.51	144.66	111.30
1	A	303	VAL	N-CA-C	5.95	116.58	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1637	0	1583	10	0
1	D	1635	0	1570	14	0
2	B	1587	0	1501	4	0
2	E	1580	0	1497	13	0
3	C	107	0	93	0	0
3	F	107	0	93	0	0
4	G	96	0	82	0	0
4	H	96	0	82	0	0
4	I	96	0	82	2	0
4	J	96	0	82	1	0
5	D	1	0	0	0	0
6	A	64	0	0	0	0
6	B	25	0	0	0	0
6	C	1	0	0	0	0
6	D	67	0	0	1	0
6	E	32	0	0	1	0
All	All	7227	0	6665	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.71	0.72
1:D:297:ASN:O	1:D:298:SER:CB	2.43	0.67
1:A:353:PRO:HD3	1:A:365:LEU:HD23	1.83	0.59
2:E:290:LYS:HE3	2:E:305:VAL:HG21	1.84	0.58
2:B:242:LEU:HG	2:B:336:ILE:HG12	1.87	0.57
2:E:320:LYS:HD3	2:E:333:GLU:OE2	2.05	0.56
2:E:322:LYS:HG3	2:E:333:GLU:HG2	1.90	0.54
1:D:276:ASN:HB2	1:D:322:LYS:HB3	1.92	0.52
1:D:346:PRO:HB3	1:D:372:PHE:HB3	1.92	0.50
1:A:245:PRO:HD3	1:A:259:VAL:HG12	1.93	0.50
1:A:246:LYS:O	1:A:247:PRO:C	2.55	0.50
1:D:297:ASN:O	1:D:298:SER:HB3	2.12	0.49
1:D:443:LEU:C	6:D:644:HOH:O	2.54	0.49
1:D:318:GLU:HA	1:D:337:SER:HB3	1.95	0.49
2:B:274:LYS:HB3	2:B:324:SER:HB2	1.93	0.49
1:D:351:LEU:HB2	1:D:366:THR:HB	1.95	0.48
2:E:358:MET:HE1	2:E:417:TRP:HE1	1.78	0.47
2:E:362:GLN:HG2	2:E:413:ASP:HA	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:424:SER:OG	1:A:440:SER:OG	2.31	0.47
2:E:246:LYS:HG2	4:J:6:GAL:O4	2.16	0.46
2:E:353:PRO:HA	6:E:616:HOH:O	2.16	0.46
1:A:406:LEU:C	1:A:406:LEU:HD12	2.41	0.45
2:E:261:CYS:HB2	2:E:277:TRP:CZ2	2.52	0.45
2:E:367:CYS:HB2	2:E:381:TRP:CZ2	2.51	0.45
1:D:238:PRO:HD2	1:D:328:LEU:HD13	1.99	0.44
2:E:379:VAL:HG21	2:E:406:LEU:HD11	1.99	0.44
1:A:278:TYR:HB2	1:A:320:LYS:HB3	1.99	0.44
2:E:366:VAL:HG22	2:E:409:PHE:CD2	2.53	0.44
2:E:358:MET:HE1	2:E:417:TRP:NE1	2.33	0.43
2:B:238:PRO:HA	2:B:265:ASP:HB2	2.00	0.43
1:D:242:LEU:HG	1:D:336:ILE:HG12	2.00	0.43
1:D:295:GLN:OE1	4:I:1:NAG:H62	2.18	0.42
1:D:297:ASN:O	1:D:298:SER:HB2	2.19	0.42
1:A:238:PRO:HD2	1:A:328:LEU:HD13	2.01	0.42
1:D:405:PHE:CD1	1:D:405:PHE:C	2.97	0.42
1:A:322:LYS:HG3	1:A:333:GLU:HG2	2.02	0.41
2:E:373:TYR:CG	2:E:374:PRO:HA	2.55	0.41
2:B:391:TYR:HB3	2:B:410:LEU:HD12	2.03	0.41
1:A:257:PRO:HG2	1:A:308:VAL:O	2.21	0.40
1:D:356:GLU:O	1:D:359:THR:HG22	2.22	0.40
1:D:246:LYS:HG2	4:I:6:GAL:O4	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/227 (90%)	201 (98%)	4 (2%)	0	100	100
1	D	205/227 (90%)	200 (98%)	4 (2%)	1 (0%)	24	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	202/240 (84%)	195 (96%)	7 (4%)	0	100	100
2	E	205/240 (85%)	201 (98%)	4 (2%)	0	100	100
3	C	11/13 (85%)	10 (91%)	1 (9%)	0	100	100
3	F	11/13 (85%)	10 (91%)	1 (9%)	0	100	100
All	All	839/960 (87%)	817 (97%)	21 (2%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	298	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/209 (90%)	185 (98%)	3 (2%)	55	64
1	D	185/209 (88%)	182 (98%)	3 (2%)	55	64
2	B	172/219 (78%)	169 (98%)	3 (2%)	53	62
2	E	172/219 (78%)	167 (97%)	5 (3%)	37	42
3	C	11/11 (100%)	10 (91%)	1 (9%)	9	6
3	F	11/11 (100%)	11 (100%)	0	100	100
All	All	739/878 (84%)	724 (98%)	15 (2%)	48	56

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	293	GLU
1	A	336	ILE
1	A	359	THR
2	B	336	ILE
2	B	370	LYS
2	B	440	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	8	GLU
1	D	328	LEU
1	D	336	ILE
1	D	418	GLN
2	E	288	LYS
2	E	289	THR
2	E	328	LEU
2	E	382	GLU
2	E	443	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	311	GLN
1	A	347	GLN
1	A	362	GLN
1	A	438	GLN
2	B	285	HIS
2	B	362	GLN
2	B	418	GLN
1	D	268	HIS
2	E	389	ASN
2	E	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

32 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	1	4,1	14,14,15	0.60	0	17,19,21	0.71	0
4	NAG	G	2	4	14,14,15	0.51	0	17,19,21	0.79	0
4	BMA	G	3	4	11,11,12	0.32	0	15,15,17	1.19	1 (6%)
4	MAN	G	4	4	11,11,12	0.60	0	15,15,17	1.20	2 (13%)
4	NAG	G	5	4	14,14,15	0.57	0	17,19,21	0.73	0
4	GAL	G	6	4	11,11,12	0.55	0	15,15,17	0.63	0
4	MAN	G	7	4	11,11,12	0.59	0	15,15,17	0.66	0
4	FUC	G	8	4	10,10,11	0.58	0	14,14,16	0.96	0
4	NAG	H	1	4,2	14,14,15	0.54	0	17,19,21	0.84	0
4	NAG	H	2	4	14,14,15	0.48	0	17,19,21	1.02	2 (11%)
4	BMA	H	3	4	11,11,12	0.58	0	15,15,17	1.55	3 (20%)
4	MAN	H	4	4	11,11,12	0.63	0	15,15,17	1.23	2 (13%)
4	NAG	H	5	4	14,14,15	0.61	0	17,19,21	1.03	1 (5%)
4	GAL	H	6	4	11,11,12	0.63	0	15,15,17	0.91	0
4	MAN	H	7	4	11,11,12	0.53	0	15,15,17	2.11	5 (33%)
4	FUC	H	8	4	10,10,11	0.63	0	14,14,16	0.57	0
4	NAG	I	1	4,1	14,14,15	0.60	0	17,19,21	0.74	0
4	NAG	I	2	4	14,14,15	0.65	0	17,19,21	1.00	0
4	BMA	I	3	4	11,11,12	0.26	0	15,15,17	1.05	1 (6%)
4	MAN	I	4	4	11,11,12	0.58	0	15,15,17	0.89	1 (6%)
4	NAG	I	5	4	14,14,15	0.67	0	17,19,21	0.78	0
4	GAL	I	6	4	11,11,12	0.64	0	15,15,17	1.03	1 (6%)
4	MAN	I	7	4	11,11,12	0.65	0	15,15,17	0.69	0
4	FUC	I	8	4	10,10,11	0.80	0	14,14,16	0.71	0
4	NAG	J	1	4,2	14,14,15	0.55	0	17,19,21	0.67	0
4	NAG	J	2	4	14,14,15	0.47	0	17,19,21	1.03	0
4	BMA	J	3	4	11,11,12	0.49	0	15,15,17	1.05	2 (13%)
4	MAN	J	4	4	11,11,12	0.62	0	15,15,17	0.82	0
4	NAG	J	5	4	14,14,15	0.59	0	17,19,21	0.80	0
4	GAL	J	6	4	11,11,12	0.61	0	15,15,17	0.91	1 (6%)
4	MAN	J	7	4	11,11,12	0.56	0	15,15,17	0.96	1 (6%)
4	FUC	J	8	4	10,10,11	0.65	0	14,14,16	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
4	NAG	G	5	4	-	0/6/23/26	0/1/1/1
4	GAL	G	6	4	-	0/2/19/22	0/1/1/1
4	MAN	G	7	4	-	2/2/19/22	0/1/1/1
4	FUC	G	8	4	-	-	0/1/1/1
4	NAG	H	1	4,2	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
4	MAN	H	4	4	-	0/2/19/22	0/1/1/1
4	NAG	H	5	4	-	0/6/23/26	0/1/1/1
4	GAL	H	6	4	-	0/2/19/22	0/1/1/1
4	MAN	H	7	4	-	1/2/19/22	0/1/1/1
4	FUC	H	8	4	-	-	0/1/1/1
4	NAG	I	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	I	2	4	-	0/6/23/26	0/1/1/1
4	BMA	I	3	4	-	0/2/19/22	0/1/1/1
4	MAN	I	4	4	-	0/2/19/22	0/1/1/1
4	NAG	I	5	4	-	0/6/23/26	0/1/1/1
4	GAL	I	6	4	-	2/2/19/22	0/1/1/1
4	MAN	I	7	4	-	2/2/19/22	0/1/1/1
4	FUC	I	8	4	-	-	0/1/1/1
4	NAG	J	1	4,2	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
4	BMA	J	3	4	-	0/2/19/22	0/1/1/1
4	MAN	J	4	4	-	0/2/19/22	0/1/1/1
4	NAG	J	5	4	-	0/6/23/26	0/1/1/1
4	GAL	J	6	4	-	0/2/19/22	0/1/1/1
4	MAN	J	7	4	-	0/2/19/22	0/1/1/1
4	FUC	J	8	4	-	-	0/1/1/1

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	7	MAN	C1-O5-C5	5.84	120.01	112.19
4	H	3	BMA	C1-C2-C3	3.78	115.15	109.64
4	G	4	MAN	C1-O5-C5	3.37	116.70	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	3	BMA	C1-O5-C5	3.28	116.58	112.19
4	H	3	BMA	C1-O5-C5	3.26	116.56	112.19
4	H	7	MAN	C3-C4-C5	3.12	115.89	110.23
4	I	3	BMA	C1-O5-C5	2.95	116.14	112.19
4	H	4	MAN	C1-O5-C5	2.65	115.73	112.19
4	J	3	BMA	C1-O5-C5	2.51	115.55	112.19
4	J	6	GAL	C1-C2-C3	2.50	113.28	109.64
4	H	7	MAN	O5-C1-C2	2.41	116.53	110.79
4	H	7	MAN	O5-C5-C6	2.38	112.29	107.66
4	H	5	NAG	C1-O5-C5	2.34	115.32	112.19
4	H	2	NAG	C2-N2-C7	-2.29	119.83	122.90
4	I	6	GAL	C1-C2-C3	2.24	112.90	109.64
4	J	3	BMA	C1-C2-C3	2.21	112.86	109.64
4	H	4	MAN	C3-C4-C5	2.20	114.23	110.23
4	I	4	MAN	C1-O5-C5	2.19	115.12	112.19
4	J	7	MAN	C1-O5-C5	2.14	115.05	112.19
4	G	4	MAN	O2-C2-C3	-2.13	105.74	110.15
4	H	7	MAN	C2-C3-C4	2.08	114.52	110.86
4	H	3	BMA	O6-C6-C5	-2.02	104.46	111.33
4	H	2	NAG	O5-C1-C2	-2.00	108.19	111.29

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	2	NAG	O5-C5-C6-O6
4	I	7	MAN	O5-C5-C6-O6
4	H	2	NAG	O5-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
4	I	7	MAN	C4-C5-C6-O6
4	H	2	NAG	C4-C5-C6-O6
4	G	7	MAN	C4-C5-C6-O6
4	I	6	GAL	C4-C5-C6-O6
4	G	7	MAN	O5-C5-C6-O6
4	I	6	GAL	O5-C5-C6-O6
4	H	7	MAN	C4-C5-C6-O6



There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	I	6	GAL	1	0
4	J	6	GAL	1	0
4	I	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/227 (91%)	0.03	1 (0%) 87 88	23, 36, 48, 67	0
1	D	207/227 (91%)	0.00	2 (0%) 79 81	23, 36, 48, 62	1 (0%)
2	B	206/240 (85%)	0.64	20 (9%) 13 14	26, 46, 87, 103	0
2	E	207/240 (86%)	0.60	18 (8%) 16 16	26, 48, 83, 98	0
3	C	13/13 (100%)	0.49	0 100 100	29, 40, 53, 70	0
3	F	13/13 (100%)	0.69	0 100 100	33, 45, 63, 68	0
All	All	853/960 (88%)	0.33	41 (4%) 35 38	23, 40, 76, 103	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	413	ASP	5.1
2	E	443	LEU	4.4
2	E	361	ASN	4.2
2	E	413	ASP	4.0
2	E	419	GLN	3.7
2	E	417	TRP	3.5
2	E	354	SER	3.4
2	B	354	SER	3.3
2	B	296	TYR	3.2
2	B	361	ASN	2.8
2	E	298	SER	2.8
2	E	415	SER	2.7
2	B	441	LEU	2.7
2	E	418	GLN	2.6
2	B	443	LEU	2.6
2	E	270	ASP	2.6
2	B	414	LYS	2.6
2	B	299	THR	2.5
2	B	251	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	270	ASP	2.5
2	B	421	ASN	2.5
2	B	419	GLN	2.4
2	B	389	ASN	2.4
2	E	416	ARG	2.3
2	E	422	VAL	2.3
2	B	412	VAL	2.2
2	E	412	VAL	2.2
1	A	441	LEU	2.2
2	E	359	THR	2.2
2	E	384	ASN	2.2
2	B	417	TRP	2.2
2	B	362	GLN	2.1
2	E	420	GLY	2.1
2	E	355	ARG	2.1
1	D	359	THR	2.1
2	B	433	HIS	2.1
2	B	371	GLY	2.1
2	B	295	GLN	2.0
2	B	241	PHE	2.0
1	D	443	LEU	2.0
2	E	296	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FUC	J	8	10/11	0.52	0.19	75,82,97,98	0
4	MAN	H	7	11/12	0.53	0.17	72,83,88,92	0
4	MAN	G	7	11/12	0.73	0.13	53,61,66,85	0
4	MAN	J	7	11/12	0.74	0.14	78,84,88,89	0
4	MAN	I	7	11/12	0.78	0.12	58,68,75,85	0
4	FUC	H	8	10/11	0.79	0.13	74,86,93,94	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GAL	I	6	11/12	0.84	0.13	46,52,65,76	0
4	BMA	H	3	11/12	0.84	0.10	48,59,65,70	0
4	GAL	G	6	11/12	0.84	0.17	43,63,75,77	0
4	NAG	H	1	14/15	0.84	0.12	58,64,78,78	0
4	NAG	H	2	14/15	0.86	0.12	47,59,72,73	0
4	MAN	H	4	11/12	0.86	0.12	42,63,70,72	0
4	GAL	H	6	11/12	0.86	0.15	56,62,73,75	0
4	MAN	J	4	11/12	0.87	0.13	50,62,75,76	0
4	NAG	J	1	14/15	0.88	0.10	53,61,79,82	0
4	FUC	G	8	10/11	0.88	0.10	47,50,59,60	0
4	BMA	J	3	11/12	0.89	0.10	51,57,67,71	0
4	NAG	J	2	14/15	0.89	0.10	43,58,65,69	0
4	FUC	I	8	10/11	0.90	0.09	36,43,45,48	0
4	GAL	J	6	11/12	0.90	0.12	50,61,69,72	0
4	NAG	G	5	14/15	0.91	0.09	33,41,49,60	0
4	BMA	G	3	11/12	0.91	0.08	35,40,49,54	0
4	NAG	J	5	14/15	0.91	0.09	44,51,60,62	0
4	NAG	I	2	14/15	0.92	0.08	31,38,47,51	0
4	NAG	G	1	14/15	0.92	0.10	37,43,55,57	0
4	NAG	I	1	14/15	0.92	0.10	34,40,44,51	0
4	NAG	G	2	14/15	0.93	0.09	38,43,53,54	0
4	MAN	G	4	11/12	0.93	0.08	30,41,50,51	0
4	BMA	I	3	11/12	0.93	0.07	41,46,53,61	0
4	NAG	I	5	14/15	0.93	0.08	27,40,52,55	0
4	NAG	H	5	14/15	0.93	0.09	46,54,70,70	0
4	MAN	I	4	11/12	0.94	0.08	31,44,56,58	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	IOD	D	509	1/1	0.99	0.03	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.