



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:09 PM UTC

PDB ID : 5DJQ / pdb_00005djq
Title : The structure of CBB3 cytochrome oxidase.
Authors : Buschmann, S.; Warkentin, E.; Xie, H.; Kohlstaedt, M.; Langer, J.D.; Ermler, U.; Michel, H.
Deposited on : 2015-09-02
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	% 61% 32% . .
1	D	474	58% 35% . .
1	G	474	% 59% 35% . .
1	K	474	% 60% 34% . .
2	B	203	63% 29% 6% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	203	
2	H	203	
2	L	203	
3	C	311	
3	F	311	
3	I	311	
3	M	311	
4	N	36	
4	O	36	
4	P	36	
4	Q	36	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	PO4	K	506	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 31974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cbb3-type cytochrome c oxidase subunit CcoN1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	466	Total	C	N	O	S	0	0	0
			3683	2461	593	607	22			
1	D	463	Total	C	N	O	S	0	0	0
			3663	2450	590	601	22			
1	G	465	Total	C	N	O	S	0	0	0
			3676	2457	592	605	22			
1	K	465	Total	C	N	O	S	0	0	0
			3676	2457	592	605	22			

- Molecule 2 is a protein called Cbb3-type cytochrome c oxidase subunit II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	197	Total	C	N	O	S	0	0	0
			1548	981	268	289	10			
2	E	197	Total	C	N	O	S	0	0	0
			1548	981	268	289	10			
2	H	197	Total	C	N	O	S	0	0	0
			1548	981	268	289	10			
2	L	197	Total	C	N	O	S	0	0	0
			1548	981	268	289	10			


- Molecule 3 is a protein called Cbb3-type cytochrome c oxidase subunit CcoP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	303	Total	C	N	O	S	0	0	0
			2312	1483	391	427	11			
3	F	303	Total	C	N	O	S	0	0	0
			2312	1483	391	427	11			
3	I	303	Total	C	N	O	S	0	0	0
			2312	1483	391	427	11			
3	M	303	Total	C	N	O	S	0	0	0
			2312	1483	391	427	11			

- Molecule 4 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	N	29	Total	C	N	O	S	0	0	0
			221	154	31	34	2			
4	O	29	Total	C	N	O	S	0	0	0
			221	154	31	34	2			
4	P	29	Total	C	N	O	S	0	0	0
			221	154	31	34	2			
4	Q	29	Total	C	N	O	S	0	0	0
			221	154	31	34	2			

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	G	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	G	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	K	1	Total	C	Fe	N	O	
			43	34	1	4	4	
5	K	1	Total	C	Fe	N	O	
			43	34	1	4	4	


- Molecule 6 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cu		
			1	1	0	0
6	D	1	Total	Cu		
			1	1	0	0
6	G	1	Total	Cu		
			1	1	0	0
6	K	1	Total	Cu		
			1	1	0	0

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Ca		
			2	2	0	0
7	D	2	Total	Ca		
			2	2	0	0
7	G	2	Total	Ca		
			2	2	0	0
7	K	2	Total	Ca		
			2	2	0	0

- Molecule 8 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H₂O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total O 2 2	0	0
8	D	1	Total O 2 2	0	0
8	G	1	Total O 2 2	0	0
8	K	1	Total O 2 2	0	0

- Molecule 9 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total 5	O 4	P 1	0	0
9	D	1	Total 5	O 4	P 1	0	0
9	G	1	Total 5	O 4	P 1	0	0
9	K	1	Total 5	O 4	P 1	0	0

- Molecule 10 is HEME C (CCD ID: HEC) (formula: $\text{C}_{34}\text{H}_{34}\text{FeN}_4\text{O}_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
10	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	H	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	I	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	I	1	Total	C	Fe	N	O	
			43	34	1	4	4	
10	L	1	Total	C	Fe	N	O	
			43	34	1	4	4	
10	M	1	Total	C	Fe	N	O	
			43	34	1	4	4	
10	M	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 11 is HEXACYANOFERRATE(3-) (CCD ID: FC6) (formula: C_6FeN_6).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	C	1	Total	C	Fe	N		
			13	6	1	6	0	0
11	F	1	Total	C	Fe	N		
			13	6	1	6	0	0
11	I	1	Total	C	Fe	N		
			13	6	1	6	0	0
11	M	1	Total	C	Fe	N		
			13	6	1	6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cbb3-type cytochrome c oxidase subunit CcoN1




- Molecule 1: Cbb3-type cytochrome c oxidase subunit CcoN1






Chain B: 




- Molecule 2: Cbb3-type cytochrome c oxidase subunit II

Chain E: 




- Molecule 2: Cbb3-type cytochrome c oxidase subunit II

Chain H: 




- Molecule 2: Cbb3-type cytochrome c oxidase subunit II

Chain L: 




- Molecule 3: Cbb3-type cytochrome c oxidase subunit CcoP1

Chain C: 




• Molecule 3: Cbb3-type cytochrome c oxidase subunit CcoP1

Chain F: 




• Molecule 3: Cbb3-type cytochrome c oxidase subunit CcoP1

Chain I: 




• Molecule 3: Cbb3-type cytochrome c oxidase subunit CcoP1




• Molecule 4: Putative uncharacterized protein




• Molecule 4: Putative uncharacterized protein



• Molecule 4: Putative uncharacterized protein



• Molecule 4: Putative uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	136.47Å 279.93Å 175.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.98 – 3.20 14.98 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.5 (14.98-3.20) 97.4 (14.98-3.20)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.187 , 0.223 0.190 , 0.224	Depositor DCC
R_{free} test set	6635 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	86.4	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 78.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	31974	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FC6, CA, HEC, CU, PEO, HEM, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/3811	0.97	14/5210 (0.3%)
1	D	0.48	0/3791	1.03	16/5182 (0.3%)
1	G	0.50	0/3804	0.93	13/5200 (0.2%)
1	K	0.48	1/3804 (0.0%)	0.97	11/5200 (0.2%)
2	B	0.44	0/1584	0.91	4/2146 (0.2%)
2	E	0.37	0/1584	0.85	2/2146 (0.1%)
2	H	0.43	0/1584	0.88	3/2146 (0.1%)
2	L	0.38	0/1584	0.85	2/2146 (0.1%)
3	C	0.65	0/2374	1.31	25/3225 (0.8%)
3	F	0.70	1/2374 (0.0%)	1.48	52/3225 (1.6%)
3	I	0.65	0/2374	1.30	26/3225 (0.8%)
3	M	0.69	0/2374	1.30	24/3225 (0.7%)
4	N	0.57	0/227	1.06	0/309
4	O	0.53	0/227	1.12	1/309 (0.3%)
4	P	0.57	0/227	1.10	0/309
4	Q	0.67	0/227	1.25	3/309 (1.0%)
All	All	0.54	2/31950 (0.0%)	1.09	196/43512 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	G	0	1
3	C	0	1
3	F	0	2
3	I	0	2
All	All	0	8

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	K	463	LYS	CA-C	6.43	1.60	1.52
3	F	36	THR	CB-OG1	5.05	1.51	1.43

The worst 5 of 196 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	462	ALA	N-CA-C	15.63	135.18	108.76
3	C	29	LYS	N-CA-C	13.58	126.08	111.28
1	A	467	TYR	N-CA-C	-13.04	87.81	109.46
1	A	468	ASP	N-CA-C	-11.19	86.96	110.80
3	F	102	GLU	N-CA-C	-10.54	99.79	111.28

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	465	ALA	Mainchain
1	A	467	TYR	Peptide
3	C	44	PHE	Sidechain
3	F	114	PHE	Sidechain
3	F	37	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3683	0	3663	142	0
1	D	3663	0	3644	172	0
1	G	3676	0	3656	156	0
1	K	3676	0	3656	168	0
2	B	1548	0	1526	50	0
2	E	1548	0	1526	50	0
2	H	1548	0	1526	49	0
2	L	1548	0	1526	44	0
3	C	2312	0	2237	103	0
3	F	2312	0	2237	199	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	I	2312	0	2237	130	0
3	M	2312	0	2237	144	0
4	N	221	0	226	9	0
4	O	221	0	226	11	0
4	P	221	0	226	13	0
4	Q	221	0	226	13	0
5	A	86	0	60	12	0
5	D	86	0	60	13	0
5	G	86	0	60	13	0
5	K	86	0	60	14	0
6	A	1	0	0	0	0
6	D	1	0	0	0	0
6	G	1	0	0	0	0
6	K	1	0	0	0	0
7	A	2	0	0	0	0
7	D	2	0	0	0	0
7	G	2	0	0	0	0
7	K	2	0	0	0	0
8	A	2	0	0	0	0
8	D	2	0	0	0	0
8	G	2	0	0	0	0
8	K	2	0	0	0	0
9	A	5	0	0	1	0
9	D	5	0	0	1	0
9	G	5	0	0	1	0
9	K	5	0	0	4	0
10	B	43	0	30	5	0
10	C	86	0	60	10	0
10	E	43	0	30	6	0
10	F	86	0	60	8	0
10	H	43	0	30	6	0
10	I	86	0	60	11	0
10	L	43	0	30	3	0
10	M	86	0	60	8	0
11	C	13	0	0	2	0
11	F	13	0	0	2	0
11	I	13	0	0	1	0
11	M	13	0	0	2	0
All	All	31974	0	31175	1293	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1293 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:207:HIS:NE2	1:D:251:TYR:CE1	1.70	1.59
1:A:207:HIS:NE2	1:A:251:TYR:CE1	1.70	1.56
1:G:207:HIS:NE2	1:G:251:TYR:CE1	1.70	1.56
1:K:207:HIS:NE2	1:K:251:TYR:CE1	1.70	1.53
1:K:207:HIS:NE2	1:K:251:TYR:HE1	1.02	1.33

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/474 (98%)	442 (95%)	20 (4%)	2 (0%)	30	62
1	D	461/474 (97%)	443 (96%)	17 (4%)	1 (0%)	43	73
1	G	463/474 (98%)	443 (96%)	19 (4%)	1 (0%)	43	73
1	K	463/474 (98%)	442 (96%)	19 (4%)	2 (0%)	30	62
2	B	195/203 (96%)	187 (96%)	8 (4%)	0	100	100
2	E	195/203 (96%)	187 (96%)	8 (4%)	0	100	100
2	H	195/203 (96%)	187 (96%)	8 (4%)	0	100	100
2	L	195/203 (96%)	187 (96%)	8 (4%)	0	100	100
3	C	301/311 (97%)	286 (95%)	15 (5%)	0	100	100
3	F	301/311 (97%)	286 (95%)	15 (5%)	0	100	100
3	I	301/311 (97%)	287 (95%)	14 (5%)	0	100	100
3	M	301/311 (97%)	284 (94%)	14 (5%)	3 (1%)	12	45
4	N	27/36 (75%)	27 (100%)	0	0	100	100
4	O	27/36 (75%)	27 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	P	27/36 (75%)	27 (100%)	0	0	100	100
4	Q	27/36 (75%)	27 (100%)	0	0	100	100
All	All	3943/4096 (96%)	3769 (96%)	165 (4%)	9 (0%)	43	73

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	466	GLU
1	A	468	ASP
1	G	465	ALA
1	K	466	GLU
1	K	468	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/379 (99%)	340 (91%)	34 (9%)	9	34
1	D	371/379 (98%)	333 (90%)	38 (10%)	7	29
1	G	373/379 (98%)	338 (91%)	35 (9%)	8	33
1	K	373/379 (98%)	337 (90%)	36 (10%)	8	32
2	B	166/172 (96%)	147 (89%)	19 (11%)	5	24
2	E	166/172 (96%)	148 (89%)	18 (11%)	6	27
2	H	166/172 (96%)	147 (89%)	19 (11%)	5	24
2	L	166/172 (96%)	149 (90%)	17 (10%)	7	29
3	C	227/234 (97%)	208 (92%)	19 (8%)	10	38
3	F	227/234 (97%)	195 (86%)	32 (14%)	3	17
3	I	227/234 (97%)	199 (88%)	28 (12%)	4	22
3	M	227/234 (97%)	197 (87%)	30 (13%)	4	19
4	N	22/28 (79%)	22 (100%)	0	100	100
4	O	22/28 (79%)	21 (96%)	1 (4%)	24	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	P	22/28 (79%)	21 (96%)	1 (4%)	24 58
4	Q	22/28 (79%)	19 (86%)	3 (14%)	3 18
All	All	3151/3252 (97%)	2821 (90%)	330 (10%)	6 28

5 of 330 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	I	123	GLU
2	L	62	ARG
3	I	193	ILE
1	K	192	LEU
3	M	35	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 118 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	282	GLN
3	M	38	GLN
1	G	431	HIS
2	L	193	GLN
1	K	414	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 44 ligands modelled in this entry, 12 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	HEC	I	402	3	46,50,50	2.53	24 (52%)	58,82,82	2.23	24 (41%)
10	HEC	B	301	2	46,50,50	2.45	20 (43%)	58,82,82	2.55	21 (36%)
10	HEC	F	402	3	46,50,50	2.66	27 (58%)	58,82,82	2.26	25 (43%)
8	PEO	G	505	6,5	1,1,1	0.36	0	-		
11	FC6	F	403	-	12,12,12	2.76	6 (50%)	-		
10	HEC	E	301	2	46,50,50	2.54	23 (50%)	58,82,82	2.39	23 (39%)
5	HEM	K	501	7,1,8	50,50,50	1.87	9 (18%)	67,82,82	1.27	2 (2%)
11	FC6	C	403	-	12,12,12	2.69	6 (50%)	-		
10	HEC	L	301	2	46,50,50	2.53	23 (50%)	58,82,82	2.39	21 (36%)
10	HEC	C	401	3	46,50,50	2.54	21 (45%)	58,82,82	2.36	22 (37%)
5	HEM	D	502	7,1	50,50,50	1.86	9 (18%)	67,82,82	1.36	5 (7%)
5	HEM	K	502	7,1	50,50,50	1.86	8 (16%)	67,82,82	1.37	5 (7%)
11	FC6	I	401	-	12,12,12	2.79	6 (50%)	-		
5	HEM	G	502	7,1	50,50,50	1.87	9 (18%)	67,82,82	1.41	7 (10%)
9	PO4	G	506	-	4,4,4	1.02	0	6,6,6	0.52	0
11	FC6	M	401	-	12,12,12	2.71	6 (50%)	-		
10	HEC	I	403	3	46,50,50	2.52	24 (52%)	58,82,82	2.27	28 (48%)
10	HEC	M	402	3	46,50,50	2.49	23 (50%)	58,82,82	2.36	23 (39%)
9	PO4	K	506	-	4,4,4	1.19	0	6,6,6	0.92	0
10	HEC	C	402	3	46,50,50	2.52	24 (52%)	58,82,82	2.39	24 (41%)
8	PEO	D	506	6,5	1,1,1	0.33	0	-		
5	HEM	G	501	7,8,1	50,50,50	1.84	9 (18%)	67,82,82	1.36	6 (8%)
8	PEO	K	507	6,5	1,1,1	0.33	0	-		
10	HEC	F	401	3	46,50,50	2.62	22 (47%)	58,82,82	2.26	24 (41%)
5	HEM	A	502	7,1	50,50,50	1.79	11 (22%)	67,82,82	1.36	8 (11%)
5	HEM	D	501	7,8,1	50,50,50	1.92	9 (18%)	67,82,82	1.29	3 (4%)
9	PO4	A	506	-	4,4,4	1.08	0	6,6,6	0.37	0
10	HEC	M	403	3	46,50,50	2.53	23 (50%)	58,82,82	2.30	26 (44%)
8	PEO	A	505	6,5	1,1,1	0.30	0	-		
5	HEM	A	501	7,8,1	50,50,50	1.84	10 (20%)	67,82,82	1.30	5 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	HEC	H	301	2	46,50,50	2.49	22 (47%)	58,82,82	2.44	22 (37%)
9	PO4	D	505	-	4,4,4	0.99	0	6,6,6	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	HEC	I	402	3	-	5/14/54/54	-
10	HEC	B	301	2	-	8/14/54/54	-
10	HEC	F	402	3	-	7/14/54/54	-
10	HEC	E	301	2	-	8/14/54/54	-
5	HEM	K	501	7,1,8	-	3/14/54/54	-
10	HEC	L	301	2	-	7/14/54/54	-
10	HEC	C	401	3	-	6/14/54/54	-
5	HEM	D	502	7,1	-	4/14/54/54	-
5	HEM	K	502	7,1	-	2/14/54/54	-
5	HEM	G	502	7,1	-	2/14/54/54	-
10	HEC	I	403	3	-	6/14/54/54	-
10	HEC	M	402	3	-	6/14/54/54	-
10	HEC	C	402	3	-	6/14/54/54	-
10	HEC	F	401	3	-	6/14/54/54	-
5	HEM	G	501	7,8,1	-	3/14/54/54	-
5	HEM	A	502	7,1	-	4/14/54/54	-
5	HEM	D	501	7,8,1	-	3/14/54/54	-
10	HEC	M	403	3	-	6/14/54/54	-
5	HEM	A	501	7,8,1	-	5/14/54/54	-
10	HEC	H	301	2	-	8/14/54/54	-

The worst 5 of 374 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	501	HEM	C3D-C2D	8.08	1.54	1.36
5	K	501	HEM	C3D-C2D	7.97	1.54	1.36
5	G	502	HEM	C3D-C2D	7.93	1.53	1.36
5	A	501	HEM	C3D-C2D	7.74	1.53	1.36
5	D	502	HEM	C3D-C2D	7.74	1.53	1.36

The worst 5 of 324 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	301	HEC	CBB-CAB-C3B	-8.40	110.64	127.43
10	E	301	HEC	CBB-CAB-C3B	-8.12	111.20	127.43
10	L	301	HEC	CBB-CAB-C3B	-8.01	111.42	127.43
10	H	301	HEC	CBB-CAB-C3B	-7.92	111.59	127.43
10	M	402	HEC	CBB-CAB-C3B	-6.96	113.51	127.43

There are no chirality outliers.

5 of 105 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	301	HEC	C2B-C3B-CAB-CBB
10	B	301	HEC	C4B-C3B-CAB-CBB
10	B	301	HEC	C4C-C3C-CAC-CBC
10	C	401	HEC	C2B-C3B-CAB-CBB
10	C	401	HEC	C4B-C3B-CAB-CBB

There are no ring outliers.

28 monomers are involved in 123 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	I	402	HEC	8	0
10	B	301	HEC	5	0
10	F	402	HEC	3	0
11	F	403	FC6	2	0
10	E	301	HEC	6	0
5	K	501	HEM	5	0
11	C	403	FC6	2	0
10	L	301	HEC	3	0
10	C	401	HEC	8	0
5	D	502	HEM	8	0
5	K	502	HEM	9	0
11	I	401	FC6	1	0
5	G	502	HEM	8	0
9	G	506	PO4	1	0
11	M	401	FC6	2	0
10	I	403	HEC	3	0
10	M	402	HEC	6	0
9	K	506	PO4	4	0
10	C	402	HEC	2	0
5	G	501	HEM	5	0
10	F	401	HEC	5	0


Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	502	HEM	9	0
5	D	501	HEM	5	0
9	A	506	PO4	1	0
10	M	403	HEC	2	0
5	A	501	HEM	3	0
10	H	301	HEC	6	0
9	D	505	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEC I 402



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC B 301



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC F 402




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC E 301



Bond lengths




Bond angles




Torsions




Rings




Ligand HEC L 301



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC C 401



Bond lengths




Bond angles





Torsions




Rings









Ligand HEC I 403



Bond lengths




Bond angles




Torsions




Rings




Ligand HEC C 402




Bond lengths




Bond angles




Torsions




Rings




Ligand HEC F 401



Bond lengths




Bond angles





Torsions





Rings











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/474 (98%)	-0.72	4 (0%) 81 64	45, 76, 129, 206	0
1	D	463/474 (97%)	-0.30	1 (0%) 91 85	76, 153, 234, 310	0
1	G	465/474 (98%)	-0.62	5 (1%) 78 61	55, 90, 142, 254	0
1	K	465/474 (98%)	-0.23	4 (0%) 81 64	61, 145, 226, 280	0
2	B	197/203 (97%)	-0.83	0 100 100	42, 70, 126, 169	0
2	E	197/203 (97%)	-0.34	0 100 100	68, 138, 213, 279	0
2	H	197/203 (97%)	-0.71	0 100 100	54, 83, 146, 202	0
2	L	197/203 (97%)	-0.43	2 (1%) 79 63	52, 110, 215, 322	0
3	C	303/311 (97%)	-0.64	0 100 100	41, 83, 138, 192	0
3	F	303/311 (97%)	-0.01	6 (1%) 65 45	72, 173, 260, 419	0
3	I	303/311 (97%)	-0.54	1 (0%) 90 81	52, 106, 166, 215	0
3	M	303/311 (97%)	-0.43	2 (0%) 84 69	43, 85, 243, 322	0
4	N	29/36 (80%)	-0.19	0 100 100	65, 92, 167, 174	0
4	O	29/36 (80%)	-0.06	0 100 100	148, 172, 249, 263	0
4	P	29/36 (80%)	-0.19	0 100 100	91, 109, 173, 191	0
4	Q	29/36 (80%)	0.36	0 100 100	119, 142, 186, 199	0
All	All	3975/4096 (97%)	-0.46	25 (0%) 85 73	41, 107, 219, 419	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	43	ALA	3.4
3	F	2	SER	3.1
1	A	465	ALA	3.1
3	M	50	TYR	3.0
3	F	49	GLU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	PO4	D	505	5/5	0.59	0.11	180,183,183,185	0
9	PO4	K	506	5/5	0.80	0.09	138,141,143,150	0
9	PO4	G	506	5/5	0.82	0.10	130,132,139,152	0
11	FC6	F	403	13/13	0.85	0.33	53,61,87,101	13
7	CA	D	504	1/1	0.89	0.10	151,151,151,151	0
9	PO4	A	506	5/5	0.91	0.07	60,68,75,88	0
8	PEO	K	507	2/2	0.92	0.09	131,131,131,131	0
7	CA	K	505	1/1	0.93	0.08	136,136,136,136	0
11	FC6	I	401	13/13	0.93	0.18	78,94,143,158	13
10	HEC	C	402	43/43	0.94	0.12	36,54,72,160	0
7	CA	A	504	1/1	0.95	0.06	99,99,99,99	0
7	CA	A	507	1/1	0.95	0.18	84,84,84,84	0
6	CU	A	503	1/1	0.95	0.20	133,133,133,133	0
6	CU	K	503	1/1	0.96	0.05	133,133,133,133	0
7	CA	G	504	1/1	0.96	0.06	87,87,87,87	0
10	HEC	B	301	43/43	0.96	0.10	39,55,69,119	0
8	PEO	D	506	2/2	0.97	0.06	145,145,145,147	0
7	CA	D	507	1/1	0.97	0.05	96,96,96,96	0
5	HEM	K	501	43/43	0.97	0.13	119,135,151,154	0
10	HEC	F	402	43/43	0.97	0.12	117,134,147,154	0
5	HEM	D	501	43/43	0.97	0.10	93,119,136,143	0
8	PEO	A	505	2/2	0.97	0.07	74,74,74,74	0
5	HEM	G	502	43/43	0.98	0.07	54,66,77,86	0
6	CU	D	503	1/1	0.98	0.03	118,118,118,118	0
5	HEM	D	502	43/43	0.98	0.08	89,110,124,130	0
10	HEC	C	401	43/43	0.98	0.08	44,57,66,73	0
8	PEO	G	505	2/2	0.98	0.05	80,80,80,83	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
10	HEC	F	401	43/43	0.98	0.09	85,103,116,121	0
5	HEM	K	502	43/43	0.98	0.10	95,108,128,130	0
10	HEC	H	301	43/43	0.98	0.08	42,53,70,79	0
10	HEC	I	402	43/43	0.98	0.08	38,54,77,91	0
10	HEC	I	403	43/43	0.98	0.09	50,61,69,85	0
10	HEC	M	402	43/43	0.98	0.07	33,42,54,63	0
10	HEC	M	403	43/43	0.98	0.08	35,51,64,84	0
11	FC6	C	403	13/13	0.98	0.10	60,75,98,101	13
7	CA	G	507	1/1	0.98	0.04	72,72,72,72	0
7	CA	K	504	1/1	0.98	0.03	95,95,95,95	0
11	FC6	M	401	13/13	0.98	0.11	43,60,98,98	13
5	HEM	A	501	43/43	0.99	0.07	45,63,72,76	0
10	HEC	E	301	43/43	0.99	0.07	62,73,96,102	0
5	HEM	A	502	43/43	0.99	0.08	41,54,79,83	0
10	HEC	L	301	43/43	0.99	0.08	49,65,78,78	0
5	HEM	G	501	43/43	0.99	0.06	57,72,84,93	0
6	CU	G	503	1/1	1.00	0.04	74,74,74,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEC C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM K 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC F 402:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM G 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC F 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM K 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC H 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC I 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC I 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC M 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC M 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around HEM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HEM A 502:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.