



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:12 PM UTC

PDB ID : 8DJ6 / pdb_00008dj6
Title : Sliding-clamp-ImuB peptide
Authors : Kapur, M.K.; Gray, O.J.; Honzatko, R.H.; Nelson, S.N.
Deposited on : 2022-06-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	8% 94%
1	B	397	9% 95%
1	C	397	6% 95%
1	D	397	11% 94%
2	E	8	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	8	 100%
2	G	8	 100%
2	H	8	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24034 atoms, of which 11879 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	389	Total	C	H	N	O	S	0	1	0
			5806	1821	2919	494	565	7			
1	C	388	Total	C	H	N	O	S	0	2	0
			5798	1820	2912	495	564	7			
1	B	389	Total	C	H	N	O	S	0	0	0
			5787	1816	2908	491	565	7			
1	D	388	Total	C	H	N	O	S	0	3	0
			5806	1823	2915	495	566	7			

- Molecule 2 is a protein called Imub-peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	8	Total	C	H	N	O	0	0	1
			106	37	52	9	8			
2	F	8	Total	C	H	N	O	0	0	1
			106	37	52	9	8			
2	G	8	Total	C	H	N	O	0	0	1
			106	37	52	9	8			
2	H	8	Total	C	H	N	O	0	0	1
			106	37	52	9	8			

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			5	1	2	2		
3	A	1	Total	C	H	O	0	0
			5	1	2	2		
3	C	1	Total	C	H	O	0	0
			5	1	2	2		
3	B	1	Total	C	H	O	0	0
			5	1	2	2		
3	B	1	Total	C	H	O	0	0
			5	1	2	2		
3	B	1	Total	C	H	O	0	0
			5	1	2	2		
3	D	1	Total	C	H	O	0	0
			5	1	2	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	95	Total	O	0	0
			95	95		
5	C	82	Total	O	0	0
			82	82		
5	B	90	Total	O	0	0
			90	90		
5	D	80	Total	O	0	0
			80	80		
5	E	9	Total	O	0	0
			9	9		
5	F	7	Total	O	0	0
			7	7		
5	G	4	Total	O	0	0
			4	4		
5	H	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta sliding clamp



- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp



- Molecule 1: Beta sliding clamp





- Molecule 2: Imub-peptide

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Imub-peptide

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Imub-peptide

Chain G:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Imub-peptide

Chain H:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	214.11Å 124.83Å 117.26Å 90.00° 113.75° 90.00°	Depositor
Resolution (Å)	46.75 – 2.50 46.75 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.3 (46.75-2.50) 96.6 (46.75-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.200 , 0.226 0.203 , 0.224	Depositor DCC
R_{free} test set	2000 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	54.9	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	24034	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, FMT, NH2, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.09	0/2942	0.25	0/4018
1	B	0.09	0/2931	0.26	0/4004
1	C	0.09	0/2945	0.26	0/4022
1	D	0.09	0/2953	0.25	0/4033
2	E	0.26	0/53	0.26	0/73
2	F	0.25	0/53	0.24	0/73
2	G	0.25	0/53	0.24	0/73
2	H	0.22	0/53	0.24	0/73
All	All	0.10	0/11983	0.25	0/16369

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2887	2919	2920	7	0
1	B	2879	2908	2907	7	0
1	C	2886	2912	2916	6	0
1	D	2891	2915	2920	9	0
2	E	54	52	53	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	54	52	53	0	0
2	G	54	52	53	0	0
2	H	54	52	53	0	0
3	A	6	4	2	0	0
3	B	9	6	3	0	0
3	C	3	2	1	0	0
3	D	3	2	1	0	0
4	B	4	3	3	0	0
5	A	95	0	0	0	0
5	B	90	0	0	1	0
5	C	82	0	0	0	0
5	D	80	0	0	2	0
5	E	9	0	0	0	0
5	F	7	0	0	0	0
5	G	4	0	0	0	0
5	H	4	0	0	0	0
All	All	12155	11879	11885	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:MET:HE3	1:B:189:LEU:HD11	1.65	0.78
1:C:295:VAL:HG23	1:C:335:PRO:HG3	1.77	0.67
1:A:295:VAL:HG23	1:A:335:PRO:HG3	1.77	0.65
1:B:155:ARG:O	1:B:158:THR:HG22	1.98	0.63
1:C:55:SER:HB3	1:C:127:LEU:HD11	1.87	0.56
1:A:29:ARG:NE	1:A:215:THR:HG21	2.21	0.55
1:A:37:VAL:HG12	1:A:39:VAL:HG22	1.94	0.50
1:C:78:LEU:HB2	1:C:121:VAL:HG12	1.94	0.49
1:D:78:LEU:HB2	1:D:121:VAL:HG12	1.94	0.48
1:B:237:LEU:N	1:B:237:LEU:HD23	2.29	0.48
1:D:254:ALA:N	5:D:506:HOH:O	2.48	0.47
1:A:272:THR:HG22	1:A:349:ARG:HD3	1.97	0.46
1:B:183:ARG:NH1	1:B:340:ASP:OD2	2.48	0.45
1:B:310:ALA:HB1	5:B:533:HOH:O	2.17	0.43
1:D:360:PRO:HB2	1:D:389:LEU:HD11	2.00	0.43
1:A:158:THR:OG1	1:A:159:LEU:N	2.51	0.43
1:C:174:MET:HE3	1:C:189:LEU:HD11	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:55:SER:HB3	1:D:127:LEU:HD11	2.00	0.43
1:D:332:ALA:HB2	1:D:395:LEU:HD21	2.00	0.42
1:D:189:LEU:N	1:D:189:LEU:HD23	2.35	0.42
1:A:174:MET:HE3	1:A:189:LEU:HD21	2.01	0.42
1:B:58:ASP:O	1:B:59:TYR:HB2	2.20	0.41
1:C:229:GLY:O	1:C:232:VAL:HG23	2.20	0.41
1:D:58:ASP:O	1:D:59:TYR:HB2	2.19	0.41
1:D:237:LEU:HD23	1:D:237:LEU:N	2.35	0.41
1:B:37:VAL:HG12	1:B:39:VAL:HG22	2.03	0.41
1:D:121:VAL:HG13	5:D:516:HOH:O	2.19	0.40
1:A:58:ASP:O	1:A:59:TYR:HB2	2.20	0.40
1:C:142:PHE:CE1	1:C:174:MET:HE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/397 (98%)	374 (96%)	14 (4%)	0	100	100
1	B	387/397 (98%)	374 (97%)	13 (3%)	0	100	100
1	C	388/397 (98%)	369 (95%)	19 (5%)	0	100	100
1	D	389/397 (98%)	372 (96%)	17 (4%)	0	100	100
2	E	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
2	F	6/8 (75%)	6 (100%)	0	0	100	100
2	G	6/8 (75%)	6 (100%)	0	0	100	100
2	H	6/8 (75%)	6 (100%)	0	0	100	100
All	All	1576/1620 (97%)	1512 (96%)	64 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/311 (99%)	307 (100%)	0	100	100
1	B	306/311 (98%)	306 (100%)	0	100	100
1	C	307/311 (99%)	306 (100%)	1 (0%)	86	94
1	D	308/311 (99%)	306 (99%)	2 (1%)	78	91
2	E	5/5 (100%)	5 (100%)	0	100	100
2	F	5/5 (100%)	5 (100%)	0	100	100
2	G	5/5 (100%)	5 (100%)	0	100	100
2	H	5/5 (100%)	5 (100%)	0	100	100
All	All	1248/1264 (99%)	1245 (100%)	3 (0%)	87	95

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	122	GLU
1	D	93	ASN
1	D	290	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	B	294	GLN
1	D	267	HIS
2	E	2	GLN
2	G	2	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FMT	C	401	-	2,2,2	0.75	0	1,1,1	0.46	0
3	FMT	D	401	-	2,2,2	0.73	0	1,1,1	0.46	0
3	FMT	A	402	-	2,2,2	0.74	0	1,1,1	0.45	0
3	FMT	B	403	-	2,2,2	0.73	0	1,1,1	0.44	0
3	FMT	A	401	-	2,2,2	0.72	0	1,1,1	0.45	0
3	FMT	B	402	-	2,2,2	0.73	0	1,1,1	0.45	0
4	ACT	B	404	-	3,3,3	1.39	1 (33%)	3,3,3	1.36	0
3	FMT	B	401	-	2,2,2	0.75	0	1,1,1	0.45	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	404	ACT	CH3-C	2.00	1.57	1.49

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/397 (97%)	0.45	30 (7%) 19 17	34, 63, 114, 173	1 (0%)
1	B	389/397 (97%)	0.49	35 (8%) 15 13	43, 63, 111, 175	0
1	C	388/397 (97%)	0.34	25 (6%) 25 22	40, 63, 111, 170	2 (0%)
1	D	388/397 (97%)	0.50	43 (11%) 10 8	35, 65, 121, 154	3 (0%)
2	E	6/8 (75%)	0.01	0 100 100	47, 47, 54, 77	0
2	F	6/8 (75%)	-0.06	0 100 100	43, 47, 53, 72	0
2	G	6/8 (75%)	-0.39	0 100 100	48, 55, 59, 74	0
2	H	6/8 (75%)	-0.07	0 100 100	52, 57, 60, 81	0
All	All	1578/1620 (97%)	0.44	133 (8%) 17 15	34, 63, 114, 175	6 (0%)

All (133) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	292	GLY	6.0
1	B	195	THR	4.8
1	C	292	GLY	4.8
1	A	290	ASP	4.7
1	C	310	ALA	4.7
1	D	310	ALA	4.5
1	A	196	PRO	4.5
1	B	293	ALA	4.4
1	B	310	ALA	4.3
1	A	292	GLY	4.3
1	D	396	PRO	4.2
1	B	196	PRO	4.1
1	D	82[A]	ARG	4.1
1	A	373	LEU	4.1
1	D	293	ALA	4.0
1	A	311	ASP	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	396	PRO	3.9
1	B	290	ASP	3.8
1	C	82[A]	ARG	3.8
1	A	310	ALA	3.8
1	D	369	ALA	3.8
1	C	195	THR	3.7
1	D	9	THR	3.6
1	D	292	GLY	3.6
1	C	196	PRO	3.6
1	A	396	PRO	3.5
1	C	396	PRO	3.5
1	A	89	ARG	3.5
1	C	197	ASP	3.4
1	B	375	GLY	3.4
1	A	291	ARG	3.4
1	D	373	LEU	3.4
1	C	229	GLY	3.3
1	B	229	GLY	3.3
1	D	290	ASP	3.2
1	C	373	LEU	3.2
1	C	305[A]	HIS	3.1
1	B	373	LEU	3.1
1	C	374	ASN	3.1
1	A	195	THR	3.0
1	D	294	GLN	3.0
1	B	231	SER	3.0
1	A	374	ASN	3.0
1	A	235	ASP	3.0
1	A	229	GLY	3.0
1	D	375	GLY	3.0
1	B	374	ASN	3.0
1	A	372	ALA	2.9
1	A	230	PRO	2.9
1	B	313	VAL	2.9
1	D	230	PRO	2.9
1	B	235	ASP	2.9
1	D	158	THR	2.9
1	D	156	ASP	2.9
1	B	376	ASN	2.9
1	C	293	ALA	2.9
1	C	33	SER	2.8
1	D	374	ASN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	371	ALA	2.8
1	A	157	ASP	2.8
1	B	230	PRO	2.8
1	A	8	LEU	2.8
1	B	34	ARG	2.8
1	D	196	PRO	2.7
1	C	159	LEU	2.7
1	C	9	THR	2.7
1	B	197	ASP	2.7
1	D	159	LEU	2.6
1	D	235	ASP	2.6
1	B	383	GLU	2.6
1	B	194	LYS	2.6
1	D	372	ALA	2.6
1	C	34	ARG	2.5
1	D	291	ARG	2.5
1	C	230	PRO	2.5
1	A	198	VAL	2.5
1	A	197	ASP	2.5
1	D	155	ARG	2.5
1	A	194	LYS	2.5
1	B	33	SER	2.5
1	C	253	ASP	2.4
1	A	34	ARG	2.4
1	A	315	ARG	2.4
1	D	34	ARG	2.4
1	D	300	ALA	2.4
1	D	311	ASP	2.4
1	D	312	ASP	2.4
1	B	8	LEU	2.4
1	A	158	THR	2.3
1	A	156	ASP	2.3
1	A	228	ALA	2.3
1	D	195	THR	2.3
1	D	237	LEU	2.3
1	A	289	ALA	2.3
1	A	371	ALA	2.3
1	B	158	THR	2.3
1	B	89	ARG	2.3
1	D	395	LEU	2.3
1	C	301	ASP	2.3
1	B	157	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	376	ASN	2.3
1	B	311	ASP	2.3
1	B	233	GLY	2.2
1	D	229	GLY	2.2
1	D	243	GLU	2.2
1	B	159	LEU	2.2
1	C	290	ASP	2.2
1	D	376	ASN	2.2
1	B	294	GLN	2.2
1	D	356	THR	2.2
1	D	329	LEU	2.2
1	A	375	GLY	2.2
1	C	156	ASP	2.2
1	C	255	GLU	2.2
1	D	50	ASP	2.2
1	C	372	ALA	2.1
1	B	371	ALA	2.1
1	D	305[A]	HIS	2.1
1	B	173	ARG	2.1
1	B	349	ARG	2.1
1	D	368	GLU	2.1
1	B	242	SER	2.1
1	D	301	ASP	2.1
1	C	194	LYS	2.1
1	B	372	ALA	2.1
1	C	375	GLY	2.1
1	B	156	ASP	2.1
1	D	197	ASP	2.1
1	D	48	SER	2.1
1	D	194	LYS	2.0
1	A	377	GLY	2.0
1	D	157	ASP	2.0
1	D	378	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	B	404	4/4	0.69	0.30	56,67,86,91	0
3	FMT	A	401	3/3	0.75	0.31	86,87,104,105	0
3	FMT	B	403	3/3	0.80	0.34	71,81,97,102	0
3	FMT	A	402	3/3	0.80	0.24	66,69,88,105	0
3	FMT	B	402	3/3	0.82	0.26	75,77,93,98	0
3	FMT	C	401	3/3	0.84	0.20	70,75,84,95	0
3	FMT	D	401	3/3	0.88	0.17	65,68,82,88	0
3	FMT	B	401	3/3	0.88	0.21	73,81,88,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.