



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:11 PM UTC

PDB ID : 3DKQ / pdb_00003dkq
Title : Crystal structure of Putative Oxygenase (YP_001051978.1) from SHE-
WANELLA BALTICA OS155 at 2.26 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2008-06-25
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	11% 88% 6% • 5%
1	B	243	30% 80% 5% 15%
1	C	243	18% 83% 7% • 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PKHD-type hydroxylase Sbal_3634.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	Se	0	7	0
			1867	1180	332	349	6			
1	B	207	Total	C	N	O	Se	0	2	0
			1640	1036	288	312	4			
1	C	221	Total	C	N	O	Se	0	4	0
			1778	1125	309	339	5			

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	expression tag	UNP A3D8P6
A	-17	GLY	-	expression tag	UNP A3D8P6
A	-16	SER	-	expression tag	UNP A3D8P6
A	-15	ASP	-	expression tag	UNP A3D8P6
A	-14	LYS	-	expression tag	UNP A3D8P6
A	-13	ILE	-	expression tag	UNP A3D8P6
A	-12	HIS	-	expression tag	UNP A3D8P6
A	-11	HIS	-	expression tag	UNP A3D8P6
A	-10	HIS	-	expression tag	UNP A3D8P6
A	-9	HIS	-	expression tag	UNP A3D8P6
A	-8	HIS	-	expression tag	UNP A3D8P6
A	-7	HIS	-	expression tag	UNP A3D8P6
A	-6	GLU	-	expression tag	UNP A3D8P6
A	-5	ASN	-	expression tag	UNP A3D8P6
A	-4	LEU	-	expression tag	UNP A3D8P6
A	-3	TYR	-	expression tag	UNP A3D8P6
A	-2	PHE	-	expression tag	UNP A3D8P6
A	-1	GLN	-	expression tag	UNP A3D8P6
A	0	GLY	-	expression tag	UNP A3D8P6
B	-18	MSE	-	expression tag	UNP A3D8P6
B	-17	GLY	-	expression tag	UNP A3D8P6
B	-16	SER	-	expression tag	UNP A3D8P6
B	-15	ASP	-	expression tag	UNP A3D8P6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	LYS	-	expression tag	UNP A3D8P6
B	-13	ILE	-	expression tag	UNP A3D8P6
B	-12	HIS	-	expression tag	UNP A3D8P6
B	-11	HIS	-	expression tag	UNP A3D8P6
B	-10	HIS	-	expression tag	UNP A3D8P6
B	-9	HIS	-	expression tag	UNP A3D8P6
B	-8	HIS	-	expression tag	UNP A3D8P6
B	-7	HIS	-	expression tag	UNP A3D8P6
B	-6	GLU	-	expression tag	UNP A3D8P6
B	-5	ASN	-	expression tag	UNP A3D8P6
B	-4	LEU	-	expression tag	UNP A3D8P6
B	-3	TYR	-	expression tag	UNP A3D8P6
B	-2	PHE	-	expression tag	UNP A3D8P6
B	-1	GLN	-	expression tag	UNP A3D8P6
B	0	GLY	-	expression tag	UNP A3D8P6
C	-18	MSE	-	expression tag	UNP A3D8P6
C	-17	GLY	-	expression tag	UNP A3D8P6
C	-16	SER	-	expression tag	UNP A3D8P6
C	-15	ASP	-	expression tag	UNP A3D8P6
C	-14	LYS	-	expression tag	UNP A3D8P6
C	-13	ILE	-	expression tag	UNP A3D8P6
C	-12	HIS	-	expression tag	UNP A3D8P6
C	-11	HIS	-	expression tag	UNP A3D8P6
C	-10	HIS	-	expression tag	UNP A3D8P6
C	-9	HIS	-	expression tag	UNP A3D8P6
C	-8	HIS	-	expression tag	UNP A3D8P6
C	-7	HIS	-	expression tag	UNP A3D8P6
C	-6	GLU	-	expression tag	UNP A3D8P6
C	-5	ASN	-	expression tag	UNP A3D8P6
C	-4	LEU	-	expression tag	UNP A3D8P6
C	-3	TYR	-	expression tag	UNP A3D8P6
C	-2	PHE	-	expression tag	UNP A3D8P6
C	-1	GLN	-	expression tag	UNP A3D8P6
C	0	GLY	-	expression tag	UNP A3D8P6

- Molecule 2 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0
2	B	1	Total Ni 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Ni	0	0
			1	1		

- Molecule 3 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	A	1	Total	C	N	0	0
			5	3	2		
3	B	1	Total	C	N	0	0
			5	3	2		
3	B	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	119	Total	O	0	0
			119	119		
5	B	52	Total	O	0	2
			54	54		
5	C	73	Total	O	0	0
			73	73		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PKHD-type hydroxylase Sbal_3634



• Molecule 1: PKHD-type hydroxylase Sbal_3634



• Molecule 1: PKHD-type hydroxylase Sbal_3634



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	153.45Å 153.45Å 84.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.86 – 2.26 28.86 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.86-2.26) 99.9 (28.86-2.26)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, R_{free}	0.196 , 0.232 0.204 , 0.238	Depositor DCC
R_{free} test set	2424 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5600	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NI, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	3/1919 (0.2%)	1.05	2/2586 (0.1%)
1	B	0.83	0/1670	1.04	5/2255 (0.2%)
1	C	0.83	1/1821 (0.1%)	1.02	2/2458 (0.1%)
All	All	0.85	4/5410 (0.1%)	1.04	9/7299 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	177	MSE	SE-CE	-8.14	1.71	1.95
1	A	36	MSE	SE-CE	-6.16	1.76	1.95
1	A	108	MSE	SE-CE	5.43	2.11	1.95
1	C	177	MSE	SE-CE	-5.39	1.79	1.95

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	80	TYR	N-CA-C	-9.91	97.34	109.72
1	A	80	TYR	N-CA-C	-8.16	97.90	109.24
1	A	137	GLY	N-CA-C	6.48	120.01	111.52
1	B	80	TYR	N-CA-C	-6.18	99.65	109.04
1	B	157	HIS	N-CA-C	5.67	117.47	108.79
1	C	157	HIS	N-CA-C	5.43	117.05	108.96
1	B	80	TYR	CA-C-N	5.07	125.61	120.38
1	B	80	TYR	C-N-CA	5.07	125.61	120.38
1	B	97	ILE	N-CA-C	-5.06	101.19	108.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1867	0	1849	11	0
1	B	1640	0	1590	5	0
1	C	1778	0	1740	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	10	0	10	0	0
3	B	10	0	10	0	0
3	C	10	0	10	0	0
4	A	30	0	40	0	0
4	C	6	0	8	0	0
5	A	119	0	0	1	0
5	B	54	0	0	0	0
5	C	73	0	0	0	0
All	All	5600	0	5257	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:MSE:HE2	1:A:97:ILE:HB	1.44	0.98
1:C:-4:LEU:HD21	1:C:3:ILE:HD11	1.55	0.86
1:A:36:MSE:HE2	1:A:97:ILE:CB	2.11	0.80
1:A:75:LEU:HB3	1:A:177:MSE:HE3	1.68	0.75
1:A:32:THR:HG21	1:A:35:ALA:HB3	1.77	0.66
1:A:36:MSE:HE2	1:A:97:ILE:CG2	2.34	0.57
1:A:171[B]:PHE:HE2	1:A:173:TRP:CZ2	2.24	0.55
1:C:-4:LEU:HD12	1:C:69:GLN:CD	2.32	0.55
1:A:186:LEU:HD21	1:B:213:VAL:HG13	1.89	0.54
1:C:2:LEU:HD11	1:C:139:GLN:HE22	1.74	0.53
1:A:32:THR:CG2	1:A:35:ALA:HB3	2.38	0.52
1:C:14:VAL:HG11	1:C:145:ALA:HB1	1.91	0.52
1:A:100:ALA:C	1:A:101:ILE:HD12	2.36	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:177:MSE:CE	1:C:218:LEU:HD11	2.41	0.50
1:C:-4:LEU:HD21	1:C:3:ILE:CD1	2.37	0.49
1:B:75:LEU:HB3	1:B:177:MSE:HE3	1.94	0.48
1:B:186:LEU:HD23	1:B:186:LEU:C	2.40	0.46
1:B:14:VAL:HG11	1:B:145:ALA:HB1	1.97	0.45
1:C:-4:LEU:HD22	1:C:-2:PHE:CE1	2.51	0.45
1:C:3:ILE:HG23	1:C:150:LEU:HB3	1.98	0.45
1:C:114:SER:HG	1:C:173:TRP:CD1	2.36	0.43
1:C:132:ILE:HG23	1:C:155:SER:HB3	2.02	0.42
1:A:174:LEU:HD12	1:A:174:LEU:C	2.45	0.42
1:B:203:ALA:HB3	1:B:206:GLU:HG2	2.03	0.41
1:C:177:MSE:HE2	1:C:218:LEU:HD11	2.03	0.41
1:A:177:MSE:HE2	5:A:588:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/243 (97%)	234 (100%)	1 (0%)	0	100	100
1	B	203/243 (84%)	200 (98%)	3 (2%)	0	100	100
1	C	221/243 (91%)	216 (98%)	5 (2%)	0	100	100
All	All	659/729 (90%)	650 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/204 (99%)	200 (99%)	2 (1%)	68	77
1	B	174/204 (85%)	172 (99%)	2 (1%)	65	75
1	C	193/204 (95%)	189 (98%)	4 (2%)	47	58
All	All	569/612 (93%)	561 (99%)	8 (1%)	61	70

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	148	LEU
1	B	83	LEU
1	B	179	ARG
1	C	-4	LEU
1	C	19[A]	GLU
1	C	19[B]	GLU
1	C	123	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	HIS
1	A	99	ASN
1	A	139	GLN
1	A	189	GLN
1	A	192	GLN
1	B	69	GLN
1	B	88	GLN
1	B	124	ASN
1	C	85	ASN
1	C	124	ASN
1	C	139	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 3 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	505	-	5,5,5	0.39	0	5,5,5	0.68	0
4	GOL	A	507	-	5,5,5	0.44	0	5,5,5	0.65	0
3	IMD	C	501	-	5,5,5	0.38	0	5,5,5	0.60	0
4	GOL	A	506	-	5,5,5	0.50	0	5,5,5	0.76	0
3	IMD	C	502	-	5,5,5	0.38	0	5,5,5	0.60	0
4	GOL	C	503	-	5,5,5	0.48	0	5,5,5	0.25	0
3	IMD	B	501	-	5,5,5	0.60	0	5,5,5	0.53	0
4	GOL	A	504	-	5,5,5	0.35	0	5,5,5	0.28	0
4	GOL	A	503	-	5,5,5	0.49	0	5,5,5	0.83	0
3	IMD	A	501	2	5,5,5	0.35	0	5,5,5	0.67	0
3	IMD	B	502	-	5,5,5	0.57	0	5,5,5	0.62	0
3	IMD	A	502	-	5,5,5	0.38	0	5,5,5	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	507	-	-	2/4/4/4	-
3	IMD	C	501	-	-	-	0/1/1/1
4	GOL	A	506	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	503	-	-	0/4/4/4	-
3	IMD	C	502	-	-	-	0/1/1/1
3	IMD	B	502	-	-	-	0/1/1/1
3	IMD	B	501	-	-	-	0/1/1/1
4	GOL	A	504	-	-	0/4/4/4	-
4	GOL	A	503	-	-	2/4/4/4	-
3	IMD	A	501	2	-	-	0/1/1/1
4	GOL	A	505	-	-	2/4/4/4	-
3	IMD	A	502	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	GOL	C1-C2-C3-O3
4	A	503	GOL	O2-C2-C3-O3
4	A	506	GOL	O1-C1-C2-C3
4	A	507	GOL	O1-C1-C2-C3
4	A	505	GOL	O2-C2-C3-O3
4	A	505	GOL	C1-C2-C3-O3
4	A	506	GOL	O1-C1-C2-O2
4	A	507	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/243 (92%)	0.73	26 (11%) 9 9	26, 52, 83, 99	7 (3%)
1	B	203/243 (83%)	1.76	73 (35%) 1 0	26, 52, 74, 109	2 (0%)
1	C	216/243 (88%)	1.45	44 (20%) 3 2	30, 52, 78, 104	4 (1%)
All	All	643/729 (88%)	1.30	143 (22%) 2 2	26, 52, 79, 109	13 (2%)

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	-2	PHE	7.3
1	B	-3	TYR	6.9
1	B	-4	LEU	6.8
1	A	-5	ASN	5.9
1	B	29	GLY	5.8
1	B	-2	PHE	5.7
1	B	-1	GLN	5.5
1	C	135	THR	5.4
1	B	203	ALA	5.4
1	B	199	ALA	5.3
1	B	171	PHE	5.2
1	A	-3	TYR	5.1
1	B	136	TYR	5.0
1	A	-2	PHE	5.0
1	C	-4	LEU	4.9
1	B	-5	ASN	4.7
1	C	136	TYR	4.6
1	C	134	ASP	4.6
1	C	-3	TYR	4.5
1	B	92	THR	4.5
1	B	95	TYR	4.4
1	A	-4	LEU	4.4
1	A	136	TYR	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	135	THR	4.3
1	B	137	GLY	4.2
1	C	31	GLN	4.1
1	A	32	THR	4.1
1	C	162	VAL	4.1
1	A	-1	GLN	4.0
1	B	97	ILE	4.0
1	C	40	ARG	3.9
1	B	202	ALA	3.8
1	A	33	SER	3.7
1	A	35	ALA	3.7
1	B	110	ARG	3.6
1	B	28	ASP	3.6
1	A	31	GLN	3.5
1	B	83	LEU	3.5
1	B	3	ILE	3.5
1	A	37	ALA	3.4
1	B	27	ILE	3.4
1	C	30	ASN	3.4
1	B	42	ARG	3.4
1	B	90	GLY	3.4
1	B	44	GLN	3.4
1	A	30	ASN	3.4
1	B	43	ASN	3.3
1	A	0	GLY	3.3
1	A	38	THR	3.3
1	C	41	LYS	3.2
1	C	139	GLN	3.2
1	B	26	TRP	3.2
1	C	127	GLY	3.1
1	A	135	THR	3.1
1	C	-1	GLN	3.1
1	C	3	ILE	3.0
1	C	137	GLY	3.0
1	C	2	LEU	2.9
1	B	197	LEU	2.9
1	B	196	SER	2.9
1	A	29	GLY	2.9
1	B	23	ALA	2.9
1	B	45	GLN	2.9
1	B	215	HIS	2.8
1	B	69	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	207	LEU	2.8
1	B	0	GLY	2.8
1	B	96	HIS	2.8
1	B	52	VAL	2.8
1	B	201	THR	2.8
1	C	165	GLY	2.8
1	A	34	GLY	2.7
1	C	29	GLY	2.7
1	B	208	PHE	2.7
1	C	208	PHE	2.7
1	B	94	GLY	2.7
1	C	171	PHE	2.7
1	B	133	GLN	2.7
1	B	98	ASP	2.7
1	B	209	ASN	2.7
1	B	156	LEU	2.7
1	B	80	TYR	2.7
1	C	138	GLN	2.7
1	B	205	GLN	2.6
1	B	206	GLU	2.6
1	B	213	VAL	2.6
1	B	47	ASP	2.6
1	B	134	ASP	2.6
1	B	79	PHE	2.6
1	C	125	TYR	2.6
1	C	222[A]	SER	2.6
1	A	40	ARG	2.5
1	B	190	LEU	2.5
1	C	155	SER	2.5
1	A	224	LEU	2.5
1	C	205	GLN	2.5
1	A	171[A]	PHE	2.4
1	B	210	LEU	2.4
1	A	42[A]	ARG	2.4
1	B	132	ILE	2.4
1	B	221	TRP	2.4
1	B	155	SER	2.3
1	B	195	GLN	2.3
1	C	202	ALA	2.3
1	C	219	ARG	2.3
1	C	163	LEU	2.3
1	C	200	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	69	GLN	2.3
1	B	21	LEU	2.2
1	B	93	PHE	2.2
1	C	0	GLY	2.2
1	B	187	LEU	2.2
1	B	89	GLY	2.2
1	C	146	GLY	2.2
1	B	157	HIS	2.2
1	C	27	ILE	2.2
1	B	46	LEU	2.2
1	B	186	LEU	2.2
1	B	179	ARG	2.2
1	C	204	GLU	2.2
1	B	175	GLN	2.2
1	C	124	ASN	2.2
1	C	128	GLY	2.2
1	C	103	SER	2.2
1	A	27	ILE	2.1
1	C	213	VAL	2.1
1	B	194	ILE	2.1
1	C	140	SER	2.1
1	C	197	LEU	2.1
1	A	103	SER	2.1
1	B	2	LEU	2.1
1	C	214	TYR	2.1
1	A	102	ARG	2.1
1	B	198	THR	2.0
1	C	203	ALA	2.0
1	B	217	LEU	2.0
1	B	5	ILE	2.0
1	B	214	TYR	2.0
1	C	71	VAL	2.0
1	C	102	ARG	2.0
1	B	55	ALA	2.0
1	A	12[A]	GLN	2.0
1	B	48	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	IMD	B	501	5/5	0.72	0.32	87,88,90,90	0
3	IMD	B	502	5/5	0.72	0.28	72,76,77,78	0
4	GOL	A	503	6/6	0.74	0.28	79,81,84,84	0
4	GOL	A	505	6/6	0.79	0.21	74,80,82,83	0
4	GOL	A	507	6/6	0.80	0.21	76,84,85,87	0
3	IMD	C	502	5/5	0.81	0.25	62,63,65,68	0
4	GOL	A	506	6/6	0.83	0.19	79,83,85,88	0
4	GOL	C	503	6/6	0.87	0.25	92,96,97,99	0
3	IMD	A	502	5/5	0.88	0.17	46,48,51,53	0
4	GOL	A	504	6/6	0.91	0.14	70,75,78,82	0
3	IMD	C	501	5/5	0.93	0.12	48,49,53,53	0
3	IMD	A	501	5/5	0.93	0.17	36,40,45,46	0
2	NI	B	500	1/1	0.94	0.10	59,59,59,59	1
2	NI	A	500	1/1	0.99	0.06	42,42,42,42	0
2	NI	C	500	1/1	0.99	0.05	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.