



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:59 AM UTC

PDB ID : 4DKP / pdb_00004dkp
Title : Crystal structure of clade A/E 93TH057 HIV-1 gp120 core in complex with AWS-I-50
Authors : Kwon, Y.D.; LaLonde, J.M.; Jones, D.M.; Sun, A.W.; Courter, J.R.; Soeta, T.; Kobayashi, T.; Princiotta, A.M.; Wu, X.; Mascola, J.; Schon, A.; Freire, E.; Sodroski, J.; Madani, N.; Smith III, A.B.; Kwong, P.D.
Deposited on : 2012-02-03
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	9% 88% 7% .
1	C	353	13% 90% 6% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6171 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	0	0
			2654	1666	460	507	21			
1	C	339	Total	C	N	O	S	0	0	0
			2654	1666	460	507	21			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	C	1	Total	C	N	O	S	0	0
			13	8	2	2	1		

- Molecule 4 is N-[(1S,2S)-2-amino-2,3-dihydro-1H-inden-1-yl]-N'-(4-chloro-3-fluorophenyl)ethanediamide (CCD ID: 0LL) (formula: $C_{17}H_{15}ClFN_3O_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	Cl	F	N	O	0	0
			24	17	1	1	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	C	1	Total	C	Cl	F	N	O	0	0
			24	17	1	1	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	296	Total	O	0	0
			296	296		
5	C	183	Total	O	0	0
			183	183		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



- Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.67Å 68.48Å 94.75Å 90.00° 91.60° 90.00°	Depositor
Resolution (Å)	33.70 – 1.80 33.70 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.3 (33.70-1.80) 89.9 (33.70-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.7.3_928, REFMAC	Depositor
R, R_{free}	0.199 , 0.225 0.194 , 0.222	Depositor DCC
R_{free} test set	3784 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.322	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6171	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0LL, NAG, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2709	0.67	1/3678 (0.0%)
1	C	0.26	0/2709	0.65	0/3678
All	All	0.27	0/5418	0.66	1/7356 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	PRO	O-C-N	5.01	123.50	121.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2654	0	2589	21	0
1	C	2654	0	2589	13	0
2	A	154	0	143	3	0
2	C	154	0	143	1	0
3	A	15	0	17	0	0
3	C	13	0	17	0	0
4	A	24	0	15	1	0
4	C	24	0	15	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	296	0	0	13	0
5	C	183	0	0	4	0
All	All	6171	0	5528	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:GLY:O	5:A:659:HOH:O	1.94	0.85
1:A:350:LYS:NZ	5:A:622:HOH:O	2.17	0.76
1:A:246:GLN:OE1	5:A:681:HOH:O	2.05	0.74
1:A:389:GLN:HG2	2:A:510:NAG:H81	1.70	0.72
1:A:240:LYS:O	5:A:766:HOH:O	2.11	0.68
1:A:344:GLN:OE1	5:A:682:HOH:O	2.12	0.67
4:A:513:OLL:H3	5:A:868:HOH:O	1.94	0.66
1:C:270:ILE:O	1:C:348:LYS:NZ	2.31	0.63
1:A:432:GLN:OE1	5:A:886:HOH:O	2.15	0.63
1:A:333:ILE:HD13	1:A:390:LEU:HD21	1.82	0.62
1:A:342:LEU:HD23	1:A:396:ILE:HD11	1.83	0.60
2:A:508:NAG:O6	5:A:874:HOH:O	2.14	0.59
1:A:121:LYS:HB2	1:A:201:ILE:HB	1.85	0.59
1:A:389:GLN:HG3	5:A:610:HOH:O	2.04	0.58
2:C:506:NAG:H5	5:C:742:HOH:O	2.04	0.57
1:A:270:ILE:HB	1:A:348:LYS:HG3	1.87	0.55
1:C:88:ASN:ND2	5:C:782:HOH:O	2.40	0.55
1:A:472:GLY:N	5:A:850:HOH:O	2.29	0.55
1:A:82:GLN:HE22	1:A:246:GLN:HG2	1.72	0.54
1:C:477:ASP:OD1	5:C:780:HOH:O	2.19	0.54
1:A:391:PHE:O	5:A:698:HOH:O	2.19	0.52
1:C:348:LYS:NZ	5:C:720:HOH:O	2.34	0.51
1:C:101:VAL:HG21	1:C:480:ARG:HG2	1.98	0.46
1:C:492:GLU:N	1:C:492:GLU:OE1	2.48	0.46
1:C:45:TRP:NE1	1:C:91:GLU:OE2	2.45	0.45
1:A:343:LYS:O	1:A:346:THR:OG1	2.35	0.45
2:A:504:NAG:HN2	1:C:61:HIS:CG	2.37	0.43
1:A:348:LYS:HG2	5:A:801:HOH:O	2.18	0.43
1:C:46:LYS:NZ	1:C:490:GLN:OE1	2.51	0.43
1:A:93:PHE:HB2	1:A:233:PHE:HZ	1.82	0.43
1:C:378:CYS:HB3	1:C:383:PHE:CE1	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:390:LEU:HG	1:C:416:LEU:HD11	2.01	0.42
1:A:207:LYS:HD2	1:A:439:ILE:HG23	2.03	0.41
1:C:370:GLU:HG2	4:C:513:OLL:N14	2.36	0.40
1:A:361:PHE:HB3	1:A:391:PHE:HB3	2.02	0.40
1:A:472:GLY:HA3	5:A:659:HOH:O	2.21	0.40
1:C:390:LEU:HD23	1:C:390:LEU:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/353 (94%)	326 (98%)	7 (2%)	0	100	100
1	C	333/353 (94%)	321 (96%)	12 (4%)	0	100	100
All	All	666/706 (94%)	647 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/311 (97%)	301 (99%)	2 (1%)	76	73
1	C	303/311 (97%)	299 (99%)	4 (1%)	61	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	606/622 (97%)	600 (99%)	6 (1%)	68	64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	219	THR
1	C	224	VAL
1	C	411	ASN
1	C	461	ASN
1	C	463	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	GLN
1	A	344	GLN
1	C	61	HIS
1	C	82	GLN
1	C	92	ASN
1	C	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	509	1	14,14,15	0.50	0	17,19,21	0.66	0
3	EPE	C	512	-	11,13,15	0.40	0	14,15,20	1.61	3 (21%)
2	NAG	C	507	1	14,14,15	0.52	0	17,19,21	0.68	0
2	NAG	A	510	1	14,14,15	0.47	0	17,19,21	0.81	0
2	NAG	C	501	1	14,14,15	0.49	0	17,19,21	0.70	0
2	NAG	C	511	1	14,14,15	0.50	0	17,19,21	0.59	0
2	NAG	A	508	1	14,14,15	0.56	0	17,19,21	0.72	0
2	NAG	A	502	1	14,14,15	0.51	0	17,19,21	0.62	0
3	EPE	A	512	-	15,15,15	0.76	1 (6%)	19,20,20	1.67	5 (26%)
2	NAG	A	506	1	14,14,15	0.48	0	17,19,21	0.82	0
2	NAG	A	507	1	14,14,15	0.52	0	17,19,21	0.77	1 (5%)
2	NAG	C	503	1	14,14,15	0.44	0	17,19,21	0.92	1 (5%)
2	NAG	C	510	1	14,14,15	0.52	0	17,19,21	0.59	0
2	NAG	A	503	1	14,14,15	0.48	0	17,19,21	0.80	0
2	NAG	A	501	1	14,14,15	0.49	0	17,19,21	0.74	0
2	NAG	C	506	1	14,14,15	0.49	0	17,19,21	0.63	0
2	NAG	C	502	1	14,14,15	0.53	0	17,19,21	0.68	0
2	NAG	A	504	1	14,14,15	0.52	0	17,19,21	0.56	0
2	NAG	A	511	1	14,14,15	0.50	0	17,19,21	0.60	0
2	NAG	A	509	1	14,14,15	0.50	0	17,19,21	0.64	0
2	NAG	C	508	1	14,14,15	0.49	0	17,19,21	0.92	1 (5%)
4	OLL	A	513	-	26,26,26	1.40	4 (15%)	33,37,37	2.33	8 (24%)
4	OLL	C	513	-	26,26,26	1.50	4 (15%)	33,37,37	2.11	7 (21%)
2	NAG	C	504	1	14,14,15	0.53	0	17,19,21	0.67	0
2	NAG	C	505	1	14,14,15	0.48	0	17,19,21	0.72	0
2	NAG	A	505	1	14,14,15	0.51	0	17,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	509	1	-	0/6/23/26	0/1/1/1
3	EPE	C	512	-	-	1/6/17/19	0/1/1/1
2	NAG	C	507	1	-	2/6/23/26	0/1/1/1
2	NAG	A	510	1	-	3/6/23/26	0/1/1/1
2	NAG	C	501	1	-	0/6/23/26	0/1/1/1
2	NAG	C	511	1	-	2/6/23/26	0/1/1/1
2	NAG	A	508	1	-	2/6/23/26	0/1/1/1
2	NAG	A	502	1	-	0/6/23/26	0/1/1/1
3	EPE	A	512	-	-	2/9/19/19	0/1/1/1
2	NAG	A	506	1	-	0/6/23/26	0/1/1/1
2	NAG	A	507	1	-	0/6/23/26	0/1/1/1
2	NAG	C	503	1	-	0/6/23/26	0/1/1/1
2	NAG	C	510	1	-	2/6/23/26	0/1/1/1
2	NAG	A	503	1	-	0/6/23/26	0/1/1/1
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	C	506	1	-	0/6/23/26	0/1/1/1
2	NAG	C	502	1	-	0/6/23/26	0/1/1/1
2	NAG	A	504	1	-	0/6/23/26	0/1/1/1
2	NAG	A	511	1	-	2/6/23/26	0/1/1/1
2	NAG	A	509	1	-	0/6/23/26	0/1/1/1
2	NAG	C	508	1	-	2/6/23/26	0/1/1/1
4	OLL	A	513	-	-	1/12/24/24	0/3/3/3
4	OLL	C	513	-	-	0/12/24/24	0/3/3/3
2	NAG	C	504	1	-	0/6/23/26	0/1/1/1
2	NAG	C	505	1	-	0/6/23/26	0/1/1/1
2	NAG	A	505	1	-	0/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	513	OLL	C07-C08	3.62	1.59	1.55
4	A	513	OLL	C07-C08	3.13	1.58	1.55
4	A	513	OLL	C13-N14	2.97	1.41	1.35
4	C	513	OLL	C13-N14	2.84	1.41	1.35
3	A	512	EPE	C10-S	2.58	1.81	1.77
4	C	513	OLL	C12-N11	2.55	1.39	1.34
4	A	513	OLL	C12-N11	2.16	1.38	1.34
4	A	513	OLL	C18-C15	2.01	1.42	1.39
4	C	513	OLL	C12-C13	2.00	1.57	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	513	OLL	C08-C07-N11	-7.87	106.41	115.01
4	C	513	OLL	C12-C13-N14	7.81	125.44	112.25
4	A	513	OLL	C12-C13-N14	7.55	125.00	112.25
4	C	513	OLL	C08-C07-N11	-5.12	109.42	115.01
3	A	512	EPE	C5-N4-C3	4.68	118.91	108.84
3	C	512	EPE	C5-N4-C3	4.26	118.03	108.84
4	A	513	OLL	O16-C13-C12	-4.22	114.38	121.24
4	C	513	OLL	O16-C13-C12	-4.19	114.43	121.24
4	A	513	OLL	C13-C12-N11	2.65	120.08	113.73
4	C	513	OLL	C18-C15-C22	-2.58	116.53	119.66
4	C	513	OLL	C13-C12-N11	2.56	119.87	113.73
4	A	513	OLL	C18-C15-C22	-2.40	116.75	119.66
4	A	513	OLL	C02-C03-C04	-2.38	118.04	120.80
4	C	513	OLL	C02-C03-C04	-2.34	118.09	120.80
3	A	512	EPE	O1S-S-C10	2.27	110.15	106.73
3	A	512	EPE	C5-C6-N1	-2.24	106.14	110.65
2	A	507	NAG	C1-O5-C5	2.23	115.17	112.19
3	C	512	EPE	C7-N4-C3	2.23	117.17	111.24
3	C	512	EPE	C7-N4-C5	2.23	117.17	111.24
3	A	512	EPE	C7-N4-C3	2.21	117.12	111.24
4	C	513	OLL	O17-C12-C13	-2.14	117.76	121.24
4	A	513	OLL	O17-C12-C13	-2.12	117.79	121.24
3	A	512	EPE	C7-N4-C5	2.07	116.77	111.24
4	A	513	OLL	C15-C22-C21	2.05	120.53	118.82
2	C	503	NAG	C1-O5-C5	2.04	114.92	112.19
2	C	508	NAG	C1-O5-C5	2.03	114.91	112.19

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	511	NAG	C8-C7-N2-C2
2	A	511	NAG	O7-C7-N2-C2
2	C	510	NAG	C8-C7-N2-C2
2	C	510	NAG	O7-C7-N2-C2
4	A	513	OLL	C03-C07-N11-C12
2	A	508	NAG	O5-C5-C6-O6
2	A	508	NAG	C4-C5-C6-O6
2	C	508	NAG	O5-C5-C6-O6
2	C	508	NAG	C4-C5-C6-O6
2	A	510	NAG	C8-C7-N2-C2
2	A	510	NAG	O7-C7-N2-C2
2	C	507	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	507	NAG	O7-C7-N2-C2
2	C	511	NAG	C8-C7-N2-C2
2	C	511	NAG	O7-C7-N2-C2
2	A	510	NAG	O5-C5-C6-O6
3	A	512	EPE	N4-C7-C8-O8
3	A	512	EPE	C10-C9-N1-C2
3	C	512	EPE	N4-C7-C8-O8

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	510	NAG	1	0
2	A	508	NAG	1	0
2	C	506	NAG	1	0
2	A	504	NAG	1	0
4	A	513	OLL	1	0
4	C	513	OLL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	339/353 (96%)	0.43	31 (9%) 15 13	17, 29, 61, 90	0
1	C	339/353 (96%)	0.89	46 (13%) 7 5	21, 43, 83, 111	0
All	All	678/706 (96%)	0.66	77 (11%) 10 8	17, 35, 75, 111	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	396	ILE	5.0
1	A	430	THR	4.9
1	C	471	GLY	4.8
1	A	473	GLY	4.7
1	C	44	VAL	4.6
1	A	472	GLY	4.6
1	C	460	ALA	4.5
1	C	492	GLU	4.2
1	C	123	THR	4.1
1	C	429	GLY	3.7
1	A	44	VAL	3.7
1	A	200	VAL	3.7
1	A	124	GLY	3.7
1	A	123	THR	3.7
1	A	396	ILE	3.6
1	C	430	THR	3.6
1	C	459	GLY	3.6
1	C	119	CYS	3.6
1	A	411	ASN	3.5
1	C	224	VAL	3.5
1	C	473	GLY	3.4
1	A	324	GLY	3.3
1	C	61	HIS	3.3
1	C	359	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	324	GLY	3.1
1	A	119	CYS	3.0
1	A	471	GLY	3.0
1	C	360	ILE	3.0
1	A	45	TRP	3.0
1	A	300	SER	3.0
1	A	392	ASN	3.0
1	A	394	THR	2.9
1	A	82	GLN	2.9
1	A	492	GLU	2.9
1	C	89	VAL	2.8
1	C	277	LEU	2.8
1	A	89	VAL	2.7
1	C	472	GLY	2.7
1	C	411	ASN	2.7
1	A	198	GLY	2.7
1	C	278	THR	2.7
1	A	301	ASN	2.6
1	C	301	ASN	2.6
1	A	366	GLY	2.6
1	A	201	ILE	2.6
1	C	232	ASN	2.6
1	C	491	ILE	2.5
1	C	124	GLY	2.5
1	A	121	LYS	2.5
1	C	45	TRP	2.5
1	C	395	CYS	2.5
1	C	349	LEU	2.4
1	A	445	CYS	2.4
1	C	79	PRO	2.4
1	C	235	GLY	2.4
1	A	116	LEU	2.4
1	C	85	HIS	2.4
1	A	87	GLU	2.3
1	C	353	PHE	2.3
1	C	355	ASN	2.3
1	C	120	VAL	2.3
1	C	84	ILE	2.2
1	C	336	THR	2.2
1	A	429	GLY	2.2
1	C	271	ILE	2.2
1	C	350	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	236	THR	2.2
1	C	463	THR	2.2
1	C	198	GLY	2.1
1	A	393	ASN	2.1
1	C	81	PRO	2.1
1	A	371	ILE	2.1
1	C	266	ALA	2.1
1	C	431	GLY	2.0
1	C	366	GLY	2.0
1	A	395	CYS	2.0
1	C	228	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	502	14/15	0.30	0.22	79,100,108,108	0
2	NAG	A	511	14/15	0.38	0.20	73,84,96,97	0
2	NAG	C	508	14/15	0.43	0.16	82,89,100,105	0
2	NAG	C	510	14/15	0.48	0.15	70,80,92,98	0
2	NAG	C	507	14/15	0.53	0.18	63,83,94,96	0
2	NAG	A	502	14/15	0.56	0.18	48,69,79,85	0
2	NAG	A	510	14/15	0.59	0.21	93,99,107,108	0
2	NAG	C	504	14/15	0.65	0.16	63,75,81,82	0
2	NAG	A	507	14/15	0.66	0.20	45,60,74,83	0
4	0LL	C	513	24/24	0.69	0.21	32,77,83,86	0
2	NAG	C	511	14/15	0.78	0.14	37,53,63,69	0
4	0LL	A	513	24/24	0.78	0.16	37,57,74,78	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	506	14/15	0.78	0.13	34,50,63,64	0
2	NAG	C	509	14/15	0.81	0.14	41,50,61,64	0
2	NAG	A	508	14/15	0.82	0.14	26,44,60,70	0
2	NAG	A	501	14/15	0.86	0.10	34,45,54,57	0
2	NAG	C	505	14/15	0.86	0.10	34,45,49,50	0
2	NAG	C	501	14/15	0.87	0.10	48,61,71,74	0
2	NAG	A	504	14/15	0.89	0.10	22,40,49,49	0
2	NAG	A	506	14/15	0.89	0.11	31,46,60,61	0
2	NAG	A	505	14/15	0.90	0.11	28,40,50,58	0
2	NAG	A	509	14/15	0.92	0.09	33,47,52,62	0
2	NAG	A	503	14/15	0.93	0.07	19,26,30,36	0
3	EPE	C	512	13/15	0.94	0.09	23,25,32,37	0
3	EPE	A	512	15/15	0.97	0.08	19,22,27,33	0
2	NAG	C	503	14/15	0.97	0.05	19,23,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 0LL C 513:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 0LL A 513:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.