



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:44 PM UTC

PDB ID : 8DK9 / pdb_00008dk9
Title : Sliding-clamp-DinX peptide
Authors : Kapur, M.K.; Gray, O.J.; Honzatko, R.H.; Nelson, S.N.
Deposited on : 2022-07-05
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	 2% 91% 6%
1	B	397	 % 92% 6%
1	C	397	 4% 89% 9%
1	D	397	 2% 92% 6%
2	E	6	 33% 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	6	 83% 17%
2	G	6	 17% 83% 17%
2	H	6	 83% 17%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 24227 atoms, of which 11854 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	388	5789	1815	2910	493	564	7	0	1	0
1	B	388	5824	1828	2928	496	565	7	0	4	0
1	C	388	5816	1824	2923	496	566	7	0	3	0
1	D	388	5788	1815	2909	493	564	7	0	1	0

- Molecule 2 is a protein called DNA polymerase IV 1 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	E	6	92	31	44	7	10	0	0	0
2	F	6	92	31	44	7	10	0	0	0
2	G	6	92	31	44	7	10	0	0	0
2	H	6	92	31	44	7	10	0	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ACETYL GROUP (CCD ID: ACE) (formula: C₂H₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			3	2	1		
5	F	1	Total	C	O	0	0
			3	2	1		
5	G	1	Total	C	O	0	0
			3	2	1		
5	H	1	Total	C	O	0	0
			3	2	1		

- Molecule 6 is AMINO GROUP (CCD ID: NH2) (formula: H₂N) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	E	1	Total	H	N	0	0
			3	2	1		
6	F	1	Total	H	N	0	0
			3	2	1		
6	G	1	Total	H	N	0	0
			3	2	1		
6	H	1	Total	H	N	0	0
			3	2	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	152	Total	O	0	0
			152	152		
7	B	189	Total	O	0	0
			189	189		
7	E	3	Total	O	0	0
			3	3		
7	F	5	Total	O	0	0
			5	5		
7	C	113	Total	O	0	0
			113	113		
7	D	126	Total	O	0	0
			126	126		
7	G	4	Total	O	0	0
			4	4		
7	H	5	Total	O	0	0
			5	5		




- Molecule 2: DNA polymerase IV 1 peptide




- Molecule 2: DNA polymerase IV 1 peptide



- Molecule 2: DNA polymerase IV 1 peptide



- Molecule 2: DNA polymerase IV 1 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.07Å 73.94Å 81.34Å 82.77° 77.57° 87.75°	Depositor
Resolution (Å)	47.35 – 2.50 47.35 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.0 (47.35-2.50) 95.0 (47.35-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.211 , 0.264 0.210 , 0.264	Depositor DCC
R_{free} test set	2005 reflections (3.60%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtrriage
Anisotropy	0.239	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24227	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACE, NH2, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/2934	0.25	0/4007
1	B	0.10	0/2961	0.25	0/4044
1	C	0.10	0/2954	0.25	0/4033
1	D	0.10	0/2934	0.25	0/4007
2	E	0.07	0/48	0.29	0/63
2	F	0.10	0/48	0.26	0/63
2	G	0.14	0/48	0.51	0/63
2	H	0.50	0/48	0.63	0/63
All	All	0.11	0/11975	0.26	0/16343

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2879	2910	2909	18	0
1	B	2896	2928	2933	14	0
1	C	2893	2923	2928	25	0
1	D	2879	2909	2909	16	0
2	E	48	44	43	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	48	44	43	0	0
2	G	48	44	43	1	0
2	H	48	44	43	0	0
3	A	5	0	0	0	0
3	B	10	0	0	0	0
3	D	5	0	0	0	0
4	B	1	0	0	0	0
5	E	3	0	3	0	0
5	F	3	0	3	0	0
5	G	3	0	3	0	0
5	H	3	0	3	0	0
6	E	1	2	0	0	0
6	F	1	2	0	0	0
6	G	1	2	0	0	0
6	H	1	2	0	0	0
7	A	152	0	0	0	0
7	B	189	0	0	0	0
7	C	113	0	0	0	0
7	D	126	0	0	0	0
7	E	3	0	0	0	0
7	F	5	0	0	0	0
7	G	4	0	0	0	0
7	H	5	0	0	0	0
All	All	12373	11854	11863	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:VAL:HG22	1:A:191:TRP:HB2	1.69	0.73
1:B:295:VAL:HG23	1:B:335:PRO:HG3	1.70	0.71
1:B:373:LEU:HD12	1:B:373:LEU:O	1.92	0.69
1:B:142:PHE:CZ	1:B:146:ILE:HD11	2.29	0.68
1:C:237:LEU:HD21	1:C:248:THR:HB	1.79	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/397 (98%)	372 (96%)	15 (4%)	0	100	100
1	B	390/397 (98%)	376 (96%)	14 (4%)	0	100	100
1	C	389/397 (98%)	369 (95%)	20 (5%)	0	100	100
1	D	387/397 (98%)	369 (95%)	18 (5%)	0	100	100
2	E	4/6 (67%)	4 (100%)	0	0	100	100
2	F	4/6 (67%)	4 (100%)	0	0	100	100
2	G	4/6 (67%)	4 (100%)	0	0	100	100
2	H	4/6 (67%)	4 (100%)	0	0	100	100
All	All	1569/1612 (97%)	1502 (96%)	67 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/311 (98%)	305 (100%)	1 (0%)	86	94
1	B	309/311 (99%)	309 (100%)	0	100	100
1	C	308/311 (99%)	308 (100%)	0	100	100
1	D	306/311 (98%)	304 (99%)	2 (1%)	76	89
2	E	5/5 (100%)	5 (100%)	0	100	100
2	F	5/5 (100%)	4 (80%)	1 (20%)	1	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	G	5/5 (100%)	4 (80%)	1 (20%)	1 2
2	H	5/5 (100%)	4 (80%)	1 (20%)	1 2
All	All	1249/1264 (99%)	1243 (100%)	6 (0%)	81 92

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	223	GLN
2	G	6	PHE
2	H	6	PHE
2	F	6	PHE
1	A	161	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	223	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic and 4 are modelled with single atom - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACE	E	101	2	1,2,2	0.81	0	0,1,1	-	-
5	ACE	H	101	2	1,2,2	0.82	0	0,1,1	-	-
5	ACE	F	101	2	1,2,2	0.81	0	0,1,1	-	-
3	SO4	A	401	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	D	401	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	B	401	-	4,4,4	0.24	0	6,6,6	0.09	0
5	ACE	G	101	2	1,2,2	0.81	0	0,1,1	-	-
3	SO4	B	402	-	4,4,4	0.24	0	6,6,6	0.09	0

There are no bond length outliers.

There are no bond angle outliers.




There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/397 (97%)	0.08	6 (1%) 72 68	36, 56, 84, 111	1 (0%)
1	B	388/397 (97%)	-0.04	5 (1%) 75 71	22, 53, 81, 94	4 (1%)
1	C	388/397 (97%)	0.34	14 (3%) 46 41	25, 70, 104, 116	3 (0%)
1	D	388/397 (97%)	0.23	9 (2%) 61 57	32, 60, 92, 109	1 (0%)
2	E	6/6 (100%)	1.03	2 (33%) 1 1	59, 67, 74, 93	0
2	F	6/6 (100%)	-0.12	0 100 100	49, 50, 56, 59	0
2	G	6/6 (100%)	0.73	1 (16%) 4 3	46, 59, 66, 86	0
2	H	6/6 (100%)	0.26	0 100 100	65, 72, 78, 82	0
All	All	1576/1612 (97%)	0.16	37 (2%) 61 57	22, 58, 96, 116	9 (0%)

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	233	GLY	3.4
1	C	217	LEU	3.3
1	C	195	THR	3.2
1	A	229	GLY	3.1
1	D	33	SER	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NH2	G	102	1/1	0.41	0.12	87,87,105,105	0
6	NH2	H	102	1/1	0.58	0.10	83,83,99,99	0
3	SO4	B	402	5/5	0.71	0.25	38,46,54,61	5
3	SO4	D	401	5/5	0.71	0.16	64,65,73,73	5
3	SO4	B	401	5/5	0.73	0.22	52,55,56,57	5
3	SO4	A	401	5/5	0.79	0.15	58,58,61,63	5
6	NH2	E	102	1/1	0.80	0.12	109,109,131,131	0
6	NH2	F	102	1/1	0.80	0.14	68,68,82,82	0
5	ACE	H	101	3/3	0.86	0.15	69,69,72,73	0
5	ACE	E	101	3/3	0.88	0.12	59,59,61,62	0
5	ACE	G	101	3/3	0.93	0.13	41,41,44,46	0
5	ACE	F	101	3/3	0.95	0.08	49,49,49,49	0
4	CL	B	403	1/1	0.98	0.09	43,43,43,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around SO4 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around SO4 A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NH2 E 102:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NH2 F 102:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around ACE E 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ACE G 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ACE F 101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.