



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:04 AM UTC

PDB ID : 8DKS / pdb_00008dks
Title : IRAK4 IN COMPLEX WITH COMPOUND #3
Authors : Chen, Y.; Lin, N.
Deposited on : 2022-07-06
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	 3% 58% 38%
1	B	460	 7% 57% 40%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	P	S	122	0	0
			2226	1400	374	437	1	14			
1	B	278	Total	C	N	O	P	S	143	1	0
			2204	1387	373	428	1	15			

- Molecule 2 is (1S,2S,3R,4R)-3-({2-[3-(pyrrolidine-1-carbonyl)anilino]thieno[3,2-d]pyrimidin-4-yl}amino)bicyclo[2.2.1]hept-5-ene-2-carboxamide (CCD ID: SO9) (formula: C₂₅H₂₆N₆O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			34	25	6	2	1		
2	B	1	Total	C	N	O	S	0	0
			34	25	6	2	1		

- Molecule 3 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total	O	0	0
			26	26		
4	B	29	Total	O	0	0
			29	29		

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	88.91Å 119.49Å 139.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.71 – 2.45 90.71 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.5 (90.71-2.45) 97.5 (90.71-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.214 , 0.267 0.272 , 0.313	Depositor DCC
R_{free} test set	907 reflections (3.28%)	wwPDB-VP
Wilson B-factor (Å ²)	55.9	Xtriage
Anisotropy	0.565	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4561	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BME, SO9, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2252	0.85	1/3034 (0.0%)
1	B	0.65	0/2229	0.86	1/2998 (0.0%)
All	All	0.64	0/4481	0.85	2/6032 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	256	ASP	CA-CB-CG	6.60	119.20	112.60
1	B	256	ASP	CA-CB-CG	5.15	117.75	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2226	0	2195	8	0
1	B	2204	0	2176	10	0
2	A	34	0	0	0	0
2	B	34	0	0	1	0
3	A	4	0	5	0	0
3	B	4	0	5	0	0
4	A	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	29	0	0	0	0
All	All	4561	0	4381	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:MET:HE2	1:A:363:GLU:CG	2.33	0.58
1:B:283:LEU:HD23	1:B:287[B]:MET:HE2	1.85	0.57
1:B:200:VAL:HG22	1:B:213:LYS:HG3	1.90	0.54
1:B:246:VAL:HG11	1:B:318:LEU:HD12	1.89	0.54
1:A:265:MET:HE2	1:A:265:MET:HA	1.89	0.53
1:B:265:MET:HE2	1:B:265:MET:HA	1.90	0.52
1:B:173:LEU:HA	1:B:176:VAL:HG12	1.95	0.49
1:A:246:VAL:HG11	1:A:318:LEU:HD12	1.95	0.48
1:B:283:LEU:CD2	1:B:287[B]:MET:HE2	2.44	0.47
1:B:280:THR:HG21	2:B:501:SO9:C33	2.44	0.47
1:A:233:GLU:HG2	1:A:260:LEU:HD13	1.96	0.46
1:A:344:MET:HE2	1:A:363:GLU:HG2	1.97	0.46
1:B:246:VAL:CG1	1:B:318:LEU:HD12	2.46	0.45
1:A:298:GLY:HA3	1:A:327:ILE:HD12	1.99	0.43
1:B:455:GLN:O	1:B:458:THR:HG22	2.19	0.43
1:A:300:ASN:HA	1:A:447:ILE:HG21	2.00	0.43
1:B:288:ARG:HB3	1:B:380:ILE:CG2	2.51	0.40
1:A:308:ILE:HD12	1:A:364:ILE:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/460 (60%)	268 (98%)	7 (2%)	0	100	100
1	B	269/460 (58%)	260 (97%)	9 (3%)	0	100	100
All	All	544/920 (59%)	528 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/406 (60%)	244 (100%)	1 (0%)	84	89
1	B	242/406 (60%)	242 (100%)	0	100	100
All	All	487/812 (60%)	486 (100%)	1 (0%)	87	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	451	GLN
1	B	241	GLN
1	B	293	GLN
1	B	435	GLN
1	B	451	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	B	346	1	8,9,10	1.67	1 (12%)	7,12,14	1.29	1 (14%)
1	SEP	A	346	1	8,9,10	1.63	1 (12%)	7,12,14	1.36	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	346	1	-	1/6/8/10	-
1	SEP	A	346	1	-	0/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	346	SEP	P-O1P	3.65	1.61	1.50
1	A	346	SEP	P-O1P	3.57	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	SEP	OG-CB-CA	3.05	111.11	108.14
1	B	346	SEP	OG-CB-CA	2.70	110.77	108.14

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	346	SEP	C-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SO9	B	501	-	37,39,39	1.11	4 (10%)	48,57,57	1.70	10 (20%)
3	BME	A	502	1	3,3,3	0.51	0	2,2,2	0.21	0
2	SO9	A	501	-	37,39,39	1.10	3 (8%)	48,57,57	1.74	11 (22%)
3	BME	B	502	1	3,3,3	0.48	0	2,2,2	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO9	B	501	-	-	2/20/48/48	0/7/6/6
3	BME	A	502	1	-	1/1/1/1	-
2	SO9	A	501	-	-	2/20/48/48	0/7/6/6
3	BME	B	502	1	-	1/1/1/1	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	SO9	C16-N15	2.51	1.39	1.35
2	A	501	SO9	C20-C24	-2.44	1.39	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	SO9	C12-C10	2.39	1.59	1.54
2	B	501	SO9	C16-N15	2.20	1.38	1.35
2	B	501	SO9	C12-C10	2.18	1.59	1.54
2	B	501	SO9	C18-N25	2.14	1.41	1.36
2	B	501	SO9	C21-S22	2.01	1.76	1.73

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	SO9	C10-C13-N15	-5.33	106.34	113.60
2	B	501	SO9	C20-N19-C18	4.75	121.77	116.69
2	B	501	SO9	N19-C18-N17	-4.23	119.33	126.26
2	A	501	SO9	C4-C2-N3	3.62	121.00	115.62
2	A	501	SO9	C20-N19-C18	3.54	120.48	116.69
2	B	501	SO9	C4-C2-N3	3.32	120.55	115.62
2	A	501	SO9	N19-C18-N17	-3.30	120.85	126.26
2	B	501	SO9	C28-C29-N31	3.28	122.76	118.66
2	A	501	SO9	C33-C32-N31	3.20	108.83	103.44
2	B	501	SO9	C18-N17-C16	2.74	123.09	116.36
2	A	501	SO9	C34-C35-N31	2.70	108.00	103.44
2	B	501	SO9	C10-C13-N15	-2.66	109.98	113.60
2	A	501	SO9	C18-N17-C16	2.60	122.74	116.36
2	A	501	SO9	N15-C16-N17	2.26	122.91	118.55
2	B	501	SO9	O1-C2-C4	-2.25	118.61	121.71
2	A	501	SO9	O1-C2-N3	-2.23	119.10	123.04
2	A	501	SO9	C35-N31-C32	-2.15	108.53	111.37
2	B	501	SO9	C33-C32-N31	2.07	106.92	103.44
2	A	501	SO9	C13-C4-C2	2.06	115.27	111.42
2	B	501	SO9	C12-C6-C4	2.04	102.60	99.90
2	B	501	SO9	C35-N31-C32	-2.04	108.67	111.37

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	502	BME	O1-C1-C2-S2
2	B	501	SO9	C21-C16-N15-C13
2	A	501	SO9	N17-C18-N25-C26
2	B	501	SO9	N17-C16-N15-C13
2	A	501	SO9	N19-C18-N25-C26
3	A	502	BME	O1-C1-C2-S2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	SO9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/460 (61%)	0.56	13 (4%) 37 36	17, 36, 52, 66	43 (15%)
1	B	277/460 (60%)	0.77	33 (11%) 9 7	11, 35, 51, 73	47 (16%)
All	All	559/920 (60%)	0.66	46 (8%) 17 15	11, 35, 52, 73	90 (16%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	258	LEU	5.6
1	B	185	ILE	4.5
1	B	230	PHE	4.4
1	B	182	GLU	4.2
1	B	256	ASP	3.9
1	B	259	CYS	3.9
1	B	203	GLY	3.7
1	B	195	GLY	3.5
1	B	201	TYR	3.3
1	B	184	PRO	3.3
1	A	343	VAL	3.3
1	B	189	GLY	3.1
1	A	240	CYS	3.0
1	B	254	ASP	3.0
1	A	165	PHE	3.0
1	B	197	PHE	2.9
1	B	252	SER	2.9
1	A	254	ASP	2.8
1	B	229	GLN	2.8
1	B	257	ASP	2.8
1	A	186	SER	2.6
1	B	179	ASN	2.6
1	B	169	SER	2.5
1	B	458	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	255	GLY	2.5
1	B	407	GLU	2.5
1	B	343	VAL	2.5
1	A	179	ASN	2.4
1	A	166	HIS	2.4
1	B	226	LEU	2.4
1	A	178	ASN	2.4
1	A	222	THR	2.3
1	A	230	PHE	2.3
1	B	193	GLY	2.2
1	B	260	LEU	2.2
1	B	171	TYR	2.2
1	B	287[A]	MET	2.2
1	B	336	SER	2.2
1	B	337	GLU	2.2
1	A	256	ASP	2.2
1	B	199	VAL	2.2
1	A	187	VAL	2.1
1	A	171	TYR	2.1
1	B	215	LEU	2.1
1	B	232	GLN	2.0
1	B	181	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	B	346	10/11	0.47	0.17	46,48,58,61	0
1	SEP	A	346	10/11	0.65	0.13	49,51,55,57	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BME	A	502	4/4	0.83	0.22	72,75,78,78	0
3	BME	B	502	4/4	0.85	0.22	69,70,71,73	0
2	SO9	B	501	34/34	0.93	0.13	32,46,51,59	0
2	SO9	A	501	34/34	0.94	0.12	33,45,51,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SO9 B 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SO9 A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.