



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 03:00 AM UTC

PDB ID : 1DLH / pdb_00001dlh
Title : CRYSTAL STRUCTURE OF THE HUMAN CLASS II MHC PROTEIN
HLA-DR1 COMPLEXED WITH AN INFLUENZA VIRUS PEPTIDE
Authors : Stern, L.J.
Deposited on : 1994-02-15
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	180	42% 51% 7%
1	D	180	44% 49% 7%
2	B	188	43% 49% 9%
2	E	188	37% 55% 8%
3	C	13	31% 69%
3	F	13	23% 69% 8%
4	G	2	100%
4	H	2	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	B	521	X	-	-	-
5	NAG	E	521	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6520 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	10	0	0
			1478	957	240	276	5			
1	D	180	Total	C	N	O	S	10	0	0
			1478	957	240	276	5			


- Molecule 2 is a protein called CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (BETA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	188	Total	C	N	O	S	14	0	0
			1544	973	277	288	6			
2	E	188	Total	C	N	O	S	14	0	0
			1543	973	277	287	6			

- Molecule 3 is a protein called ENTEROTOXIN TYPE B PRECURSOR.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	0	0
			106	69	18	19			
3	F	13	Total	C	N	O	0	0	0
			106	69	18	19			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	38	Total	O	0	0
			38	38		
6	B	40	Total	O	0	0
			40	40		
6	C	2	Total	O	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	34	Total 34	O 34	0	0
6	E	36	Total 36	O 36	0	0
6	F	3	Total 3	O 3	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (ALPHA CHAIN)



• Molecule 1: CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (ALPHA CHAIN)




• Molecule 2: CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (BETA CHAIN)







• Molecule 2: CLASS II HISTOCOMPATIBILITY ANTIGEN (HLA-DR1) (BETA CHAIN)



• Molecule 3: ENTEROTOXIN TYPE B PRECURSOR




• Molecule 3: ENTEROTOXIN TYPE B PRECURSOR



• Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.60Å 94.60Å 247.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-2.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.205 , 0.304	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6520	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	4/1523 (0.3%)	1.40	24/2077 (1.2%)
1	D	1.22	3/1523 (0.2%)	1.42	24/2077 (1.2%)
2	B	1.15	3/1584 (0.2%)	1.40	16/2152 (0.7%)
2	E	1.17	2/1583 (0.1%)	1.39	16/2150 (0.7%)
3	C	1.00	0/107	1.27	1/141 (0.7%)
3	F	1.30	1/107 (0.9%)	1.21	1/141 (0.7%)
All	All	1.17	13/6427 (0.2%)	1.39	82/8738 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2
2	E	0	1
All	All	0	3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	34	VAL	CA-CB	6.59	1.61	1.54
1	D	165	VAL	CA-CB	6.35	1.61	1.54
2	E	133	ARG	CA-C	-6.24	1.45	1.53
3	F	308	TYR	CA-C	-5.84	1.45	1.52
2	B	31	ILE	CA-CB	5.60	1.61	1.54

The worst 5 of 82 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	86	PRO	CA-C-N	9.43	129.26	120.21
1	D	86	PRO	C-N-CA	9.43	129.26	120.21
2	E	166	ARG	N-CA-C	-8.16	96.97	109.95
2	B	182	SER	CA-C-N	7.82	129.61	119.84
2	B	182	SER	C-N-CA	7.82	129.61	119.84

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	123	TYR	Sidechain
2	B	22	GLU	Mainchain
2	E	123	TYR	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1478	0	1410	99	0
1	D	1478	0	1410	94	0
2	B	1544	0	1476	133	0
2	E	1543	0	1476	142	0
3	C	106	0	119	13	0
3	F	106	0	119	14	0
4	G	28	0	24	0	0
4	H	28	0	24	1	0
5	A	14	0	13	2	0
5	B	14	0	13	4	0
5	D	14	0	13	0	0
5	E	14	0	13	3	0
6	A	38	0	0	18	0
6	B	40	0	0	20	0
6	C	2	0	0	0	0
6	D	34	0	0	16	0
6	E	36	0	0	14	0
6	F	3	0	0	1	0
All	All	6520	0	6110	432	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:THR:HG22	2:B:6:ARG:NH2	1.67	1.07
1:D:164:ARG:HE	1:D:173:PRO:HB3	1.25	0.99
2:E:3:THR:HG22	2:E:6:ARG:NH2	1.76	0.99
1:A:164:ARG:HE	1:A:173:PRO:HB3	1.24	0.98
1:A:175:LEU:HG	6:A:534:HOH:O	1.67	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/180 (99%)	155 (87%)	18 (10%)	5 (3%)	4	14
1	D	178/180 (99%)	159 (89%)	13 (7%)	6 (3%)	3	11
2	B	186/188 (99%)	163 (88%)	19 (10%)	4 (2%)	5	19
2	E	186/188 (99%)	160 (86%)	20 (11%)	6 (3%)	3	11
3	C	11/13 (85%)	11 (100%)	0	0	100	100
3	F	11/13 (85%)	11 (100%)	0	0	100	100
All	All	750/762 (98%)	659 (88%)	70 (9%)	21 (3%)	4	14

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	THR
1	D	78	ASN
1	D	111	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	45	GLY
1	A	17	ASP

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	1	1,4	14,14,15	1.09	0	17,19,21	1.48	3 (17%)
4	NDG	G	2	4	14,14,15	1.26	1 (7%)	17,19,21	1.12	1 (5%)
4	NAG	H	1	1,4	14,14,15	0.90	1 (7%)	17,19,21	1.14	1 (5%)
4	NDG	H	2	4	14,14,15	1.25	1 (7%)	17,19,21	1.15	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NDG	G	2	4	-	0/6/23/26	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NDG	H	2	4	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	2	NDG	C1-C2	4.14	1.58	1.52
4	G	2	NDG	C1-C2	3.87	1.57	1.52
4	H	1	NAG	O4-C4	2.07	1.48	1.43

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1	NAG	C4-C3-C2	-4.09	105.03	111.02
4	G	2	NDG	C4-C3-C2	3.74	116.50	111.02
4	H	2	NDG	C3-C4-C5	-2.81	105.14	110.23
4	H	1	NAG	C1-C2-N2	-2.52	106.47	110.43
4	G	1	NAG	O4-C4-C3	2.51	116.29	110.38

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	2	NDG	C1-C2-N2-C7
4	H	2	NDG	C3-C2-N2-C7


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	E	521	2	14,14,15	1.30	2 (14%)	17,19,21	0.88	1 (5%)
5	NAG	A	501	1	14,14,15	1.38	1 (7%)	17,19,21	1.36	1 (5%)
5	NAG	D	501	1	14,14,15	1.41	1 (7%)	17,19,21	1.19	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	521	2	14,14,15	1.28	1 (7%)	17,19,21	1.74	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	521	2	1/1/5/7	2/6/23/26	0/1/1/1
5	NAG	A	501	1	-	2/6/23/26	0/1/1/1
5	NAG	D	501	1	-	2/6/23/26	0/1/1/1
5	NAG	B	521	2	1/1/5/7	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	501	NAG	C1-C2	4.71	1.58	1.52
5	A	501	NAG	C1-C2	4.05	1.57	1.52
5	B	521	NAG	C1-C2	3.75	1.57	1.52
5	E	521	NAG	C1-C2	3.38	1.56	1.52
5	E	521	NAG	O5-C1	2.64	1.48	1.43

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	521	NAG	C1-O5-C5	5.39	119.42	112.19
5	A	501	NAG	C4-C3-C2	-3.54	105.83	111.02
5	E	521	NAG	C1-O5-C5	3.07	116.30	112.19
5	D	501	NAG	C1-O5-C5	2.82	115.96	112.19
5	B	521	NAG	O5-C1-C2	2.76	115.56	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	521	NAG	C1
5	E	521	NAG	C1

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	521	NAG	C4-C5-C6-O6
5	B	521	NAG	O5-C5-C6-O6
5	E	521	NAG	O5-C5-C6-O6
5	A	501	NAG	O5-C5-C6-O6
5	A	501	NAG	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	521	NAG	3	0
5	A	501	NAG	2	0
5	B	521	NAG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.