



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:52 AM UTC

PDB ID : 5DL7 / pdb_00005dl7
Title : Crystal structure of Acinetobacter baumannii OccAB3
Authors : Zahn, M.; Basle, A.; van den Berg, B.
Deposited on : 2015-09-04
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	5% 73% 21% • •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Porin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	402	Total	C	N	O	S	0	0	0
			3260	2070	559	623	8			


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	ALA	-	expression tag	UNP A0A0D5YKR4
A	-13	ASN	-	expression tag	UNP A0A0D5YKR4
A	-12	VAL	-	expression tag	UNP A0A0D5YKR4
A	-11	ARG	-	expression tag	UNP A0A0D5YKR4
A	-10	LEU	-	expression tag	UNP A0A0D5YKR4
A	-9	GLN	-	expression tag	UNP A0A0D5YKR4
A	-8	HIS	-	expression tag	UNP A0A0D5YKR4
A	-7	HIS	-	expression tag	UNP A0A0D5YKR4
A	-6	HIS	-	expression tag	UNP A0A0D5YKR4
A	-5	HIS	-	expression tag	UNP A0A0D5YKR4
A	-4	HIS	-	expression tag	UNP A0A0D5YKR4
A	-3	HIS	-	expression tag	UNP A0A0D5YKR4
A	-2	HIS	-	expression tag	UNP A0A0D5YKR4
A	-1	LEU	-	expression tag	UNP A0A0D5YKR4
A	0	GLU	-	expression tag	UNP A0A0D5YKR4

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		

- Molecule 3 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (CCD ID: C8E) (formula: C₁₆H₃₄O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			21	16	5		
3	A	1	Total	C	O	0	0
			21	16	5		
3	A	1	Total	C	O	0	0
			16	13	3		
3	A	1	Total	C	O	0	0
			15	12	3		
3	A	1	Total	C	O	0	0
			12	8	4		
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			11	7	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	216	Total	O	0	0
			216	216		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Porin



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	81.56Å 125.72Å 137.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.14 – 1.75 70.14 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.14-1.75) 99.9 (70.14-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.168 , 0.191 0.178 , 0.198	Depositor DCC
R_{free} test set	1831 reflections (2.57%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3586	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.07	74/3337 (2.2%)	1.64	28/4498 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (74) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	PRO	N-CA	-15.21	1.36	1.47
1	A	111	ASP	CB-CG	10.65	1.78	1.52
1	A	291	ASP	CG-OD1	8.45	1.41	1.25
1	A	56	ASP	CA-C	8.31	1.62	1.52
1	A	111	ASP	CG-OD1	8.08	1.40	1.25
1	A	338	ILE	CA-CB	-7.84	1.44	1.53
1	A	13	ARG	CD-NE	-7.51	1.35	1.46
1	A	261	GLN	CA-C	7.46	1.62	1.53
1	A	181	ASP	CA-CB	-7.45	1.40	1.53
1	A	173	ARG	CZ-NH2	-7.12	1.24	1.33
1	A	317	ARG	CD-NE	-7.05	1.36	1.46
1	A	150	TYR	CE1-CZ	6.93	1.54	1.38
1	A	57	VAL	CA-CB	-6.92	1.44	1.54
1	A	79	LEU	CA-C	6.90	1.59	1.52
1	A	58	LEU	CA-C	6.90	1.60	1.52
1	A	276	THR	CA-C	-6.82	1.44	1.52
1	A	329	MET	SD-CE	6.81	1.96	1.79
1	A	161	ARG	N-CA	6.78	1.54	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	LEU	CA-CB	-6.76	1.43	1.53
1	A	123	PRO	CA-C	6.68	1.60	1.52
1	A	262	ASN	CG-ND2	-6.53	1.19	1.33
1	A	403	LYS	CA-C	-6.52	1.45	1.52
1	A	7	LYS	N-CA	-6.35	1.38	1.45
1	A	342	GLN	CA-CB	6.33	1.63	1.53
1	A	91	TYR	C-O	6.29	1.31	1.23
1	A	330	THR	N-CA	6.28	1.53	1.45
1	A	41	LYS	CA-C	-6.09	1.45	1.52
1	A	284	THR	CB-OG1	6.08	1.53	1.43
1	A	42	GLY	CA-C	6.07	1.56	1.52
1	A	162	TYR	CA-C	-5.95	1.45	1.52
1	A	329	MET	C-O	-5.90	1.17	1.23
1	A	99	LYS	CA-C	-5.87	1.45	1.52
1	A	361	ILE	CA-C	5.83	1.59	1.52
1	A	126	LEU	N-CA	-5.78	1.39	1.46
1	A	211	GLN	C-O	5.76	1.30	1.23
1	A	19	ARG	CA-C	5.71	1.60	1.53
1	A	383	PRO	N-CA	-5.70	1.40	1.47
1	A	140	GLU	C-N	-5.67	1.27	1.33
1	A	123	PRO	CA-CB	5.67	1.60	1.53
1	A	47	THR	CA-CB	5.62	1.62	1.53
1	A	5	ASP	N-CA	5.62	1.53	1.46
1	A	244	ILE	C-N	-5.61	1.25	1.33
1	A	336	GLU	CB-CG	-5.59	1.35	1.52
1	A	397	ASN	CA-C	-5.59	1.45	1.52
1	A	107	LEU	CA-C	5.54	1.59	1.52
1	A	361	ILE	C-O	-5.45	1.17	1.24
1	A	135	ARG	CZ-NH1	-5.42	1.25	1.32
1	A	214	GLY	N-CA	5.40	1.51	1.45
1	A	60	MET	CA-CB	-5.40	1.44	1.53
1	A	144	LEU	C-O	-5.39	1.17	1.24
1	A	209	GLN	CD-OE1	5.39	1.33	1.23
1	A	47	THR	CA-C	5.36	1.60	1.53
1	A	159	SER	CA-C	-5.34	1.46	1.52
1	A	33	ASP	CG-OD2	5.34	1.35	1.25
1	A	116	LEU	CA-C	-5.33	1.45	1.52
1	A	157	ARG	CA-CB	5.31	1.61	1.53
1	A	20	ASP	CA-C	-5.28	1.46	1.53
1	A	250	ARG	CA-C	-5.28	1.46	1.52
1	A	310	GLU	CB-CG	-5.27	1.36	1.52
1	A	143	ASN	CG-OD1	5.22	1.33	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	TYR	N-CA	5.19	1.52	1.45
1	A	247	VAL	C-N	-5.18	1.27	1.33
1	A	180	THR	CA-CB	5.17	1.61	1.53
1	A	290	LEU	N-CA	5.13	1.52	1.46
1	A	224	GLN	C-O	-5.11	1.17	1.24
1	A	284	THR	CA-C	5.09	1.59	1.52
1	A	138	SER	N-CA	5.08	1.52	1.46
1	A	323	LEU	N-CA	-5.07	1.39	1.46
1	A	181	ASP	CB-CG	-5.05	1.39	1.52
1	A	141	ILE	C-N	-5.03	1.27	1.33
1	A	143	ASN	N-CA	5.02	1.52	1.46
1	A	306	SER	C-N	-5.01	1.26	1.33
1	A	311	TYR	N-CA	5.01	1.52	1.46
1	A	382	MET	CG-SD	5.01	1.93	1.80

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	13	ARG	NE-CZ-NH2	17.97	135.37	119.20
1	A	13	ARG	NE-CZ-NH1	-13.58	107.92	121.50
1	A	13	ARG	CD-NE-CZ	11.43	140.41	124.40
1	A	173	ARG	NE-CZ-NH2	7.49	125.94	119.20
1	A	274	GLY	CA-C-O	-6.21	117.98	122.45
1	A	284	THR	O-C-N	6.08	130.62	122.96
1	A	49	GLY	N-CA-C	-6.04	102.06	111.00
1	A	402	PHE	CB-CA-C	-5.90	100.02	109.75
1	A	300	LYS	CB-CG-CD	5.75	124.52	111.30
1	A	180	THR	CA-C-N	-5.73	113.08	122.54
1	A	180	THR	C-N-CA	-5.73	113.08	122.54
1	A	48	GLU	N-CA-C	-5.70	102.05	110.48
1	A	198	ARG	NE-CZ-NH2	5.63	124.26	119.20
1	A	342	GLN	CG-CD-NE2	5.61	124.82	116.40
1	A	292	PHE	CA-C-N	5.58	128.21	120.29
1	A	292	PHE	C-N-CA	5.58	128.21	120.29
1	A	92	GLY	CA-C-O	-5.54	116.48	121.47
1	A	268	GLY	O-C-N	-5.46	118.56	123.37
1	A	162	TYR	CE1-CZ-OH	-5.44	103.58	119.90
1	A	304	VAL	O-C-N	5.38	129.01	123.20
1	A	25	PRO	N-CA-C	-5.36	101.44	112.47
1	A	175	ASN	OD1-CG-ND2	5.28	127.88	122.60
1	A	111	ASP	CA-C-O	-5.23	114.72	120.69
1	A	289	VAL	CG1-CB-CG2	-5.20	99.36	110.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7	LYS	N-CA-CB	-5.17	101.91	111.53
1	A	75	ARG	N-CA-CB	5.14	117.56	109.69
1	A	111	ASP	O-C-N	5.09	129.00	123.04
1	A	96	ILE	CB-CA-C	-5.02	103.09	110.82

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	TYR	Sidechain
1	A	296	ASP	Sidechain
1	A	74	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3260	0	3123	18	0
2	A	4	0	0	0	0
3	A	106	0	159	7	0
4	A	216	0	0	2	0
All	All	3586	0	3282	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ASP:CB	1:A:111:ASP:CG	1.78	1.50
3:A:505:C8E:H201	3:A:505:C8E:H161	1.53	0.88
1:A:13:ARG:HD2	1:A:399:ASP:CG	2.14	0.73
3:A:505:C8E:H161	3:A:505:C8E:C20	2.23	0.68
1:A:13:ARG:HD2	1:A:399:ASP:OD2	1.95	0.65
1:A:253:SER:HB2	1:A:289:VAL:HG12	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ASP:CG	1:A:111:ASP:CA	2.71	0.58
1:A:13:ARG:HD2	1:A:399:ASP:OD1	2.05	0.56
1:A:261:GLN:O	1:A:262:ASN:HB2	2.05	0.55
1:A:69:ARG:NH1	1:A:72:ASP:OD2	2.39	0.55
1:A:13:ARG:CD	1:A:399:ASP:OD2	2.55	0.54
1:A:361:ILE:HD12	1:A:369:LEU:HB3	1.90	0.53
3:A:505:C8E:H112	3:A:506:C8E:H102	1.95	0.48
1:A:334:LYS:NZ	4:A:604:HOH:O	2.45	0.48
3:A:506:C8E:H72	3:A:508:C8E:C10	2.44	0.48
3:A:505:C8E:C20	3:A:505:C8E:C16	2.91	0.47
1:A:190:TYR:CE2	1:A:192:LEU:HD21	2.51	0.44
1:A:181:ASP:O	1:A:182:SER:HB3	2.17	0.44
3:A:506:C8E:H72	3:A:508:C8E:H102	2.00	0.43
3:A:506:C8E:H191	4:A:777:HOH:O	2.18	0.43
1:A:22:LYS:NZ	1:A:377:HIS:HE1	2.16	0.43
1:A:12:LEU:CD1	1:A:12:LEU:N	2.83	0.42
1:A:3:ILE:O	1:A:4:ALA:C	2.63	0.41
1:A:300:LYS:HD2	1:A:341:LEU:HD21	2.03	0.41
1:A:150:TYR:CD2	1:A:150:TYR:C	2.98	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	400/419 (96%)	386 (96%)	13 (3%)	1 (0%)	36 21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	347/363 (96%)	343 (99%)	4 (1%)	63 49

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	123	PRO
1	A	310	GLU
1	A	360	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	211	GLN
1	A	226	ASN
1	A	235	ASN
1	A	259	ASN
1	A	315	ASN
1	A	346	GLN
1	A	377	HIS
1	A	389	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	C8E	A	509	-	11,11,20	0.95	0	10,10,19	1.95	5 (50%)
3	C8E	A	507	-	15,15,20	0.75	0	14,14,19	0.98	1 (7%)
3	C8E	A	510	-	9,9,20	0.64	0	8,8,19	0.79	0
3	C8E	A	508	-	14,14,20	0.91	1 (7%)	13,13,19	1.56	2 (15%)
3	C8E	A	511	-	10,10,20	0.93	0	9,9,19	0.79	0
3	C8E	A	505	-	20,20,20	0.90	0	19,19,19	2.18	6 (31%)
3	C8E	A	506	-	20,20,20	0.74	0	19,19,19	1.33	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C8E	A	509	-	-	1/9/9/18	-
3	C8E	A	507	-	-	5/13/13/18	-
3	C8E	A	510	-	-	4/7/7/18	-
3	C8E	A	508	-	-	4/12/12/18	-
3	C8E	A	511	-	-	0/8/8/18	-
3	C8E	A	505	-	-	8/18/18/18	-
3	C8E	A	506	-	-	4/18/18/18	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	508	C8E	O12-C13	2.02	1.50	1.42

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	505	C8E	C10-O9-C8	5.54	137.51	113.26
3	A	505	C8E	O9-C8-C7	-5.10	83.54	110.32
3	A	508	C8E	C13-O12-C11	-3.41	98.33	113.26
3	A	508	C8E	O12-C13-C14	3.41	125.12	110.11
3	A	509	C8E	C16-O15-C14	3.20	127.26	113.26
3	A	506	C8E	O18-C17-C16	3.13	124.60	110.35
3	A	509	C8E	O18-C17-C16	-3.02	96.58	110.35
3	A	505	C8E	O18-C17-C16	2.94	123.73	110.35
3	A	506	C8E	C13-O12-C11	2.82	125.60	113.26
3	A	505	C8E	O9-C10-C11	-2.64	98.34	110.35
3	A	507	C8E	C10-O9-C8	2.40	123.77	113.26
3	A	509	C8E	O18-C19-C20	-2.40	99.53	110.11
3	A	505	C8E	O12-C11-C10	2.29	120.80	110.35
3	A	509	C8E	O12-C13-C14	2.21	120.44	110.35
3	A	505	C8E	O21-C20-C19	2.07	123.98	111.82
3	A	509	C8E	C11-O12-C13	2.05	120.24	113.06
3	A	506	C8E	O18-C19-C20	2.01	118.99	110.11

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	505	C8E	O12-C13-C14-O15
3	A	505	C8E	O15-C16-C17-O18
3	A	505	C8E	C20-C19-O18-C17
3	A	508	C8E	C2-C3-C4-C5
3	A	508	C8E	C6-C7-C8-O9
3	A	506	C8E	O15-C16-C17-O18
3	A	505	C8E	O18-C19-C20-O21
3	A	508	C8E	O12-C13-C14-O15
3	A	510	C8E	O18-C19-C20-O21
3	A	508	C8E	C4-C5-C6-C7
3	A	507	C8E	C13-C14-O15-C16
3	A	506	C8E	C1-C2-C3-C4
3	A	507	C8E	O9-C10-C11-O12
3	A	507	C8E	C3-C4-C5-C6
3	A	505	C8E	C11-C10-O9-C8

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	505	C8E	C16-C17-O18-C19
3	A	507	C8E	C4-C5-C6-C7
3	A	505	C8E	C10-C11-O12-C13
3	A	510	C8E	C17-C16-O15-C14
3	A	506	C8E	O9-C10-C11-O12
3	A	510	C8E	C16-C17-O18-C19
3	A	506	C8E	O12-C13-C14-O15
3	A	507	C8E	O12-C13-C14-O15
3	A	509	C8E	C10-C11-O12-C13
3	A	510	C8E	O15-C16-C17-O18
3	A	505	C8E	O9-C10-C11-O12


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	508	C8E	2	0
3	A	505	C8E	4	0
3	A	506	C8E	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	402/419 (95%)	0.26	20 (4%) 34 39	18, 30, 58, 82	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ILE	5.8
1	A	4	ALA	5.5
1	A	404	PHE	4.3
1	A	402	PHE	3.8
1	A	26	TYR	3.7
1	A	55	VAL	3.5
1	A	23	LYS	2.8
1	A	368	GLY	2.8
1	A	366	LEU	2.6
1	A	28	TYR	2.5
1	A	365	LYS	2.4
1	A	403	LYS	2.3
1	A	221	LEU	2.3
1	A	27	PRO	2.3
1	A	161	ARG	2.2
1	A	8	ALA	2.2
1	A	6	SER	2.2
1	A	225	LEU	2.1
1	A	162	TYR	2.1
1	A	69	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	C8E	A	509	12/21	0.79	0.23	43,52,64,67	0
3	C8E	A	507	16/21	0.86	0.19	39,53,63,63	0
3	C8E	A	506	21/21	0.87	0.18	32,40,61,73	0
3	C8E	A	505	21/21	0.88	0.18	29,44,56,63	0
3	C8E	A	511	11/21	0.88	0.15	30,42,46,48	0
3	C8E	A	508	15/21	0.89	0.18	40,52,66,69	0
2	CA	A	504	1/1	0.91	0.10	54,54,54,54	0
3	C8E	A	510	10/21	0.92	0.14	49,56,63,64	0
2	CA	A	502	1/1	0.95	0.18	52,52,52,52	0
2	CA	A	503	1/1	0.98	0.05	35,35,35,35	0
2	CA	A	501	1/1	1.00	0.04	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around C8E A 509:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C8E A 507:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C8E A 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C8E A 505:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C8E A 508:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C8E A 510:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.