



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:25 PM UTC

PDB ID : 5DLM / pdb_00005dlm
Title : Complex of Influenza M2e and Antibody
Authors : Cho, K.J.; Schepens, B.; Moonens, K.; Deng, L.; Fiers, W.; Remaut, H.; Saelens, X.
Deposited on : 2015-09-07
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	216	
1	I	216	
2	L	217	
2	M	217	
3	X	23	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Y	23	4% 26% 9% 61%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of monoclonal antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	216	Total	C	N	O	S	0	2	0
			1624	1021	270	324	9			
1	I	214	Total	C	N	O	S	0	0	0
			1597	1006	266	316	9			

- Molecule 2 is a protein called Light chain of monoclonal antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	217	Total	C	N	O	S	0	0	0
			1690	1063	283	339	5			
2	M	217	Total	C	N	O	S	0	1	0
			1698	1068	286	339	5			


- Molecule 3 is a protein called Matrix protein 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	X	9	Total	C	N	O	0	0	0
			68	43	9	16			
3	Y	9	Total	C	N	O	0	0	0
			68	43	9	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	15	GLY	TRP	engineered mutation	UNP A4U6V3
Y	15	GLY	TRP	engineered mutation	UNP A4U6V3

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	O	S	0	0
			5	4	1		
4	M	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	95	Total	O	0	0
			95	95		
5	L	72	Total	O	0	0
			72	72		
5	I	47	Total	O	0	0
			47	47		
5	M	84	Total	O	0	0
			84	84		
5	X	5	Total	O	0	0
			5	5		
5	Y	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heavy chain of monoclonal antibody




- Molecule 1: Heavy chain of monoclonal antibody



- Molecule 2: Light chain of monoclonal antibody




- Molecule 2: Light chain of monoclonal antibody






• Molecule 3: Matrix protein 2



• Molecule 3: Matrix protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.64Å 101.44Å 212.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 2.20 49.21 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.0 (49.21-2.20) 98.0 (49.21-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.187 , 0.240 0.194 , 0.241	Depositor DCC
R_{free} test set	2494 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7060	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	1.60	12/1667 (0.7%)	1.29	11/2276 (0.5%)
1	I	1.37	4/1637 (0.2%)	1.23	10/2235 (0.4%)
2	L	1.49	5/1730 (0.3%)	1.31	6/2347 (0.3%)
2	M	1.53	10/1741 (0.6%)	1.30	8/2361 (0.3%)
3	X	1.34	0/68	1.33	0/93
3	Y	1.18	0/68	1.44	1/93 (1.1%)
All	All	1.49	31/6911 (0.4%)	1.28	36/9405 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1
2	M	0	1
All	All	0	2

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	5	GLN	CA-C	-8.66	1.42	1.52
1	H	209	ASP	C-O	-6.76	1.16	1.23
2	M	49	PRO	N-CA	-6.71	1.38	1.47
2	M	17	GLU	CD-OE1	6.66	1.38	1.25
2	M	12	PRO	C-O	-6.54	1.16	1.23
1	I	110	THR	CB-CG2	-6.49	1.31	1.52
2	M	113	ARG	N-CA	6.40	1.53	1.46
1	H	118	THR	C-O	-6.22	1.16	1.24
1	H	103	SER	N-CA	6.21	1.53	1.46
1	I	167	THR	CA-C	6.11	1.60	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	82	ARG	CA-C	-6.00	1.46	1.53
1	I	42	GLU	CA-C	5.69	1.60	1.52
2	M	60	PHE	CA-C	5.64	1.60	1.52
1	H	121	PRO	C-O	-5.62	1.17	1.23
2	L	151	VAL	C-O	-5.57	1.18	1.24
2	L	90	ILE	N-CA	5.52	1.53	1.46
1	H	30	SER	N-CA	-5.48	1.39	1.46
2	M	66	ARG	CA-C	5.45	1.60	1.52
2	M	150	ASN	CA-C	-5.45	1.46	1.52
2	M	102	THR	CA-C	5.43	1.59	1.52
2	M	6	GLN	N-CA	-5.38	1.39	1.46
2	L	49	PRO	N-CA	-5.36	1.40	1.47
1	I	31	THR	CA-C	5.32	1.59	1.52
1	H	186	PRO	CA-CB	5.15	1.60	1.53
1	H	110	THR	N-CA	5.13	1.52	1.46
2	L	136	SER	N-CA	5.12	1.52	1.46
1	H	123	VAL	C-O	-5.12	1.18	1.24
1	H	119	THR	C-O	-5.10	1.17	1.25
1	H	127	ALA	C-O	-5.08	1.17	1.24
1	H	142	CYS	CA-C	5.01	1.58	1.52
2	M	54	TYR	C-O	-5.01	1.17	1.23

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	81	SER	N-CA-C	12.41	124.89	111.36
2	M	30	VAL	N-CA-C	-10.90	93.34	108.27
2	L	30	VAL	N-CA-C	-10.70	93.61	108.27
2	L	29	ILE	CB-CA-C	-8.34	100.57	110.91
2	M	29	ILE	CB-CA-C	-8.12	100.84	110.91
1	I	135	ASN	N-CA-C	7.48	121.82	112.24
3	Y	9	THR	CB-CA-C	-6.76	101.00	109.31
1	H	175	ASP	N-CA-C	-6.53	103.65	112.26
1	H	29	LEU	CA-C-N	6.30	129.35	120.28
1	H	29	LEU	C-N-CA	6.30	129.35	120.28
1	H	179	LEU	CA-CB-CG	6.13	137.74	116.30
1	H	110	THR	CA-CB-CG2	5.84	120.44	110.50
2	L	142	ASN	N-CA-C	5.81	119.29	109.76
1	H	5	GLN	CB-CA-C	-5.76	100.92	110.14
1	H	78	VAL	CB-CA-C	-5.75	102.07	110.81
1	I	12	VAL	CB-CA-C	-5.74	100.84	111.18
1	I	211	LYS	CG-CD-CE	5.69	124.39	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	113	ARG	CG-CD-NE	5.65	124.43	112.00
2	M	205	THR	N-CA-C	5.57	119.34	112.54
2	L	110	GLU	CB-CG-CD	5.51	121.96	112.60
1	I	174	SER	N-CA-C	5.47	118.88	111.17
1	I	78	VAL	CB-CA-C	-5.47	102.50	110.81
1	H	185	VAL	CA-C-N	-5.43	114.33	119.76
1	H	185	VAL	C-N-CA	-5.43	114.33	119.76
1	I	29	LEU	N-CA-C	5.23	117.66	111.33
1	I	110	THR	CB-CA-C	5.21	118.34	109.80
1	I	29	LEU	CA-C-N	5.20	127.77	120.28
1	I	29	LEU	C-N-CA	5.20	127.77	120.28
2	M	113	ARG	CG-CD-NE	5.16	123.36	112.00
1	I	82	MET	CG-SD-CE	-5.12	89.64	100.90
2	M	63	VAL	CA-C-O	5.11	122.17	119.15
1	H	5	GLN	N-CA-CB	5.05	118.70	110.77
2	M	73	GLY	CA-C-N	5.05	131.46	122.06
2	M	73	GLY	C-N-CA	5.05	131.46	122.06
1	H	82	MET	CG-SD-CE	-5.04	89.80	100.90
2	M	139	CYS	N-CA-CB	5.02	118.49	110.65

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	29	ILE	Peptide
2	M	29	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1624	0	1600	13	0
1	I	1597	0	1574	14	0
2	L	1690	0	1634	13	0
2	M	1698	0	1647	12	0
3	X	68	0	68	1	0
3	Y	68	0	68	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	L	5	0	0	0	0
4	M	5	0	0	0	0
5	H	95	0	0	0	0
5	I	47	0	0	1	0
5	L	72	0	0	1	0
5	M	84	0	0	1	0
5	X	5	0	0	0	0
5	Y	2	0	0	0	0
All	All	7060	0	6591	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:174[B]:SER:OG	1:H:175:ASP:N	2.13	0.82
1:I:6:GLN:HE21	1:I:95:CYS:H	1.25	0.80
1:H:6:GLN:HE21	1:H:95:CYS:H	1.25	0.80
1:I:171:VAL:HG11	2:M:165:LEU:HD11	1.85	0.59
2:M:180:MET:HE3	2:M:181:SER:C	2.28	0.58
1:I:44:ARG:NH1	2:M:105:GLY:O	2.39	0.56
1:H:5:GLN:N	1:H:5:GLN:OE1	2.39	0.56
2:M:194:HIS:O	2:M:216:ARG:NH1	2.40	0.54
3:X:3:LEU:HG	3:X:9:THR:HG23	1.90	0.53
1:I:53:SER:CB	3:Y:7:VAL:HG13	2.38	0.53
2:M:124:PRO:HB3	2:M:214:PHE:CE2	2.45	0.52
2:L:193:ARG:HG2	2:L:193:ARG:HH11	1.74	0.51
1:H:44:ARG:NH1	2:L:105:GLY:O	2.43	0.51
2:L:88:LEU:HD23	5:L:428:HOH:O	2.10	0.51
2:L:113:ARG:NH1	2:L:114:ALA:O	2.41	0.51
1:H:153[A]:THR:CG2	1:H:200:ALA:HB3	2.40	0.51
1:I:82:MET:HE1	1:I:93:TYR:CZ	2.46	0.50
2:L:180:MET:CE	2:L:181:SER:C	2.85	0.49
1:H:67:PHE:CZ	1:H:82:MET:HE2	2.47	0.49
1:I:53:SER:HB2	3:Y:7:VAL:HG13	1.95	0.48
1:H:67:PHE:CZ	1:H:82:MET:CE	2.97	0.48
1:H:153[A]:THR:HG22	1:H:200:ALA:HB3	1.97	0.47
1:H:11:SER:HB2	1:H:149:PRO:HG3	1.97	0.47
1:I:67:PHE:CZ	1:I:82:MET:CE	2.98	0.47
1:H:147:TYR:CE2	1:H:152:VAL:HG13	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:180:MET:HE2	2:M:180:MET:C	2.41	0.46
2:L:180:MET:HE3	2:L:181:SER:O	2.15	0.46
1:H:171:VAL:HG11	2:L:165:LEU:HD11	1.98	0.45
1:H:82:MET:HE1	1:H:93:TYR:CZ	2.51	0.45
2:M:30:VAL:HB	5:M:455:HOH:O	2.16	0.45
2:M:88:LEU:C	2:M:88:LEU:HD12	2.41	0.45
2:L:59:ARG:NE	2:L:65:ASP:HA	2.32	0.45
2:M:14:SER:HB2	2:M:17:GLU:HG2	1.99	0.45
1:I:210:LYS:NZ	5:I:304:HOH:O	2.45	0.45
2:L:29:ILE:HG23	2:L:97:SER:CB	2.47	0.45
2:L:193:ARG:HG2	2:L:193:ARG:NH1	2.32	0.45
2:M:29:ILE:HG23	2:M:97:SER:CB	2.48	0.44
2:L:29:ILE:O	2:L:30:VAL:HG12	2.17	0.44
1:I:191:PRO:O	1:I:192:SER:C	2.61	0.43
2:L:29:ILE:O	2:L:30:VAL:CG1	2.67	0.43
1:I:12:VAL:CG2	1:I:18:LEU:HD22	2.49	0.43
1:I:67:PHE:CE2	1:I:82:MET:HE3	2.54	0.43
1:I:67:PHE:CZ	1:I:82:MET:HE2	2.53	0.43
1:I:147:TYR:CE2	1:I:152:VAL:HG13	2.54	0.43
1:I:11:SER:HB2	1:I:149:PRO:HG3	2.01	0.42
1:H:67:PHE:CE2	1:H:82:MET:HE3	2.55	0.42
2:M:42:LEU:HD13	2:M:91:TYR:CE1	2.54	0.42
2:M:113:ARG:NH1	2:M:114:ALA:O	2.44	0.41
3:Y:7:VAL:HG12	3:Y:8:GLU:HG2	2.01	0.41
2:L:180:MET:CE	2:L:181:SER:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	216/216 (100%)	208 (96%)	5 (2%)	3 (1%)	9 7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	212/216 (98%)	206 (97%)	6 (3%)	0	100	100
2	L	215/217 (99%)	206 (96%)	9 (4%)	0	100	100
2	M	216/217 (100%)	211 (98%)	4 (2%)	1 (0%)	24	27
3	X	7/23 (30%)	7 (100%)	0	0	100	100
3	Y	7/23 (30%)	7 (100%)	0	0	100	100
All	All	873/912 (96%)	845 (97%)	24 (3%)	4 (0%)	36	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	174[A]	SER
1	H	174[B]	SER
1	H	135	ASN
2	M	204	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	188/186 (101%)	179 (95%)	9 (5%)	23	30
1	I	184/186 (99%)	176 (96%)	8 (4%)	26	35
2	L	194/194 (100%)	184 (95%)	10 (5%)	21	26
2	M	195/194 (100%)	185 (95%)	10 (5%)	21	27
3	X	9/21 (43%)	9 (100%)	0	100	100
3	Y	9/21 (43%)	7 (78%)	2 (22%)	1	1
All	All	779/802 (97%)	740 (95%)	39 (5%)	22	28

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	31	THR
1	H	44	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	48	VAL
1	H	115	SER
1	H	134	THR
1	H	136	SER
1	H	161	LEU
1	H	179	LEU
1	H	216	ASP
2	L	22	SER
2	L	30	VAL
2	L	38	LEU
2	L	82	ARG
2	L	113	ARG
2	L	174	LYS
2	L	180	MET
2	L	186	LEU
2	L	208	SER
2	L	213	SER
1	I	12	VAL
1	I	13	LYS
1	I	20	LEU
1	I	44	ARG
1	I	48	VAL
1	I	74	VAL
1	I	192	SER
1	I	205	SER
2	M	17	GLU
2	M	38	LEU
2	M	82	ARG
2	M	113	ARG
2	M	160	ARG
2	M	173	SER
2	M	186	LEU
2	M	196	SER
2	M	208	SER
2	M	217	ASN
3	Y	7	VAL
3	Y	9	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	166	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	143	ASN
1	I	3	GLN
1	I	107	GLN
2	M	47	GLN
2	M	195	ASN
2	M	217	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	L	301	-	4,4,4	0.49	0	6,6,6	0.47	0
4	SO4	M	301	-	4,4,4	0.43	0	6,6,6	0.62	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	216/216 (100%)	-0.27	5 (2%) 61 58	23, 38, 60, 123	2 (0%)
1	I	214/216 (99%)	0.12	5 (2%) 61 58	30, 52, 76, 122	0
2	L	217/217 (100%)	-0.19	4 (1%) 67 64	28, 40, 63, 95	0
2	M	217/217 (100%)	-0.05	6 (2%) 55 52	24, 42, 79, 107	1 (0%)
3	X	9/23 (39%)	-0.06	0 100 100	33, 43, 63, 72	0
3	Y	9/23 (39%)	0.39	1 (11%) 10 8	38, 46, 71, 77	0
All	All	882/912 (96%)	-0.09	21 (2%) 59 56	23, 43, 74, 123	3 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	30	VAL	4.5
1	H	134	THR	4.0
2	M	31	HIS	3.8
2	L	30	VAL	3.5
2	L	29	ILE	3.1
3	Y	10	PRO	3.1
1	I	5	GLN	2.9
1	I	2	VAL	2.9
1	I	134	THR	2.7
2	M	131	THR	2.7
2	M	29	ILE	2.7
1	H	131	ALA	2.6
1	H	132	ALA	2.6
1	H	174[A]	SER	2.5
1	H	5	GLN	2.5
2	L	31	HIS	2.4
2	M	217	ASN	2.3
1	I	132	ALA	2.3
1	I	131	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	M	32	SER	2.2
2	L	72	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	L	301	5/5	0.89	0.09	69,72,79,84	0
4	SO4	M	301	5/5	0.92	0.08	55,63,73,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.