



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:56 PM EDT

PDB ID : 7DMM / pdb_00007dmm
Title : Structure of a glucose isomerase crystal grown in an aqueous glycerol solution without any precipitants
Authors : Suzuki, Y.; Maita, N.
Deposited on : 2020-12-04
Resolution : 0.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1358 (1.04-0.96)
Clashscore	190562	1411 (1.04-0.96)
Ramachandran outliers	187476	1347 (1.04-0.96)
Sidechain outliers	187428	1348 (1.04-0.96)
RSRZ outliers	180081	1355 (1.04-0.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	9% 85% 14%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	17	0
			3142	1983	558	592	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	GLN	ARG	variant	UNP P24300

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Ca 1	0	0

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total 2	Mn 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	303	Total 303	O 303	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.61Å 98.46Å 102.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.16 – 0.99 51.16 – 0.99	Depositor EDS
% Data completeness (in resolution range)	99.7 (51.16-0.99) 99.7 (51.16-0.99)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.08 (at 0.99Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.157 , 0.166 0.167 , 0.174	Depositor DCC
R_{free} test set	13077 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	8.2	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 27.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.010 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3460	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.69	58/3261 (1.8%)	1.28	20/4413 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (58) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	253[A]	LYS	CA-C	-9.44	1.41	1.52
1	A	253[B]	LYS	CA-C	-9.44	1.41	1.52
1	A	289[A]	LYS	CA-C	-8.98	1.45	1.52
1	A	289[B]	LYS	CA-C	-8.98	1.45	1.52
1	A	250	ASN	C-O	8.95	1.38	1.23
1	A	368	ARG	CZ-NH1	8.92	1.45	1.32
1	A	67	GLU	C-N	-8.55	1.23	1.33
1	A	113	ARG	CZ-NH1	-8.41	1.21	1.32
1	A	266	ARG	CZ-NH2	-7.95	1.23	1.33
1	A	56	ASP	CB-CG	7.82	1.71	1.52
1	A	253[A]	LYS	N-CA	-7.28	1.35	1.46
1	A	253[B]	LYS	N-CA	-7.28	1.35	1.46
1	A	347	LEU	C-O	-6.84	1.16	1.24
1	A	68	ARG	CZ-NH2	-6.80	1.24	1.33
1	A	349	ALA	C-O	-6.68	1.16	1.24
1	A	70	GLU	N-CA	-6.63	1.38	1.46
1	A	76	ARG	NE-CZ	-6.60	1.25	1.33
1	A	340[A]	ARG	CZ-NH1	6.59	1.42	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	340[B]	ARG	CZ-NH1	6.59	1.42	1.32
1	A	175	ASP	C-N	-6.59	1.26	1.33
1	A	351	LEU	C-O	-6.48	1.16	1.24
1	A	343	ALA	CA-CB	-6.44	1.48	1.53
1	A	252	ILE	N-CA	-6.43	1.38	1.46
1	A	266	ARG	NE-CZ	6.38	1.40	1.33
1	A	361	ASP	C-N	-6.29	1.25	1.33
1	A	26	PHE	CB-CG	6.28	1.65	1.50
1	A	157	ARG	NE-CZ	-6.17	1.26	1.33
1	A	266	ARG	CD-NE	-6.09	1.37	1.46
1	A	167	GLU	CA-C	-6.06	1.44	1.52
1	A	140	ARG	CZ-NH2	-6.05	1.25	1.33
1	A	29	ALA	CA-C	-6.03	1.45	1.52
1	A	250	ASN	CG-ND2	-6.02	1.20	1.33
1	A	65	ASP	CA-C	-6.00	1.45	1.52
1	A	130	GLY	N-CA	-5.98	1.36	1.45
1	A	6	THR	N-CA	5.79	1.50	1.46
1	A	353	ASP	C-O	5.73	1.31	1.23
1	A	6	THR	C-O	5.69	1.28	1.24
1	A	354	ARG	N-CA	-5.65	1.39	1.46
1	A	362	VAL	N-CA	5.65	1.53	1.46
1	A	348	GLN	C-O	-5.63	1.17	1.24
1	A	172	GLN	C-N	-5.60	1.26	1.33
1	A	76	ARG	CZ-NH1	-5.54	1.25	1.32
1	A	368	ARG	C-N	-5.53	1.26	1.33
1	A	31	ARG	C-O	-5.47	1.17	1.23
1	A	345	ASP	N-CA	5.38	1.53	1.46
1	A	63	SER	C-O	-5.32	1.17	1.23
1	A	65	ASP	C-O	5.32	1.30	1.24
1	A	88	MET	C-O	-5.22	1.17	1.23
1	A	341	PRO	CA-C	-5.18	1.46	1.52
1	A	169	VAL	CB-CG2	-5.17	1.35	1.52
1	A	69	GLU	CA-C	-5.13	1.46	1.52
1	A	220	HIS	CE1-NE2	-5.09	1.27	1.32
1	A	252	ILE	C-O	5.09	1.30	1.23
1	A	167	GLU	C-O	5.07	1.30	1.24
1	A	109	ARG	CZ-NH1	-5.05	1.25	1.32
1	A	26	PHE	CG-CD2	-5.03	1.28	1.38
1	A	365	ALA	C-N	-5.03	1.27	1.34
1	A	177	ARG	CZ-NH2	-5.02	1.26	1.33

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	374	ARG	NE-CZ-NH2	9.07	127.36	119.20
1	A	368	ARG	NE-CZ-NH2	-8.46	111.59	119.20
1	A	55	ASP	CA-CB-CG	7.88	120.48	112.60
1	A	266	ARG	CD-NE-CZ	7.61	135.05	124.40
1	A	382	HIS	CB-CG-CD2	-7.28	121.74	131.20
1	A	110	ASP	CA-CB-CG	6.68	119.28	112.60
1	A	374	ARG	NE-CZ-NH1	-6.67	114.83	121.50
1	A	340[A]	ARG	NE-CZ-NH1	6.39	127.89	121.50
1	A	340[B]	ARG	NE-CZ-NH1	6.39	127.89	121.50
1	A	345	ASP	CA-CB-CG	6.38	118.98	112.60
1	A	285	HIS	CB-CG-ND1	6.20	132.00	122.70
1	A	177	ARG	NE-CZ-NH1	-6.19	115.31	121.50
1	A	285	HIS	CB-CG-CD2	-5.86	123.58	131.20
1	A	252	ILE	CB-CA-C	5.80	119.75	111.81
1	A	113	ARG	NE-CZ-NH1	5.65	127.15	121.50
1	A	368	ARG	CA-CB-CG	5.46	125.03	114.10
1	A	250	ASN	CB-CG-ND2	5.33	124.40	116.40
1	A	362	VAL	N-CA-CB	5.33	117.80	110.54
1	A	220	HIS	ND1-CE1-NE2	5.06	113.46	108.40
1	A	57	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3142	0	3043	7	0
2	A	12	0	14	0	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	A	303	0	0	2	2
All	All	3460	0	3057	7	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:323:ASP:OD2	1:A:382:HIS:HE1	1.84	0.61
1:A:152[A]:ARG:HG3	5:A:665:HOH:O	2.06	0.54
1:A:132:GLU:HG3	5:A:612:HOH:O	2.15	0.46
1:A:52:THR:OG1	1:A:285:HIS:HE1	1.99	0.44
1:A:337:GLU:HG2	1:A:340[A]:ARG:NH2	2.35	0.42
1:A:245:ASP:OD1	1:A:285:HIS:HD2	2.03	0.41
1:A:25[B]:PRO:HB2	1:A:26:PHE:CE2	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:778:HOH:O	5:A:797:HOH:O[4_554]	1.93	0.27
5:A:622:HOH:O	5:A:797:HOH:O[4_554]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	401/386 (104%)	386 (96%)	14 (4%)	1 (0%)	43 14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/303 (106%)	319 (100%)	1 (0%)	86	61

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	285	HIS
1	A	382	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	402	-	5,5,5	2.61	2 (40%)	5,5,5	1.30	1 (20%)
2	GOL	A	401	4	5,5,5	0.67	0	5,5,5	1.02	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	402	-	-	0/4/4/4	-
2	GOL	A	401	4	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	GOL	C1-C2	5.04	1.71	1.51
2	A	402	GOL	C3-C2	-2.91	1.40	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	GOL	O2-C2-C1	-2.08	100.57	109.18
2	A	401	GOL	C3-C2-C1	2.06	119.37	111.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	386/386 (100%)	0.52	33 (8%)	16 17	4, 8, 17, 26	17 (4%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	GLY	4.3
1	A	175	ASP	3.6
1	A	2	ASN	3.4
1	A	69	GLU	3.4
1	A	8	GLU	3.2
1	A	66	SER	3.2
1	A	296	PHE	3.1
1	A	26	PHE	2.9
1	A	4	GLN	2.8
1	A	6	THR	2.8
1	A	344	ALA	2.7
1	A	64	SER	2.7
1	A	349	ALA	2.6
1	A	7	PRO	2.5
1	A	62	GLY	2.4
1	A	173	GLY	2.4
1	A	252	ILE	2.4
1	A	281	SER	2.4
1	A	170	THR	2.3
1	A	10	ARG	2.3
1	A	359	GLU	2.3
1	A	354	ARG	2.3
1	A	351	LEU	2.3
1	A	357	PHE	2.2
1	A	65	ASP	2.2
1	A	343	ALA	2.2
1	A	34	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	368	ARG	2.1
1	A	364	ALA	2.1
1	A	208	ARG	2.1
1	A	32	ARG	2.0
1	A	250	ASN	2.0
1	A	350	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	402	6/6	0.91	0.12	8,14,18,22	0
3	CA	A	403	1/1	0.97	0.47	23,23,23,23	0
2	GOL	A	401	6/6	0.98	0.05	6,7,7,7	0
4	MN	A	404	1/1	0.99	0.07	7,7,7,7	0
4	MN	A	405	1/1	1.00	0.17	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.