



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:09 AM UTC

PDB ID : 8DMQ / pdb_00008dmq
Title : Crystal structure of Legionella pneumophila macrodomain MavL in complex with ubiquitin vinyl methyl ester
Authors : Zhang, Z.; Das, C.
Deposited on : 2022-07-08
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	3% 84% 8% 8%
1	B	399	2% 82% 11% 7%
2	C	75	5% 95% 5%
2	D	75	29% 81% 19%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7348 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MavL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	2	1	0
			2844	1804	476	554	10			
1	B	370	Total	C	N	O	S	6	0	0
			2859	1815	480	555	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	GLY	-	expression tag	UNP Q5ZSJ1
A	38	PRO	-	expression tag	UNP Q5ZSJ1
A	39	LEU	-	expression tag	UNP Q5ZSJ1
A	40	GLY	-	expression tag	UNP Q5ZSJ1
A	41	SER	-	expression tag	UNP Q5ZSJ1
B	37	GLY	-	expression tag	UNP Q5ZSJ1
B	38	PRO	-	expression tag	UNP Q5ZSJ1
B	39	LEU	-	expression tag	UNP Q5ZSJ1
B	40	GLY	-	expression tag	UNP Q5ZSJ1
B	41	SER	-	expression tag	UNP Q5ZSJ1

- Molecule 2 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	75	Total	C	N	O	S	4	0	0
			585	367	101	116	1			
2	D	75	Total	C	N	O	S	0	0	0
			559	356	94	108	1			

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is METHYL 4-AMINOBUTANOATE (CCD ID: GVE) (formula: $C_5H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			8	5	1	2		
5	D	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 6 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			16	10	6		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	156	Total	O	0	0
			156	156		
7	B	209	Total	O	0	0
			209	209		
7	C	21	Total	O	0	0
			21	21		
7	D	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MavL




- Molecule 1: MavL



- Molecule 2: Ubiquitin



- Molecule 2: Ubiquitin



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.72Å 108.72Å 209.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	69.71 – 2.19 69.71 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.7 (69.71-2.19) 99.7 (69.71-2.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.178 , 0.209 0.190 , 0.219	Depositor DCC
R_{free} test set	3615 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7348	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 1PE, GVE, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2914	0.60	0/3961
1	B	0.44	0/2929	0.64	1/3979 (0.0%)
2	C	0.39	0/591	0.60	0/796
2	D	0.30	0/565	0.56	0/766
All	All	0.40	0/6999	0.61	1/9502 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	ASP	CA-CB-CG	5.52	118.12	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2844	0	2725	22	0
1	B	2859	0	2753	31	0
2	C	585	0	597	4	0
2	D	559	0	562	13	0
3	A	39	0	15	1	0
3	B	13	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	13	0	5	3	0
4	B	4	0	6	0	0
4	C	8	0	12	0	0
5	C	8	0	0	0	0
5	D	8	0	0	0	0
6	D	16	0	22	2	0
7	A	156	0	0	2	0
7	B	209	0	0	1	0
7	C	21	0	0	0	0
7	D	6	0	0	0	0
All	All	7348	0	6702	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:ALA:HB1	1:B:343:MET:HE1	1.37	1.05
1:B:324:TYR:HD2	1:B:343:MET:HE3	1.39	0.88
1:A:330:GLN:HE21	2:D:33:LYS:HA	1.40	0.86
2:D:5:VAL:HB	2:D:13:ILE:HG23	1.59	0.83
1:B:337:ALA:CB	1:B:343:MET:HE1	2.20	0.67
1:B:312:VAL:HG22	1:B:336:LYS:HD3	1.76	0.67
1:A:126:GLY:O	1:A:161:PRO:HG2	1.95	0.66
1:B:324:TYR:CD2	1:B:343:MET:HE3	2.28	0.66
2:D:1:MET:N	2:D:17:VAL:O	2.29	0.64
2:D:16:GLU:O	2:D:29:LYS:NZ	2.31	0.63
1:B:337:ALA:HB1	1:B:343:MET:CE	2.21	0.62
1:A:83:SER:OG	1:A:86:GLU:HG2	2.00	0.61
1:B:240:ARG:HD3	1:B:268:ASP:OD2	2.00	0.61
1:B:209:ARG:HG3	1:B:256:TYR:CZ	2.36	0.61
1:A:52:LEU:HD13	1:A:403:LEU:HG	1.86	0.58
1:A:265:TYR:CD2	1:A:266:MET:HG2	2.38	0.57
1:B:314:TRP:CD1	1:B:315:ASP:H	2.25	0.54
1:B:59:LYS:O	1:B:63:GLU:HG3	2.06	0.54
1:B:104:VAL:HG13	2:C:47:GLY:HA3	1.89	0.54
1:B:227:PHE:CZ	1:B:315:ASP:HB3	2.43	0.53
1:A:82:LYS:HB3	1:A:86:GLU:HG3	1.90	0.53
2:D:55:THR:HG22	2:D:58:ASP:CG	2.35	0.52
2:D:55:THR:HG23	2:D:57:SER:H	1.74	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:TRP:CG	1:A:315:ASP:H	2.29	0.51
1:B:227:PHE:HZ	1:B:315:ASP:HB3	1.76	0.50
2:D:14:THR:O	2:D:33:LYS:HE3	2.12	0.49
1:B:101:LYS:O	1:B:318:SER:HA	2.13	0.48
1:B:227:PHE:HZ	1:B:315:ASP:CB	2.25	0.48
1:A:45:LEU:HD13	1:A:97:ILE:HD13	1.96	0.48
3:A:501:FLC:OG2	3:A:501:FLC:OHB	2.30	0.48
1:A:149:HIS:CE1	1:A:153:ILE:HD12	2.49	0.47
2:D:42:ARG:HE	6:D:601:1PE:H241	1.80	0.47
1:B:370:ARG:HB3	1:B:374:SER:HB3	1.97	0.47
1:A:55:ILE:HG22	1:A:89:ILE:HD11	1.96	0.47
1:A:240:ARG:HD3	1:A:268:ASP:OD2	2.16	0.46
2:D:55:THR:HG23	2:D:57:SER:N	2.31	0.46
1:A:127:ASP:HA	1:A:196:ARG:HG2	1.98	0.46
3:C:104:FLC:HA2	3:C:104:FLC:OG1	2.16	0.45
1:B:273:LYS:HB3	1:B:273:LYS:HE3	1.54	0.45
1:B:302:ASN:HB2	1:B:303:PRO:HD2	1.97	0.45
1:A:68:THR:HG23	7:A:679:HOH:O	2.16	0.45
1:A:330:GLN:HE22	2:D:33:LYS:HG2	1.81	0.45
1:A:254:LEU:HB3	1:A:257:ILE:HD12	1.98	0.44
2:C:42:ARG:HE	3:C:104:FLC:HA1	1.82	0.44
1:A:45:LEU:HD23	1:A:130:VAL:HB	2.00	0.44
1:A:265:TYR:HD2	1:A:266:MET:HG2	1.78	0.44
1:B:253:SER:C	1:B:255:PRO:HD3	2.42	0.44
1:B:327:GLY:C	2:D:72:ARG:HD3	2.43	0.43
1:A:281:ARG:HD3	7:A:613:HOH:O	2.17	0.43
2:C:42:ARG:NE	3:C:104:FLC:HA1	2.33	0.43
1:A:353:TYR:OH	1:A:358:GLY:HA2	2.19	0.43
1:B:265:TYR:CD2	1:B:266:MET:HG2	2.53	0.43
1:B:254:LEU:N	1:B:255:PRO:HD3	2.35	0.42
1:A:302:ASN:HB2	1:A:303:PRO:HD2	2.02	0.42
1:B:107:GLU:H	1:B:107:GLU:CD	2.28	0.42
1:B:183:LYS:HE2	7:B:923:HOH:O	2.20	0.42
1:B:103:GLN:O	2:C:48:LYS:HD2	2.20	0.41
2:D:42:ARG:NH2	6:D:601:1PE:H231	2.35	0.41
1:A:101:LYS:O	1:A:318:SER:HA	2.19	0.41
1:B:295:LEU:HD21	1:B:336:LYS:HG2	2.02	0.41
1:B:395:ASP:HB3	1:B:406:VAL:HG22	2.01	0.41
1:B:362:PRO:HA	1:B:376:TRP:CZ2	2.56	0.41
1:B:49:LYS:HB2	1:B:394:LEU:O	2.21	0.40
1:A:359:ARG:NH2	2:D:31:GLN:HG2	2.36	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:368:LYS:O	1:B:369:ASP:HB2	2.22	0.40
1:B:353:TYR:OH	1:B:358:GLY:HA2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/399 (92%)	350 (95%)	17 (5%)	0	100	100
1	B	368/399 (92%)	352 (96%)	15 (4%)	1 (0%)	36	42
2	C	73/75 (97%)	70 (96%)	2 (3%)	1 (1%)	9	7
2	D	73/75 (97%)	71 (97%)	2 (3%)	0	100	100
All	All	881/948 (93%)	843 (96%)	36 (4%)	2 (0%)	43	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	322	ASN
2	C	74	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/341 (89%)	302 (100%)	1 (0%)	86	93
1	B	305/341 (89%)	305 (100%)	0	100	100
2	C	65/68 (96%)	65 (100%)	0	100	100
2	D	59/68 (87%)	59 (100%)	0	100	100
All	All	732/818 (90%)	731 (100%)	1 (0%)	88	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	103	GLN
1	A	330	GLN
1	B	109	GLN
1	B	210	GLN
2	C	40	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FLC	C	104	-	12,12,12	1.10	0	17,17,17	1.77	5 (29%)
4	EDO	C	103	-	3,3,3	0.45	0	2,2,2	0.26	0
6	1PE	D	601	-	15,15,15	0.55	0	14,14,14	0.39	0
3	FLC	A	502	-	12,12,12	1.31	0	17,17,17	2.15	5 (29%)
3	FLC	A	503	-	12,12,12	1.08	0	17,17,17	1.21	1 (5%)
5	GVE	C	101	1,2	7,7,7	1.36	2 (28%)	7,7,7	1.23	0
5	GVE	D	602	1,2	7,7,7	1.41	1 (14%)	7,7,7	1.04	1 (14%)
4	EDO	B	702	-	3,3,3	0.42	0	2,2,2	0.29	0
4	EDO	C	102	-	3,3,3	0.42	0	2,2,2	0.47	0
3	FLC	B	701	-	12,12,12	1.22	0	17,17,17	1.65	3 (17%)
3	FLC	A	501	-	12,12,12	1.07	0	17,17,17	1.32	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	C	104	-	-	9/16/16/16	-
4	EDO	C	103	-	-	0/1/1/1	-
6	1PE	D	601	-	-	6/13/13/13	-
3	FLC	A	502	-	-	4/16/16/16	-
3	FLC	A	503	-	-	6/16/16/16	-
5	GVE	C	101	1,2	-	0/6/6/6	-
5	GVE	D	602	1,2	-	3/6/6/6	-
4	EDO	B	702	-	-	1/1/1/1	-
4	EDO	C	102	-	-	1/1/1/1	-
3	FLC	B	701	-	-	8/16/16/16	-
3	FLC	A	501	-	-	10/16/16/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	602	GVE	OXT-C	2.76	1.41	1.33
5	C	101	GVE	OXT-C	2.66	1.41	1.33
5	C	101	GVE	OXT-CH3	-2.03	1.40	1.45

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	FLC	OB2-CBC-CB	4.82	122.39	113.14
3	A	502	FLC	OHB-CB-CBC	-4.14	103.09	108.96
3	C	104	FLC	OB2-CBC-CB	3.96	120.74	113.14
3	A	502	FLC	CG-CB-CBC	3.41	117.57	110.03
3	A	501	FLC	OB2-CBC-CB	3.35	119.56	113.14
3	B	701	FLC	OB2-CBC-CB	3.34	119.54	113.14
3	A	503	FLC	OB2-CBC-CB	3.03	118.95	113.14
3	B	701	FLC	OA2-CAC-CA	2.59	122.56	114.35
3	C	104	FLC	CB-CG-CGC	2.52	120.82	113.92
3	C	104	FLC	OG2-CGC-OG1	-2.39	117.19	123.33
3	C	104	FLC	CG-CB-CA	2.36	115.37	109.31
3	A	502	FLC	OHB-CB-CG	-2.28	104.18	109.38
3	C	104	FLC	OA2-CAC-CA	2.25	121.48	114.35
5	D	602	GVE	C1-CB-CG	-2.06	106.77	112.81
3	B	701	FLC	OHB-CB-CA	-2.05	104.70	109.38
3	A	502	FLC	OB1-CBC-CB	-2.01	118.19	122.09

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	FLC	CBC-CB-CG-CGC
3	A	501	FLC	OHB-CB-CG-CGC
3	A	503	FLC	CA-CB-CG-CGC
3	A	503	FLC	CBC-CB-CG-CGC
3	B	701	FLC	OHB-CB-CBC-OB1
3	B	701	FLC	OHB-CB-CBC-OB2
3	C	104	FLC	CA-CB-CBC-OB1
3	C	104	FLC	CA-CB-CBC-OB2
3	C	104	FLC	OHB-CB-CBC-OB1
3	C	104	FLC	OHB-CB-CBC-OB2
3	C	104	FLC	CA-CB-CG-CGC
3	C	104	FLC	CBC-CB-CG-CGC
3	C	104	FLC	OHB-CB-CG-CGC
6	D	601	1PE	OH5-C14-C24-OH4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	501	FLC	CA-CB-CG-CGC
3	A	502	FLC	CA-CB-CG-CGC
3	A	502	FLC	CBC-CB-CG-CGC
3	A	502	FLC	OHB-CB-CG-CGC
3	A	503	FLC	CAC-CA-CB-CBC
5	D	602	GVE	C1-CB-CG-C
3	A	503	FLC	CAC-CA-CB-CG
3	A	503	FLC	OHB-CB-CG-CGC
3	A	503	FLC	CAC-CA-CB-OHB
3	C	104	FLC	CB-CA-CAC-OA1
6	D	601	1PE	OH2-C12-C22-OH3
3	C	104	FLC	CB-CA-CAC-OA2
4	C	102	EDO	O1-C1-C2-O2
6	D	601	1PE	C15-C25-OH5-C14
6	D	601	1PE	C23-C13-OH4-C24
4	B	702	EDO	O1-C1-C2-O2
3	A	501	FLC	CG-CB-CBC-OB1
3	B	701	FLC	CA-CB-CBC-OB1
3	B	701	FLC	CA-CB-CBC-OB2
3	B	701	FLC	CG-CB-CBC-OB2
5	D	602	GVE	CG-C-OXT-CH3
3	B	701	FLC	CAC-CA-CB-OHB
6	D	601	1PE	C12-C22-OH3-C23
3	A	501	FLC	CA-CB-CBC-OB2
3	A	501	FLC	CG-CB-CBC-OB2
3	A	501	FLC	CA-CB-CBC-OB1
5	D	602	GVE	O-C-OXT-CH3
3	A	501	FLC	CB-CG-CGC-OG1
3	A	501	FLC	OHB-CB-CBC-OB1
3	B	701	FLC	CG-CB-CBC-OB1
3	A	501	FLC	CB-CG-CGC-OG2
3	A	502	FLC	CB-CA-CAC-OA1
6	D	601	1PE	C24-C14-OH5-C25
3	B	701	FLC	CB-CA-CAC-OA1

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	104	FLC	3	0
6	D	601	1PE	2	0
3	A	501	FLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/399 (92%)	0.25	10 (2%) 56 53	32, 45, 60, 80	2 (0%)
1	B	370/399 (92%)	0.08	7 (1%) 66 63	29, 40, 56, 77	2 (0%)
2	C	75/75 (100%)	0.54	4 (5%) 32 29	36, 49, 67, 74	3 (4%)
2	D	75/75 (100%)	1.45	22 (29%) 1 1	49, 68, 85, 94	0
All	All	888/948 (93%)	0.31	43 (4%) 35 32	29, 44, 72, 94	7 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	41	SER	6.9
1	A	409	ARG	5.5
2	C	73	LEU	4.9
2	D	13	ILE	4.3
2	D	14	THR	4.1
1	A	266	MET	3.7
2	D	73	LEU	3.5
2	D	60	ASN	3.5
1	A	330	GLN	3.4
2	D	46	ALA	3.3
2	D	52	ASP	3.1
2	D	63	LYS	2.8
2	D	20	SER	2.8
2	D	47	GLY	2.8
1	A	368	LYS	2.8
1	B	410	SER	2.8
2	D	23	ILE	2.8
1	A	42	ALA	2.7
1	B	42	ALA	2.7
2	D	59	TYR	2.7
2	D	39	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	8	LEU	2.6
2	D	12	THR	2.5
2	C	74	ARG	2.5
1	A	230	ALA	2.4
1	B	357	TRP	2.4
2	D	64	GLU	2.3
2	D	62	GLN	2.3
2	D	45	PHE	2.3
2	D	53	GLY	2.3
1	A	153	ILE	2.3
2	D	65	SER	2.2
1	A	373	MET	2.2
1	A	329	ARG	2.2
1	B	269	GLU	2.2
2	D	1	MET	2.2
2	D	17	VAL	2.2
2	C	10	GLY	2.2
1	B	60	GLN	2.1
1	B	368	LYS	2.1
1	A	371	LYS	2.1
2	D	3	ILE	2.0
2	D	57	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GVE	D	602	8/8	0.60	0.33	91,92,99,101	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GVE	C	101	8/8	0.65	0.26	65,67,73,74	0
4	EDO	C	102	4/4	0.75	0.24	68,71,73,74	0
3	FLC	A	501	13/13	0.76	0.15	91,104,113,115	0
3	FLC	C	104	13/13	0.78	0.18	91,93,96,98	0
3	FLC	B	701	13/13	0.78	0.14	91,93,101,102	0
3	FLC	A	503	13/13	0.83	0.15	90,98,105,107	0
6	1PE	D	601	16/16	0.83	0.20	68,71,78,80	0
3	FLC	A	502	13/13	0.86	0.14	58,62,65,68	0
4	EDO	C	103	4/4	0.88	0.16	59,60,61,61	0
4	EDO	B	702	4/4	0.89	0.17	67,71,71,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.