



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:32 PM UTC

PDB ID : 3DNF / pdb_00003dnf
Title : Structure of (E)-4-Hydroxy-3-methyl-but-2-enyl Diphosphate Reductase, the Terminal Enzyme of the Non-Mevalonate Pathway
Authors : Re kittke, I.
Deposited on : 2008-07-02
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	
1	B	297	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	281	Total	C	N	O	S	0	3	0
			2215	1408	383	415	9			
1	B	282	Total	C	N	O	S	0	12	0
			2274	1455	390	420	9			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	ARG	-	expression tag	UNP O67625
A	291	SER	-	expression tag	UNP O67625
A	292	HIS	-	expression tag	UNP O67625
A	293	HIS	-	expression tag	UNP O67625
A	294	HIS	-	expression tag	UNP O67625
A	295	HIS	-	expression tag	UNP O67625
A	296	HIS	-	expression tag	UNP O67625
A	297	HIS	-	expression tag	UNP O67625
B	290	ARG	-	expression tag	UNP O67625
B	291	SER	-	expression tag	UNP O67625
B	292	HIS	-	expression tag	UNP O67625
B	293	HIS	-	expression tag	UNP O67625
B	294	HIS	-	expression tag	UNP O67625
B	295	HIS	-	expression tag	UNP O67625
B	296	HIS	-	expression tag	UNP O67625
B	297	HIS	-	expression tag	UNP O67625

- Molecule 2 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			7	3	4		
2	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	243	Total 243	O 243	0	0
4	B	318	Total 318	O 318	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase



- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.37Å 87.76Å 72.40Å 90.00° 95.13° 90.00°	Depositor
Resolution (Å)	9.90 – 1.65 9.90 – 1.65	Depositor EDS
% Data completeness (in resolution range)	91.3 (9.90-1.65) 90.9 (9.90-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.226 0.201 , 0.231	Depositor DCC
R_{free} test set	4120 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	22.5	Xtrriage
Anisotropy	0.077	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5076	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	7/2259 (0.3%)	1.01	2/3052 (0.1%)
1	B	0.87	5/2348 (0.2%)	0.91	0/3170
All	All	0.94	12/4607 (0.3%)	0.96	2/6222 (0.0%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	ASN	CB-CG	11.38	1.80	1.52
1	A	237	ASN	CG-OD1	10.79	1.44	1.23
1	A	246	GLU	C-N	9.50	1.47	1.33
1	A	246	GLU	C-O	9.03	1.35	1.24
1	B	211	ASP	CG-OD2	8.35	1.41	1.25
1	B	253	ARG	C-O	7.13	1.32	1.24
1	A	236	PRO	C-O	-6.72	1.15	1.24
1	B	211	ASP	CG-OD1	6.68	1.38	1.25
1	B	253	ARG	C-N	6.04	1.40	1.33
1	A	238	THR	C-O	5.71	1.30	1.24
1	A	251	TRP	C-O	5.59	1.31	1.24
1	B	235	ASN	CG-OD1	5.08	1.33	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	ASN	OD1-CG-ND2	-12.07	110.53	122.60
1	A	102	VAL	N-CA-C	-5.33	105.52	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	0	2261	25	0
1	B	2274	0	2365	40	0
2	A	7	0	0	1	0
2	B	7	0	0	0	0
3	A	6	0	8	2	0
3	B	6	0	8	1	0
4	A	243	0	0	1	0
4	B	318	0	0	9	0
All	All	5076	0	4642	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:ASN:CB	1:A:237:ASN:CG	1.80	1.54
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:CG1	1.72	1.18
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:CG1	2.26	1.13
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:HG12	1.87	1.02
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:HG13	1.47	0.95
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:CG1	2.03	0.88
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:HG12	1.48	0.86
1:B:124:HIS:HD2	1:B:126:GLU:H	1.22	0.86
1:A:124:HIS:HD2	1:A:126:GLU:H	1.32	0.75
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:HG12	1.67	0.71
1:B:104:GLU:OE1	4:B:4136:HOH:O	2.11	0.68
1:B:124:HIS:CD2	1:B:126:GLU:H	2.08	0.68
1:A:247:LEU:HD13	1:A:278:ILE:HG13	1.79	0.64
1:A:124:HIS:CD2	1:A:126:GLU:H	2.14	0.64
1:B:243:THR:HG22	1:B:245:GLU:H	1.63	0.63
1:A:247:LEU:HD11	1:A:274:VAL:HG13	1.81	0.62
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:HG11	1.80	0.62
1:B:211:ASP:HB2	1:B:255:VAL:HG13	1.82	0.61
1:A:40:ILE:HG23	1:A:41:ILE:HD12	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:LYS:NZ	4:B:4008:HOH:O	2.21	0.59
1:B:48:ASN:OD1	4:B:4175:HOH:O	2.15	0.58
1:B:59:GLN:OE1	4:B:4176:HOH:O	2.16	0.58
1:B:223:ASN:HB2	4:B:4165:HOH:O	2.03	0.58
1:A:237:ASN:CG	1:A:237:ASN:CA	2.72	0.57
1:B:277:ARG:O	1:B:277:ARG:HD3	2.05	0.56
1:B:33:LYS:HE3	1:B:56:PHE:CD2	2.41	0.55
1:B:247:LEU:HD11	1:B:274:VAL:HG13	1.89	0.55
1:B:104:GLU:HG2	4:B:4101:HOH:O	2.06	0.54
1:B:204:LYS:HG3	1:B:230:ILE:HD11	1.89	0.54
1:A:248:GLN:HB2	1:A:251:TRP:CE2	2.43	0.53
1:B:196:THR:HG23	4:B:3891:HOH:O	2.08	0.53
1:B:156:HIS:HD2	3:B:3732:GOL:O2	1.92	0.53
1:A:4:ILE:O	1:A:4:ILE:HG13	2.10	0.51
1:B:217:GLY:HA3	1:B:224:THR:HG21	1.93	0.50
1:B:203:VAL:HG21	1:B:227:LEU:HD22	1.94	0.50
1:B:204:LYS:CG	1:B:230:ILE:HD11	2.42	0.50
1:B:156:HIS:HE1	4:B:3908:HOH:O	1.95	0.48
1:B:4:ILE:H	1:B:4:ILE:HD12	1.78	0.47
1:A:40:ILE:O	1:A:41:ILE:HG23	2.15	0.47
1:A:156:HIS:HD2	3:A:3731:GOL:O2	1.97	0.47
1:A:142:ILE:HD13	1:A:152:GLU:HB3	1.95	0.47
1:A:187:LYS:HE3	1:B:185[A]:GLU:CD	2.40	0.47
1:A:28:LYS:HG2	1:A:29:GLU:HG3	1.96	0.47
1:A:256:LYS:HE3	1:A:257:ARG:HG3	1.97	0.47
1:A:43:ASN:HD22	1:A:43:ASN:C	2.22	0.46
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:HG11	2.31	0.46
1:A:72:ARG:NH2	1:A:74:HIS:HE1	2.14	0.46
1:B:267:PRO:HD2	1:B:270:ILE:HD12	1.98	0.46
1:A:74:HIS:HD2	1:A:75:GLY:O	1.98	0.45
1:A:137[B]:CYS:SG	4:A:3800:HOH:O	2.61	0.45
1:A:217:GLY:HA3	1:A:224:THR:HG21	2.00	0.44
1:B:210:VAL:HA	1:B:257:ARG:HE	1.83	0.43
1:A:3:ASP:HB3	1:A:5:ILE:HG12	1.99	0.43
1:A:15:GLY:HA3	2:A:2302:F3S:S4	2.58	0.43
1:A:248:GLN:HE21	1:A:251:TRP:HZ2	1.65	0.43
1:B:64:LYS:O	1:B:65:GLU:C	2.61	0.42
1:B:6:ILE:HD11	1:B:275:LYS:HD2	2.02	0.42
1:A:187:LYS:HE3	1:B:185[A]:GLU:OE2	2.20	0.42
1:B:74:HIS:HE1	4:B:4021:HOH:O	2.02	0.41
1:B:124:HIS:HD2	1:B:126:GLU:N	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLY:O	3:A:3731:GOL:H31	2.21	0.41
1:B:107:CYS:SG	1:B:137[A]:CYS:SG	3.14	0.40
1:B:227:LEU:O	1:B:230:ILE:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	282/297 (95%)	278 (99%)	3 (1%)	1 (0%)	30 15
1	B	292/297 (98%)	287 (98%)	5 (2%)	0	100 100
All	All	574/594 (97%)	565 (98%)	8 (1%)	1 (0%)	43 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	243/255 (95%)	232 (96%)	11 (4%)	24 5
1	B	253/255 (99%)	247 (98%)	6 (2%)	43 20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	496/510 (97%)	479 (97%)	17 (3%)	33 10

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	VAL
1	A	4	ILE
1	A	5	ILE
1	A	41	ILE
1	A	43	ASN
1	A	197[A]	SER
1	A	197[B]	SER
1	A	198	LEU
1	A	233	GLU
1	A	234	LEU
1	B	1	MET
1	B	4	ILE
1	B	92	ILE
1	B	212	VAL
1	B	277	ARG
1	B	282	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	74	HIS
1	A	123	ASN
1	A	124	HIS
1	A	156	HIS
1	A	235	ASN
1	B	9	HIS
1	B	59	GLN
1	B	124	HIS
1	B	156	HIS
1	B	223	ASN
1	B	273	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	3732	-	5,5,5	0.25	0	5,5,5	0.72	0
2	F3S	B	3302	1	0,9,9	-	-	-		
2	F3S	A	2302	1	0,9,9	-	-	-		
3	GOL	A	3731	-	5,5,5	0.30	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	3732	-	-	4/4/4/4	-
2	F3S	A	2302	1	-	-	0/3/3/3
2	F3S	B	3302	1	-	-	0/3/3/3
3	GOL	A	3731	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	3732	GOL	O1-C1-C2-C3
3	B	3732	GOL	C1-C2-C3-O3
3	B	3732	GOL	O1-C1-C2-O2
3	B	3732	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	3732	GOL	1	0
2	A	2302	F3S	1	0
3	A	3731	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/297 (94%)	0.83	36 (12%) 7 8	16, 33, 41, 75	3 (1%)
1	B	282/297 (94%)	0.95	56 (19%) 3 3	18, 33, 40, 71	12 (4%)
All	All	563/594 (94%)	0.89	92 (16%) 4 5	16, 33, 40, 75	15 (2%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	13	CYS	9.4
1	A	2	VAL	9.1
1	B	2	VAL	6.0
1	A	1	MET	5.7
1	A	13	CYS	5.7
1	B	264	ALA	5.7
1	B	197	SER	5.6
1	B	3	ASP	5.2
1	A	4	ILE	4.8
1	B	1	MET	4.8
1	B	236	PRO	4.7
1	B	249	PRO	4.7
1	B	222	GLY	4.6
1	B	11	GLY	4.4
1	B	12	PHE	4.3
1	A	3	ASP	4.3
1	B	255	VAL	4.2
1	A	42	HIS	4.1
1	A	251	TRP	3.9
1	A	244	ALA	3.8
1	B	251	TRP	3.7
1	B	252	PHE	3.6
1	B	248	GLN	3.6
1	B	196	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	10	ALA	3.5
1	B	282	CYS	3.5
1	B	254	GLY	3.5
1	B	281	ILE	3.4
1	A	5	ILE	3.4
1	A	247	LEU	3.4
1	B	280	GLU	3.4
1	B	262	ALA	3.4
1	A	207	ALA	3.3
1	B	279	GLN	3.3
1	B	265	SER	3.3
1	A	198	LEU	3.2
1	B	253	ARG	3.2
1	B	4	ILE	3.1
1	A	196	THR	3.1
1	B	202	SER	3.1
1	B	239	TYR	3.1
1	B	245	GLU	3.0
1	B	263	GLY	3.0
1	A	248	GLN	2.9
1	A	237	ASN	2.9
1	B	199	ARG	2.8
1	A	279	GLN	2.8
1	B	226	ARG	2.8
1	A	221	SER	2.8
1	B	198	LEU	2.7
1	A	6	ILE	2.7
1	B	231	SER	2.7
1	B	229	TYR	2.7
1	B	230	ILE	2.7
1	A	222	GLY	2.6
1	B	250	GLU	2.6
1	B	266	THR	2.6
1	B	9	HIS	2.5
1	A	256	LYS	2.5
1	B	258	VAL	2.5
1	A	41	ILE	2.5
1	A	280	GLU	2.4
1	A	197[A]	SER	2.4
1	B	123[A]	ASN	2.4
1	B	5	ILE	2.4
1	A	29	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	264	ALA	2.4
1	B	224	THR	2.3
1	A	12	PHE	2.3
1	B	42	HIS	2.3
1	A	255	VAL	2.3
1	B	247	LEU	2.3
1	B	98	TYR	2.3
1	A	7	ALA	2.3
1	B	276	SER	2.2
1	A	101	ALA	2.2
1	B	269	TRP	2.2
1	B	200	GLN	2.2
1	A	194	ASN	2.2
1	B	223	ASN	2.2
1	A	205	LYS	2.1
1	B	221	SER	2.1
1	B	228	TYR	2.1
1	A	202	SER	2.1
1	A	281	ILE	2.1
1	A	220	ASN	2.1
1	B	237	ASN	2.1
1	A	9	HIS	2.1
1	B	210	VAL	2.0
1	B	212	VAL	2.0
1	A	249	PRO	2.0
1	B	267	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	3732	6/6	0.91	0.15	32,37,41,45	0
3	GOL	A	3731	6/6	0.94	0.13	24,33,37,39	0
2	F3S	A	2302	7/7	0.95	0.12	21,25,29,35	0
2	F3S	B	3302	7/7	0.96	0.15	24,27,33,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.