



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:16 PM UTC

PDB ID : 5DN5 / pdb_00005dn5
Title : Structure of a C-terminally truncated glycoside hydrolase domain from
Salmonella typhimurium FlgJ
Authors : Zaloba, P.; Bailey-Elkin, B.A.; Mark, B.L.
Deposited on : 2015-09-09
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	
1	B	160	
1	C	160	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6522 atoms, of which 3080 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidoglycan hydrolase FlgJ.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	151	Total	C	H	N	O	S	0	0	0
			2188	707	1071	189	219	2			
1	B	151	Total	C	H	N	O	S	0	0	0
			2178	706	1059	192	219	2			
1	C	141	Total	C	H	N	O	S	0	0	0
			1964	645	950	174	194	1			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	MET	-	initiating methionine	UNP P15931
A	302	GLY	-	expression tag	UNP P15931
A	303	SER	-	expression tag	UNP P15931
A	304	HIS	-	expression tag	UNP P15931
A	305	HIS	-	expression tag	UNP P15931
A	306	HIS	-	expression tag	UNP P15931
A	307	HIS	-	expression tag	UNP P15931
A	308	HIS	-	expression tag	UNP P15931
A	309	HIS	-	expression tag	UNP P15931
B	150	MET	-	initiating methionine	UNP P15931
B	302	GLY	-	expression tag	UNP P15931
B	303	SER	-	expression tag	UNP P15931
B	304	HIS	-	expression tag	UNP P15931
B	305	HIS	-	expression tag	UNP P15931
B	306	HIS	-	expression tag	UNP P15931
B	307	HIS	-	expression tag	UNP P15931
B	308	HIS	-	expression tag	UNP P15931
B	309	HIS	-	expression tag	UNP P15931
C	150	MET	-	initiating methionine	UNP P15931
C	302	GLY	-	expression tag	UNP P15931
C	303	SER	-	expression tag	UNP P15931
C	304	HIS	-	expression tag	UNP P15931
C	305	HIS	-	expression tag	UNP P15931

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	306	HIS	-	expression tag	UNP P15931
C	307	HIS	-	expression tag	UNP P15931
C	308	HIS	-	expression tag	UNP P15931
C	309	HIS	-	expression tag	UNP P15931

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total I 2 2	0	0
2	B	3	Total I 3 3	0	0
2	C	5	Total I 5 5	0	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	2	Total Cl 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	65	Total O 65 65	0	0
5	B	66	Total O 66 66	0	0
5	C	48	Total O 48 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Peptidoglycan hydrolase FlgJ



- Molecule 1: Peptidoglycan hydrolase FlgJ



- Molecule 1: Peptidoglycan hydrolase FlgJ



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.70Å 61.12Å 65.22Å 90.00° 106.70° 90.00°	Depositor
Resolution (Å)	29.76 – 2.15 29.76 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.76-2.15) 99.6 (29.76-2.15)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.175 , 0.252 0.180 , 0.257	Depositor DCC
R_{free} test set	1096 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 62.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6522	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/1139	0.90	0/1554
1	B	0.91	0/1141	0.97	4/1559 (0.3%)
1	C	0.82	0/1032	0.95	1/1408 (0.1%)
All	All	0.87	0/3312	0.94	5/4521 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	152	SER	N-CA-C	-8.59	101.88	111.07
1	C	153	LYS	N-CA-C	-5.50	105.93	113.30
1	B	272	ALA	N-CA-C	-5.29	105.51	111.28
1	B	270	GLN	CA-C-N	-5.03	113.72	122.97
1	B	270	GLN	C-N-CA	-5.03	113.72	122.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1117	1071	1071	8	0
1	B	1119	1059	1059	15	0
1	C	1014	950	949	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	5	0	0	0	0
3	A	1	0	0	0	0
4	C	2	0	0	0	0
5	A	65	0	0	1	0
5	B	66	0	0	1	0
5	C	48	0	0	1	0
All	All	3442	3080	3079	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:195:GLU:OE1	5:C:501:HOH:O	2.00	0.79
1:C:190:ARG:NH2	1:C:203:VAL:O	2.22	0.72
1:C:290:LEU:O	1:C:293:MET:N	2.25	0.68
1:C:269:GLU:N	1:C:269:GLU:OE2	2.26	0.68
1:B:194:ARG:HH22	1:C:198:GLU:CD	2.13	0.57
1:B:155:PHE:CZ	1:B:159:LEU:HD12	2.40	0.56
1:A:289:LYS:NZ	5:A:501:HOH:O	2.27	0.56
1:B:194:ARG:HD3	1:B:238:VAL:HG21	1.89	0.54
1:B:285:ASN:OD1	1:B:286:TYR:N	2.41	0.54
1:B:283:ASP:N	1:B:283:ASP:OD1	2.44	0.51
1:A:247:SER:HB2	1:B:161:LEU:HD13	1.91	0.51
1:B:255:ARG:NH2	1:C:154:ASP:OD2	2.38	0.51
1:A:261:ALA:HB3	1:A:274:ALA:HB1	1.94	0.49
1:B:271:GLY:HA2	1:B:274:ALA:HB3	1.95	0.49
1:B:156:LEU:HD12	1:B:297:LEU:HD22	1.97	0.47
1:A:151:ASP:O	1:A:153:LYS:N	2.48	0.47
1:C:296:GLN:O	1:C:300:MET:HG3	2.15	0.47
1:C:290:LEU:O	1:C:291:THR:C	2.59	0.45
1:B:296:GLN:O	1:B:300:MET:HE3	2.17	0.44
1:B:176:LEU:HD21	1:B:275:LEU:HD11	2.00	0.44
1:B:278:ALA:HA	5:B:522:HOH:O	2.18	0.44
1:B:184:GLU:OE2	1:B:286:TYR:OH	2.27	0.43
1:A:194:ARG:HA	1:A:217:THR:OG1	2.19	0.43
1:A:161:LEU:C	1:A:161:LEU:HD23	2.44	0.43
1:C:277:ASN:O	1:C:278:ALA:HB2	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:PHE:CE1	1:A:159:LEU:HD22	2.55	0.42
1:A:178:LEU:HB3	1:A:297:LEU:CD1	2.49	0.42
1:B:276:GLN:C	1:B:278:ALA:H	2.27	0.41
1:C:221:THR:OG1	1:C:232:VAL:HG13	2.19	0.41
1:B:267:THR:O	1:B:270:GLN:O	2.38	0.41
1:C:254:THR:O	1:C:255:ARG:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/160 (93%)	144 (97%)	3 (2%)	2 (1%)	9	5
1	B	149/160 (93%)	144 (97%)	5 (3%)	0	100	100
1	C	137/160 (86%)	129 (94%)	8 (6%)	0	100	100
All	All	435/480 (91%)	417 (96%)	16 (4%)	2 (0%)	24	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	SER
1	A	278	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/127 (86%)	107 (98%)	2 (2%)	51	58
1	B	107/127 (84%)	104 (97%)	3 (3%)	38	40
1	C	91/127 (72%)	90 (99%)	1 (1%)	65	72
All	All	307/381 (81%)	301 (98%)	6 (2%)	48	54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	SER
1	A	273	VAL
1	B	194	ARG
1	B	203	VAL
1	B	219	ILE
1	C	247	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	GLN
1	A	189	GLN
1	B	276	GLN
1	B	295	GLN
1	C	180	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 13 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/160 (94%)	0.06	3 (1%) 65 69	16, 34, 61, 73	0
1	B	151/160 (94%)	0.09	9 (5%) 27 31	16, 31, 65, 80	0
1	C	141/160 (88%)	0.40	12 (8%) 16 18	14, 41, 64, 75	0
All	All	443/480 (92%)	0.18	24 (5%) 31 35	14, 36, 63, 80	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	301	SER	3.1
1	B	156	LEU	3.1
1	B	279	GLY	3.0
1	B	282	THR	3.0
1	C	273	VAL	2.9
1	B	301	SER	2.8
1	C	278	ALA	2.8
1	C	289	LYS	2.7
1	B	280	TYR	2.6
1	A	284	PRO	2.6
1	C	266	ALA	2.5
1	C	254	THR	2.4
1	A	226	ASN	2.4
1	C	277	ASN	2.4
1	B	281	ALA	2.4
1	B	285	ASN	2.4
1	B	286	TYR	2.3
1	C	288	ARG	2.3
1	C	299	ALA	2.1
1	C	297	LEU	2.1
1	B	299	ALA	2.1
1	C	156	LEU	2.0
1	C	292	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	282	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	C	406	1/1	0.89	0.17	37,37,37,37	0
4	CL	C	407	1/1	0.95	0.13	44,44,44,44	0
3	NA	A	403	1/1	0.98	0.11	33,33,33,33	0
2	IOD	C	401	1/1	0.98	0.04	47,47,47,47	1
2	IOD	C	402	1/1	0.98	0.04	40,40,40,40	1
2	IOD	C	403	1/1	0.99	0.02	36,36,36,36	1
2	IOD	B	401	1/1	0.99	0.03	39,39,39,39	1
2	IOD	B	402	1/1	1.00	0.01	25,25,25,25	0
2	IOD	C	404	1/1	1.00	0.02	23,23,23,23	0
2	IOD	C	405	1/1	1.00	0.01	22,22,22,22	0
2	IOD	B	403	1/1	1.00	0.01	22,22,22,22	0
2	IOD	A	402	1/1	1.00	0.02	26,26,26,26	0
2	IOD	A	401	1/1	1.00	0.01	20,20,20,20	1

6.5 Other polymers [i](#)

There are no such residues in this entry.