



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 11:11 PM UTC

PDB ID : 6DNA / pdb_00006dna
Title : Crystal structure of T110A mutant human Glutamate oxaloacetate transaminase 1 (GOT1)
Authors : Assar, Z.; Holt, M.C.; Stein, A.J.; Lairson, L.; Lyssiotis, C.A.
Deposited on : 2018-06-06
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	4% 70% 19% 5% . .
1	B	408	6% 72% 19% 5% . .
1	C	408	3% 73% 19% . . .
1	D	408	5% 71% 20% 5% . .
1	E	408	7% 68% 23% . . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	408	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PLP	E	501	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18520 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase, cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	2	0
			3029	1932	529	559	9			
1	B	395	Total	C	N	O	S	0	4	0
			3104	1985	537	573	9			
1	C	391	Total	C	N	O	S	0	2	0
			3049	1947	531	562	9			
1	D	396	Total	C	N	O	S	0	3	0
			3089	1968	542	570	9			
1	E	393	Total	C	N	O	S	0	3	0
			3095	1979	536	571	9			
1	F	383	Total	C	N	O	S	0	4	0
			3013	1925	523	556	9			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	THR	engineered mutation	UNP P17174
A	411	LYS	-	expression tag	UNP P17174
A	412	ILE	-	expression tag	UNP P17174
A	413	GLN	-	expression tag	UNP P17174
B	110	ALA	THR	engineered mutation	UNP P17174
B	411	LYS	-	expression tag	UNP P17174
B	412	ILE	-	expression tag	UNP P17174
B	413	GLN	-	expression tag	UNP P17174
C	110	ALA	THR	engineered mutation	UNP P17174
C	411	LYS	-	expression tag	UNP P17174
C	412	ILE	-	expression tag	UNP P17174
C	413	GLN	-	expression tag	UNP P17174
D	110	ALA	THR	engineered mutation	UNP P17174
D	411	LYS	-	expression tag	UNP P17174
D	412	ILE	-	expression tag	UNP P17174
D	413	GLN	-	expression tag	UNP P17174
E	110	ALA	THR	engineered mutation	UNP P17174

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	411	LYS	-	expression tag	UNP P17174
E	412	ILE	-	expression tag	UNP P17174
E	413	GLN	-	expression tag	UNP P17174
F	110	ALA	THR	engineered mutation	UNP P17174
F	411	LYS	-	expression tag	UNP P17174
F	412	ILE	-	expression tag	UNP P17174
F	413	GLN	-	expression tag	UNP P17174

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			10	6	4		
2	D	1	Total	C	O	0	0
			13	8	5		

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	C	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	D	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	E	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	F	1	Total	C	N	O	P	0	0
			16	8	1	6	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	O	0	0
			1	1		
4	C	3	Total	O	0	0
			3	3		
4	E	3	Total	O	0	0
			3	3		
4	F	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aspartate aminotransferase, cytoplasmic




- Molecule 1: Aspartate aminotransferase, cytoplasmic




- Molecule 1: Aspartate aminotransferase, cytoplasmic






- Molecule 1: Aspartate aminotransferase, cytoplasmic




- Molecule 1: Aspartate aminotransferase, cytoplasmic



- Molecule 1: Aspartate aminotransferase, cytoplasmic





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	257.70Å 148.63Å 83.18Å 90.00° 89.98° 90.00°	Depositor
Resolution (Å)	48.70 – 3.00 48.70 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.70-3.00) 99.1 (48.70-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	65.47 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.184 , 0.211 0.206 , 0.236	Depositor DCC
R_{free} test set	3047 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	83.8	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.000 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.076 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.077 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.000 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	18520	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.82	0/3108	1.15	4/4230 (0.1%)
1	B	0.87	2/3192 (0.1%)	1.11	4/4343 (0.1%)
1	C	0.83	2/3129 (0.1%)	1.07	0/4260
1	D	0.81	0/3171	1.15	7/4315 (0.2%)
1	E	0.86	2/3180 (0.1%)	1.10	3/4326 (0.1%)
1	F	0.79	0/3097	1.10	5/4211 (0.1%)
All	All	0.83	6/18877 (0.0%)	1.11	23/25685 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	4
1	C	0	2
1	D	0	1
1	E	0	6
1	F	0	3
All	All	0	22

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	370	GLN	N-CA	8.41	1.57	1.46
1	B	144	HIS	CE1-NE2	6.32	1.38	1.32
1	C	14	GLN	N-CA	6.20	1.54	1.46
1	E	226	TYR	C-O	-6.06	1.15	1.24
1	C	17	LEU	CA-C	5.95	1.55	1.52

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	318	GLU	CB-CG-CD	7.27	124.95	112.60
1	A	399	ASP	CA-CB-CG	-6.61	105.99	112.60
1	B	15	PRO	CA-N-CD	-6.43	103.00	112.00
1	F	342	LEU	CA-C-O	-6.21	113.24	120.20
1	D	347	THR	CA-C-N	6.09	127.45	119.84

There are no chirality outliers.

5 of 22 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	167	ARG	Sidechain
1	A	216	ARG	Sidechain
1	A	236	ARG	Sidechain
1	A	350	THR	Peptide
1	A	86	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3029	0	2902	88	0
1	B	3104	0	2993	125	0
1	C	3049	0	2934	72	0
1	D	3089	0	2985	114	0
1	E	3095	0	2995	139	0
1	F	3013	0	2915	191	0
2	A	13	0	18	0	0
2	B	10	0	13	0	0
2	D	13	0	18	0	0
3	A	16	0	7	1	0
3	B	16	0	7	2	0
3	C	16	0	7	2	0
3	D	16	0	7	2	0
3	E	16	0	7	9	0
3	F	16	0	7	4	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	3	0	0	0	0
4	E	3	0	0	0	0
4	F	2	0	0	0	0
All	All	18520	0	17815	691	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 691 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:21:LEU:HA	1:F:25:PHE:CD2	1.41	1.56
1:B:382:LEU:CD2	1:B:388:ILE:HG23	1.49	1.41
1:F:21:LEU:HA	1:F:25:PHE:CE2	1.59	1.37
1:F:17:LEU:O	1:F:20:LYS:CD	1.76	1.33
1:B:373:TYR:C	1:B:381:TYR:CE1	1.88	1.33

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/408 (94%)	344 (89%)	22 (6%)	19 (5%)	1	10
1	B	395/408 (97%)	358 (91%)	28 (7%)	9 (2%)	5	25
1	C	389/408 (95%)	355 (91%)	25 (6%)	9 (2%)	5	25
1	D	395/408 (97%)	355 (90%)	31 (8%)	9 (2%)	5	25
1	E	394/408 (97%)	347 (88%)	35 (9%)	12 (3%)	3	19
1	F	383/408 (94%)	353 (92%)	23 (6%)	7 (2%)	6	31
All	All	2341/2448 (96%)	2112 (90%)	164 (7%)	65 (3%)	4	21

5 of 65 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	26	ARG
1	A	29	PRO
1	A	33	LYS
1	A	347	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/346 (90%)	285 (91%)	28 (9%)	9	34
1	B	324/346 (94%)	290 (90%)	34 (10%)	6	27
1	C	316/346 (91%)	295 (93%)	21 (7%)	15	46
1	D	321/346 (93%)	287 (89%)	34 (11%)	6	27
1	E	323/346 (93%)	295 (91%)	28 (9%)	9	35
1	F	315/346 (91%)	283 (90%)	32 (10%)	7	29
All	All	1912/2076 (92%)	1735 (91%)	177 (9%)	9	32

5 of 177 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	404	SER
1	E	403	THR
1	E	97	LYS
1	E	338	LEU
1	F	24	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	35	ASN
1	F	128	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	302	GLN
1	C	323	ASN
1	C	176	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PG4	D	501	-	12,12,12	0.54	0	11,11,11	0.26	0
2	PG4	B	501	-	9,9,12	0.64	0	8,8,11	0.47	0
3	PLP	E	501	-	16,16,16	2.55	3 (18%)	20,23,23	2.05	5 (25%)
3	PLP	C	501	-	16,16,16	2.60	3 (18%)	20,23,23	1.62	5 (25%)
3	PLP	B	502	-	16,16,16	2.70	3 (18%)	20,23,23	1.42	3 (15%)
3	PLP	D	502	-	16,16,16	2.25	3 (18%)	20,23,23	1.97	5 (25%)
3	PLP	A	502	-	16,16,16	2.72	3 (18%)	20,23,23	1.82	5 (25%)
3	PLP	F	501	-	16,16,16	2.55	3 (18%)	20,23,23	1.84	7 (35%)
2	PG4	A	501	-	12,12,12	0.64	0	11,11,11	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	D	501	-	-	5/10/10/10	-
2	PG4	B	501	-	-	5/7/7/10	-
3	PLP	E	501	-	-	2/8/8/8	0/1/1/1
3	PLP	C	501	-	-	3/8/8/8	0/1/1/1
3	PLP	B	502	-	-	3/8/8/8	0/1/1/1
3	PLP	D	502	-	-	1/8/8/8	0/1/1/1
3	PLP	A	502	-	-	0/8/8/8	0/1/1/1
3	PLP	F	501	-	-	5/8/8/8	0/1/1/1
2	PG4	A	501	-	-	5/10/10/10	-

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	PLP	C3-C2	7.76	1.49	1.41
3	A	502	PLP	C3-C2	7.58	1.48	1.41
3	F	501	PLP	C3-C2	7.26	1.48	1.41
3	E	501	PLP	C3-C2	7.15	1.48	1.41
3	C	501	PLP	C3-C2	6.42	1.47	1.41

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	501	PLP	C4-C3-C2	-5.27	117.18	120.14
3	A	502	PLP	O4P-C5A-C5	5.12	118.96	109.36
3	D	502	PLP	O4P-C5A-C5	4.81	118.38	109.36
3	F	501	PLP	O4P-C5A-C5	4.60	117.98	109.36
3	E	501	PLP	O4P-C5A-C5	4.18	117.19	109.36

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	PLP	C5A-O4P-P-O1P
3	B	502	PLP	C5A-O4P-P-O2P
3	B	502	PLP	C5A-O4P-P-O3P
3	C	501	PLP	C5A-O4P-P-O2P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	E	501	PLP	C4-C5-C5A-O4P

There are no ring outliers.

6 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	501	PLP	9	0
3	C	501	PLP	2	0
3	B	502	PLP	2	0
3	D	502	PLP	2	0
3	A	502	PLP	1	0
3	F	501	PLP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	390/408 (95%)	0.36	17 (4%)	39	21	36, 68, 113, 168	3 (0%)
1	B	395/408 (96%)	0.45	23 (5%)	29	15	30, 75, 128, 228	5 (1%)
1	C	391/408 (95%)	0.31	12 (3%)	51	30	33, 72, 107, 156	3 (0%)
1	D	396/408 (97%)	0.38	20 (5%)	33	17	24, 70, 124, 208	4 (1%)
1	E	393/408 (96%)	0.47	30 (7%)	20	10	35, 75, 123, 213	3 (0%)
1	F	383/408 (93%)	0.42	14 (3%)	45	25	27, 77, 126, 159	5 (1%)
All	All	2348/2448 (95%)	0.40	116 (4%)	35	18	24, 73, 120, 228	23 (0%)

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	382	LEU	8.9
1	A	408	ALA	6.4
1	E	344	ALA	5.6
1	A	37	GLY	5.1
1	B	381	TYR	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PLP	B	502	16/16	0.80	0.19	101,111,159,163	10
2	PG4	A	501	13/13	0.82	0.14	87,99,115,123	0
2	PG4	B	501	10/13	0.85	0.11	75,81,88,91	0
3	PLP	A	502	16/16	0.87	0.21	96,121,141,145	10
2	PG4	D	501	13/13	0.87	0.14	85,99,107,108	0
3	PLP	C	501	16/16	0.89	0.21	122,136,158,166	10
3	PLP	F	501	16/16	0.90	0.13	75,90,106,110	10
3	PLP	E	501	16/16	0.91	0.18	102,116,133,137	16
3	PLP	D	502	16/16	0.92	0.15	78,101,143,148	10

6.5 Other polymers [i](#)

There are no such residues in this entry.