



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:34 PM UTC

PDB ID : 5DOR / pdb_00005dor
Title : P2 Integrase catalytic domain in space group P21
Authors : Skaar, K.; Claesson, M.; Odegrip, R.; Haggard-Ljungquist, E.; Hogbom, M.
Deposited on : 2015-09-11
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	81% 9% 8%
1	B	176	4% 83% 7% 9%
1	C	176	8% 78% 10% 10%
1	D	176	8% 81% 7% 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1264	804	227	227	6			
1	B	161	Total	C	N	O	S	0	1	0
			1266	805	229	226	6			
1	C	158	Total	C	N	O	S	0	0	0
			1235	788	223	218	6			
1	D	157	Total	C	N	O	S	0	0	0
			1224	780	219	219	6			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Zn 1	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total 10	O 10	0	0
4	B	28	Total 28	O 28	0	0
4	C	14	Total 14	O 14	0	0
4	D	22	Total 22	O 22	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Integrase





- Molecule 1: Integrase



- Molecule 1: Integrase



- Molecule 1: Integrase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.11Å 92.22Å 64.94Å 90.00° 96.34° 90.00°	Depositor
Resolution (Å)	64.54 – 2.50 64.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (64.54-2.50) 99.7 (64.54-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.230 , 0.275 0.224 , 0.270	Depositor DCC
R_{free} test set	1328 reflections (4.51%)	wwPDB-VP
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.569	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5069	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/1288	0.89	1/1740 (0.1%)
1	B	0.84	0/1290	1.01	2/1741 (0.1%)
1	C	0.76	0/1259	0.92	1/1700 (0.1%)
1	D	0.83	0/1248	0.92	0/1686
All	All	0.80	0/5085	0.94	4/6867 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	319	ASN	CB-CA-C	-8.73	92.96	110.17
1	A	200	ARG	NE-CZ-NH2	5.66	124.30	119.20
1	C	222	ARG	CD-NE-CZ	5.06	131.48	124.40
1	B	319	ASN	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1264	0	1283	12	1
1	B	1266	0	1287	9	0
1	C	1235	0	1261	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1224	0	1238	16	1
2	B	5	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	0	0	0
4	B	28	0	0	0	0
4	C	14	0	0	0	0
4	D	22	0	0	0	0
All	All	5069	0	5069	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:299:GLU:OE2	1:D:299:GLU:OE2	1.75	1.03
1:C:173:LEU:HA	1:C:258:ILE:CD1	2.05	0.84
1:A:318:LEU:HD11	1:D:253:ARG:HD2	1.73	0.71
1:C:299:GLU:CD	1:D:299:GLU:OE2	2.40	0.64
1:C:173:LEU:HA	1:C:258:ILE:HD11	1.81	0.63
1:A:326:GLU:O	1:A:327:ALA:HB2	1.98	0.62
1:C:173:LEU:HA	1:C:258:ILE:HD13	1.81	0.62
1:C:173:LEU:CA	1:C:258:ILE:HD13	2.31	0.61
1:C:299:GLU:OE2	1:D:299:GLU:CD	2.46	0.58
1:A:196:GLY:HA3	1:A:200:ARG:NH1	2.21	0.54
1:B:168:GLU:H	1:B:168:GLU:CD	2.15	0.53
1:C:222:ARG:HD3	1:C:294:GLY:HA3	1.90	0.53
1:C:297:ARG:HB3	1:C:299:GLU:OE1	2.10	0.52
1:A:322:ARG:NH1	1:D:257:ALA:HB2	2.26	0.51
1:B:315:ALA:O	1:B:319:ASN:ND2	2.45	0.49
1:B:268:THR:OG1	1:B:269:HIS:HD2	1.95	0.49
1:A:297:ARG:HB3	1:A:299:GLU:OE1	2.12	0.49
1:A:318:LEU:HD11	1:D:253:ARG:CD	2.41	0.49
1:B:297:ARG:HB3	1:B:299:GLU:OE1	2.13	0.48
1:D:297:ARG:HB3	1:D:299:GLU:OE1	2.12	0.48
1:A:190:SER:HA	1:B:319:ASN:ND2	2.28	0.48
1:C:173:LEU:CA	1:C:258:ILE:CD1	2.84	0.48
1:C:210:ARG:CG	1:C:223:THR:CG2	2.93	0.47
1:A:311:TYR:CZ	1:D:250:ARG:HD2	2.50	0.47
1:A:235:ALA:O	1:A:236:ASP:CG	2.58	0.46
1:C:256:LYS:HE2	1:C:256:LYS:HB2	1.82	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:326:GLU:O	1:A:327:ALA:CB	2.62	0.45
1:C:264:MET:O	1:C:264:MET:HG3	2.15	0.45
1:C:170:ILE:HG22	1:D:327:ALA:HB2	1.99	0.45
1:A:168:GLU:CD	1:A:168:GLU:H	2.25	0.45
1:B:214:VAL:HG22	1:B:215:LYS:H	1.83	0.43
1:C:251:PHE:CZ	1:C:255:MET:HE2	2.53	0.43
1:C:299:GLU:OE2	1:D:299:GLU:HG3	2.18	0.43
1:C:319:ASN:HB2	1:C:320:PRO:HD2	2.01	0.43
1:B:233:MET:HE2	1:B:233:MET:HB3	1.93	0.43
1:C:210:ARG:HG3	1:C:223:THR:CG2	2.48	0.43
1:C:299:GLU:CG	1:D:299:GLU:OE2	2.66	0.42
1:B:251:PHE:CZ	1:B:255:MET:HE2	2.55	0.42
1:C:233:MET:HE2	1:C:233:MET:HB3	1.93	0.42
1:D:251:PHE:CZ	1:D:255:MET:HE2	2.55	0.41
1:A:251:PHE:CZ	1:A:255:MET:HE2	2.55	0.41
1:B:214:VAL:O	1:B:215:LYS:C	2.64	0.40
1:D:235:ALA:O	1:D:236:ASP:HB2	2.20	0.40
1:D:327:ALA:C	1:D:328:GLU:HG2	2.45	0.40
1:C:299:GLU:OE2	1:D:299:GLU:CG	2.69	0.40
1:D:236:ASP:C	1:D:236:ASP:OD1	2.65	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:ARG:NH2	1:D:306:HIS:O[1_455]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	156/176 (89%)	152 (97%)	3 (2%)	1 (1%)	21 38

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	156/176 (89%)	152 (97%)	4 (3%)	0	100	100
1	C	152/176 (86%)	150 (99%)	2 (1%)	0	100	100
1	D	151/176 (86%)	150 (99%)	1 (1%)	0	100	100
All	All	615/704 (87%)	604 (98%)	10 (2%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	327	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/145 (90%)	126 (96%)	5 (4%)	29	56
1	B	131/145 (90%)	126 (96%)	5 (4%)	29	56
1	C	128/145 (88%)	123 (96%)	5 (4%)	28	55
1	D	126/145 (87%)	123 (98%)	3 (2%)	43	70
All	All	516/580 (89%)	498 (96%)	18 (4%)	32	58

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	GLU
1	A	266	GLN
1	A	321	LEU
1	A	326	GLU
1	A	328	GLU
1	B	168	GLU
1	B	311	TYR
1	B	319	ASN
1	B	321	LEU
1	B	328	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	201	LEU
1	C	253	ARG
1	C	264	MET
1	C	296	THR
1	C	312	LEU
1	D	168	GLU
1	D	312	LEU
1	D	328	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	278	HIS
1	A	300	GLN
1	B	269	HIS
1	B	300	GLN
1	B	319	ASN
1	C	278	HIS
1	D	300	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	B	401	-	4,4,4	0.68	0	6,6,6	0.83	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/176 (92%)	0.17	2 (1%) 76 73	40, 58, 83, 108	0
1	B	161/176 (91%)	0.12	7 (4%) 40 35	21, 47, 76, 93	1 (0%)
1	C	158/176 (89%)	0.54	14 (8%) 15 13	41, 63, 102, 121	0
1	D	157/176 (89%)	0.43	14 (8%) 15 13	30, 53, 97, 128	0
All	All	638/704 (90%)	0.31	37 (5%) 29 25	21, 55, 93, 128	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	240	GLY	5.7
1	D	214	VAL	4.5
1	D	264	MET	4.4
1	B	162	THR	4.2
1	D	262	LEU	4.0
1	B	311	TYR	3.9
1	C	262	LEU	3.6
1	C	163	GLY	3.5
1	C	257	ALA	3.5
1	D	292	ILE	3.4
1	D	263	PRO	3.3
1	C	267	ALA	3.1
1	B	329	SER	3.1
1	C	260	PRO	3.1
1	B	264	MET	2.9
1	D	291	ARG	2.8
1	C	255	MET	2.8
1	C	264	MET	2.7
1	C	256	LYS	2.6
1	C	235	ALA	2.6
1	C	268	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	264	MET	2.6
1	C	189	LEU	2.5
1	B	240	GLY	2.4
1	D	298	ILE	2.3
1	D	221	PRO	2.3
1	C	265	GLY	2.3
1	C	261	ASP	2.3
1	B	214	VAL	2.2
1	D	163	GLY	2.2
1	A	326	GLU	2.2
1	D	260	PRO	2.1
1	D	241	PHE	2.1
1	B	220	LYS	2.1
1	D	317	SER	2.1
1	D	208	HIS	2.1
1	C	274	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	B	401	5/5	0.87	0.16	69,75,92,95	0
3	ZN	B	402	1/1	0.91	0.14	114,114,114,114	0

6.5 Other polymers [i](#)

There are no such residues in this entry.