



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:25 AM UTC

PDB ID : 1DPD / pdb_00001dpd
Title : CRYSTALLOGRAPHIC AND ENZYMATIC INVESTIGATIONS ON THE
ROLE OF SER558, HIS610 AND ASN614 IN THE CATALYTIC
MECHANISM OF AZOTOBACTER VINELANDII DIHYDROLIPOAMIDE
ACETYLTRANSFERASE (E2P)
Authors : Hendle, J.; Hol, W.G.J.
Deposited on : 1995-02-03
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIHYDROLIPOYL-TRANSACETYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	0	0
			1811	1168	310	326	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	558	ALA	SER	conflict	UNP P10802

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	51	Total	O	0	0
			51	51		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DIHYDROLIPOYL-TRANSACETYLASE

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	F 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	224.67Å 224.67Å 224.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.70 10.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.70) 97.5 (10.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.89 (at 2.71Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.184 , (Not available) 0.168 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1862	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	8/1848 (0.4%)	1.77	35/2518 (1.4%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	502	HIS	CD2-NE2	-7.36	1.29	1.37
1	A	426	HIS	CD2-NE2	-6.41	1.30	1.37
1	A	562	HIS	CD2-NE2	-6.38	1.30	1.37
1	A	434	HIS	CD2-NE2	-6.15	1.31	1.37
1	A	610	HIS	CD2-NE2	-5.71	1.31	1.37
1	A	502	HIS	CG-ND1	-5.60	1.32	1.38
1	A	426	HIS	CG-ND1	-5.25	1.32	1.38
1	A	554	CYS	CA-CB	-5.19	1.46	1.54

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	577	VAL	N-CA-CB	-10.34	98.92	112.26
1	A	563	ILE	N-CA-C	-9.08	104.45	111.62
1	A	614	ASN	OD1-CG-ND2	-8.67	113.93	122.60
1	A	424	ASN	OD1-CG-ND2	-7.23	115.37	122.60
1	A	592	ASP	CA-CB-CG	7.04	119.64	112.60
1	A	497	ARG	CA-CB-CG	-7.01	100.08	114.10
1	A	566	THR	N-CA-CB	-6.92	100.36	110.47
1	A	568	PHE	CA-CB-CG	6.85	120.65	113.80
1	A	577	VAL	CB-CA-C	6.59	119.38	111.20
1	A	627	ASP	CA-CB-CG	6.17	118.77	112.60
1	A	614	ASN	CB-CG-ND2	6.06	125.50	116.40
1	A	599	ARG	NE-CZ-NH2	6.03	124.63	119.20
1	A	637	LEU	CA-CB-CG	5.83	136.69	116.30
1	A	402	ASP	CA-CB-CG	5.77	118.37	112.60
1	A	563	ILE	CA-C-O	-5.76	115.67	120.80
1	A	509	THR	O-C-N	-5.65	117.07	121.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	551	GLN	N-CA-C	5.65	118.27	109.52
1	A	434	HIS	CB-CG-CD2	-5.51	124.04	131.20
1	A	610	HIS	CB-CG-CD2	-5.51	124.04	131.20
1	A	481	LEU	CA-C-N	5.44	125.11	119.56
1	A	481	LEU	C-N-CA	5.44	125.11	119.56
1	A	562	HIS	CB-CG-CD2	-5.40	124.18	131.20
1	A	591	TRP	CG-CD1-NE1	-5.39	103.19	110.20
1	A	420	ILE	N-CA-C	-5.24	105.39	110.42
1	A	424	ASN	CA-CB-CG	-5.22	107.38	112.60
1	A	591	TRP	CE2-CD2-CG	-5.18	100.98	107.20
1	A	427	ARG	N-CA-CB	-5.18	102.50	110.01
1	A	453	GLN	OE1-CD-NE2	-5.16	117.44	122.60
1	A	509	THR	CA-C-N	5.16	124.47	118.85
1	A	509	THR	C-N-CA	5.16	124.47	118.85
1	A	593	GLY	N-CA-C	-5.15	108.95	115.08
1	A	597	GLN	OE1-CD-NE2	-5.15	117.45	122.60
1	A	621	PHE	CA-C-O	-5.05	115.51	120.82
1	A	566	THR	CA-CB-OG1	-5.03	102.06	109.60
1	A	588	GLN	OE1-CD-NE2	-5.00	117.60	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1811	0	1855	19	0
2	A	51	0	0	1	0
All	All	1862	0	1855	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:HIS:HD2	1:A:520:ASN:H	1.46	0.63
1:A:566:THR:HG21	2:A:715:HOH:O	2.05	0.56
1:A:417:LEU:HD12	1:A:511:ASP:HB3	1.90	0.53
1:A:599:ARG:HB2	1:A:601:MET:HE3	1.93	0.51
1:A:545:LEU:HD23	1:A:550:MET:SD	2.51	0.51
1:A:590:VAL:HG21	1:A:601:MET:HE3	1.92	0.51
1:A:416:ARG:O	1:A:420:ILE:HG12	2.10	0.50
1:A:586:SER:HB2	1:A:588:GLN:HE21	1.77	0.49
1:A:473:ALA:HA	1:A:635:ILE:HD11	1.95	0.48
1:A:562:HIS:H	1:A:562:HIS:CD2	2.35	0.45
1:A:478:LEU:HD23	1:A:501:VAL:HG13	1.99	0.45
1:A:502:HIS:CD2	1:A:520:ASN:H	2.32	0.45
1:A:446:LEU:HD21	1:A:469:LEU:HD22	2.00	0.43
1:A:446:LEU:CD2	1:A:469:LEU:HD22	2.50	0.41
1:A:632:ILE:HD11	1:A:635:ILE:HD12	2.03	0.41
1:A:634:ALA:O	1:A:637:LEU:HD23	2.20	0.41
1:A:577:VAL:HG13	1:A:613:ILE:CD1	2.52	0.40
1:A:624:ARG:HA	1:A:624:ARG:HD2	1.91	0.40
1:A:632:ILE:CD1	1:A:635:ILE:HD12	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/243 (99%)	233 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	186/194 (96%)	165 (89%)	21 (11%)	5 14

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	408	GLU
1	A	414	MET
1	A	417	LEU
1	A	446	LEU
1	A	451	VAL
1	A	467	LEU
1	A	470	LEU
1	A	472	LYS
1	A	478	LEU
1	A	479	LYS
1	A	497	ARG
1	A	511	ASP
1	A	543	LYS
1	A	550	MET
1	A	566	THR
1	A	577	VAL
1	A	580	LEU
1	A	597	GLN
1	A	599	ARG
1	A	613	ILE
1	A	636	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	419	GLN
1	A	431	ASN
1	A	453	GLN
1	A	502	HIS
1	A	523	GLN
1	A	588	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/243 (100%)	-0.76	1 (0%) 88 87	2, 14, 45, 66	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	402	ASP	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.