



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:54 AM UTC

PDB ID : 2DPP / pdb_00002dpp
Title : Crystal structure of thermostable Bacillus sp. RAPc8 nitrile hydratase
Authors : Tsekoa, T.L.; Tastan-Bishop, A.O.; Cameron, R.A.; Sewell, B.T.; Sayed, M.F.;
Cowan, D.A.
Deposited on : 2006-05-12
Resolution : 2.52 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	 70% 18% 5% • 6%
2	B	229	 71% 21% • • •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrile hydratase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	1631	1039	284	301	7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	119	CSD	CYS	modified residue	UNP Q84FS5
A	121	CSO	CYS	modified residue	UNP Q84FS5

- Molecule 2 is a protein called Nitrile hydratase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	227	1857	1193	313	343	8	0	0	0

- Molecule 3 is COBALT (II) ION (CCD ID: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Co	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	87	Total	O	0	0
			87	87		
4	B	116	Total	O	0	0
			116	116		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitrile hydratase alpha subunit

Chain A:  70% 18% 5% • 6%



- Molecule 2: Nitrile hydratase beta subunit

Chain B:  71% 21% • • •



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.62Å 106.62Å 83.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.52 20.00 – 2.52	Depositor EDS
% Data completeness (in resolution range)	93.0 (20.00-2.52) 99.6 (20.00-2.52)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.53Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.198 , 0.279 0.195 , 0.273	Depositor DCC
R_{free} test set	831 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3692	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CO, CSD, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.47	11/1660 (0.7%)	1.31	11/2256 (0.5%)
2	B	1.48	9/1907 (0.5%)	1.35	10/2577 (0.4%)
All	All	1.48	20/3567 (0.6%)	1.33	21/4833 (0.4%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	ILE	CA-CB	8.29	1.65	1.54
1	A	75	ALA	CA-CB	7.67	1.65	1.53
2	B	171	VAL	CA-CB	6.51	1.62	1.54
1	A	162	VAL	CA-CB	6.40	1.62	1.54
2	B	210	LYS	CD-CE	6.35	1.71	1.52
1	A	94	TYR	C-O	-6.07	1.16	1.23
1	A	73	ASP	CA-C	6.00	1.58	1.52
2	B	210	LYS	CB-CG	5.92	1.70	1.52
1	A	61	MET	CG-SD	5.67	1.95	1.80
2	B	13	PHE	C-O	-5.59	1.16	1.24
1	A	166	VAL	CA-CB	5.46	1.61	1.54
2	B	223	TYR	N-CA	5.45	1.53	1.46
1	A	197	VAL	CA-C	5.44	1.59	1.53
1	A	67	VAL	CA-CB	5.31	1.60	1.54
1	A	37	ILE	CA-CB	5.27	1.60	1.54
2	B	134	ARG	CA-C	-5.16	1.46	1.52
1	A	130	PRO	CA-C	5.14	1.56	1.52
2	B	210	LYS	CE-NZ	5.08	1.64	1.49
1	A	114	VAL	N-CA	-5.05	1.40	1.46
2	B	210	LYS	CG-CD	5.00	1.67	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	VAL	N-CA-CB	7.21	118.57	110.72
2	B	159	THR	CB-CA-C	-7.04	94.97	110.67
1	A	165	ARG	NE-CZ-NH2	6.88	125.39	119.20
2	B	166	ARG	NE-CZ-NH1	-6.85	114.65	121.50
2	B	166	ARG	NE-CZ-NH2	6.15	124.73	119.20
2	B	55	PHE	N-CA-C	-5.95	104.88	111.36
2	B	104	LYS	N-CA-C	5.90	117.71	111.28
1	A	165	ARG	CB-CG-CD	5.56	124.09	111.30
2	B	103	LEU	N-CA-C	-5.31	104.39	111.24
2	B	72	TYR	N-CA-C	5.30	116.74	111.07
1	A	162	VAL	CB-CA-C	5.27	117.45	110.91
1	A	165	ARG	NE-CZ-NH1	-5.21	116.28	121.50
1	A	57	GLU	N-CA-C	5.20	117.69	111.71
2	B	173	ASP	N-CA-C	-5.19	107.08	113.41
2	B	166	ARG	CB-CG-CD	5.13	123.11	111.30
1	A	188	MET	CB-CA-C	5.10	118.22	109.51
1	A	79	ARG	CA-C-N	5.08	127.40	120.54
1	A	79	ARG	C-N-CA	5.08	127.40	120.54
1	A	189	THR	N-CA-C	5.04	117.03	110.53
1	A	78	GLN	N-CA-C	5.03	116.77	111.28
2	B	65	VAL	CB-CA-C	-5.01	104.40	112.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	1611	25	0
2	B	1857	0	1805	44	0
3	A	1	0	0	0	0
4	A	87	0	0	3	0
4	B	116	0	0	8	0
All	All	3692	0	3416	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:5:HIS:HD2	2:B:60:GLU:HG2	1.05	1.10
2:B:25:ILE:HD11	2:B:28:THR:HG22	1.35	1.07
2:B:5:HIS:CD2	2:B:60:GLU:HG2	1.94	1.01
2:B:210:LYS:HE3	2:B:210:LYS:HA	1.46	0.94
1:A:20:ARG:HD2	4:B:287:HOH:O	1.68	0.93
2:B:2:ASN:HD21	2:B:166:ARG:HH12	1.07	0.93
2:B:5:HIS:HD2	2:B:60:GLU:CG	1.88	0.84
2:B:2:ASN:ND2	2:B:166:ARG:HH12	1.82	0.77
1:A:14:PHE:HB3	1:A:17:HIS:CD2	2.20	0.76
2:B:5:HIS:ND1	2:B:6:ASP:N	2.33	0.76
2:B:159:THR:HG23	2:B:183:ASP:OD1	1.88	0.73
1:A:53:HIS:CD2	1:A:58:LEU:HD22	2.24	0.73
2:B:2:ASN:HD21	2:B:166:ARG:NH1	1.85	0.72
1:A:16:HIS:H	1:A:16:HIS:CD2	2.08	0.70
2:B:54:GLU:OE1	4:B:341:HOH:O	2.09	0.69
2:B:5:HIS:ND1	2:B:5:HIS:C	2.50	0.68
1:A:159:PRO:HD2	1:A:162:VAL:HG13	1.77	0.66
2:B:5:HIS:CD2	2:B:60:GLU:CG	2.71	0.65
2:B:60:GLU:OE1	2:B:158:HIS:HD2	1.81	0.64
2:B:159:THR:HB	2:B:161:PHE:H	1.64	0.62
1:A:41:HIS:HE1	4:B:279:HOH:O	1.82	0.62
2:B:160:ARG:O	2:B:162:PRO:HD3	1.99	0.61
2:B:15:LYS:HG2	2:B:17:MET:CE	2.31	0.61
2:B:88:GLY:HA2	4:B:292:HOH:O	2.00	0.61
2:B:210:LYS:HA	2:B:210:LYS:CE	2.28	0.59
1:A:89:ARG:HG3	1:A:94:TYR:CD1	2.38	0.59
2:B:25:ILE:HD11	2:B:28:THR:CG2	2.24	0.58
1:A:36:LEU:HD13	1:A:42:LEU:HD22	1.84	0.57
2:B:171:VAL:HG22	4:B:282:HOH:O	2.02	0.57
2:B:4:ILE:HD12	2:B:4:ILE:N	2.20	0.57
2:B:93:LYS:HB2	2:B:93:LYS:NZ	2.20	0.56
1:A:16:HIS:CD2	1:A:16:HIS:N	2.73	0.55
2:B:22:GLU:HG2	4:B:255:HOH:O	2.05	0.55
1:A:165:ARG:HD3	4:A:309:HOH:O	2.08	0.54
2:B:159:THR:CG2	2:B:183:ASP:OD1	2.56	0.53
1:A:53:HIS:CD2	1:A:58:LEU:CD2	2.91	0.53
1:A:121:CSO:HB2	1:A:174:ARG:CZ	2.39	0.53
2:B:5:HIS:CE1	2:B:6:ASP:HB2	2.44	0.53
1:A:16:HIS:H	1:A:16:HIS:HD2	1.53	0.52
2:B:124:LYS:NZ	2:B:128:ASP:OD2	2.43	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:HIS:HE1	4:B:257:HOH:O	1.94	0.51
4:A:356:HOH:O	2:B:75:HIS:HD2	1.94	0.51
1:A:159:PRO:HD2	1:A:162:VAL:CG1	2.41	0.50
2:B:67:TYR:O	2:B:75:HIS:HE1	1.98	0.47
1:A:172:GLU:HG3	2:B:132:PRO:HG3	1.97	0.47
2:B:183:ASP:OD2	2:B:183:ASP:N	2.48	0.47
4:A:310:HOH:O	2:B:214:SER:HB2	2.14	0.46
1:A:115:VAL:HG22	1:A:116:CYS:N	2.31	0.46
2:B:148:ILE:C	2:B:148:ILE:HD12	2.40	0.46
2:B:210:LYS:CE	2:B:211:GLN:H	2.28	0.45
2:B:210:LYS:HE2	2:B:211:GLN:H	1.81	0.45
2:B:4:ILE:HD12	2:B:4:ILE:H	1.82	0.44
2:B:5:HIS:CD2	2:B:60:GLU:CD	2.96	0.44
1:A:188:MET:HE3	1:A:192:GLU:HB3	2.01	0.43
2:B:136:ILE:HD11	2:B:195:TYR:CE1	2.54	0.43
1:A:59:GLY:HA3	1:A:61:MET:SD	2.59	0.43
1:A:177:VAL:O	1:A:179:PRO:HD3	2.20	0.42
2:B:103:LEU:HD12	2:B:103:LEU:HA	1.96	0.42
1:A:53:HIS:HA	1:A:57:GLU:HB2	2.02	0.42
1:A:120:SER:HB2	1:A:135:LYS:HG2	2.02	0.41
1:A:165:ARG:HB3	2:B:216:TYR:CE1	2.56	0.41
2:B:91:ASP:N	4:B:313:HOH:O	2.54	0.41
2:B:150:THR:HB	2:B:161:PHE:CZ	2.56	0.41
1:A:42:LEU:HD12	1:A:42:LEU:HA	1.98	0.41
1:A:185:THR:O	1:A:188:MET:HG2	2.21	0.41
1:A:201:SER:HA	1:A:206:ALA:O	2.21	0.41
2:B:93:LYS:HB2	2:B:93:LYS:HZ2	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/216 (92%)	191 (96%)	6 (3%)	1 (0%)	24	41
2	B	225/229 (98%)	216 (96%)	9 (4%)	0	100	100
All	All	423/445 (95%)	407 (96%)	15 (4%)	1 (0%)	43	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/190 (93%)	162 (92%)	15 (8%)	10	20
2	B	194/196 (99%)	159 (82%)	35 (18%)	2	3
All	All	371/386 (96%)	321 (86%)	50 (14%)	4	7

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	HIS
1	A	20	ARG
1	A	28	ARG
1	A	32	LEU
1	A	35	LEU
1	A	42	LEU
1	A	52	LYS
1	A	58	LEU
1	A	82	GLU
1	A	114	VAL
1	A	126	LEU
1	A	162	VAL
1	A	165	ARG
1	A	186	GLU
1	A	191	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	4	ILE
2	B	5	HIS
2	B	15	LYS
2	B	19	VAL
2	B	25	ILE
2	B	34	LEU
2	B	47	LEU
2	B	64	PRO
2	B	65	VAL
2	B	84	LEU
2	B	92	GLU
2	B	93	LYS
2	B	95	LEU
2	B	97	GLU
2	B	103	LEU
2	B	104	LYS
2	B	105	LYS
2	B	107	ASP
2	B	109	LYS
2	B	112	ARG
2	B	118	LEU
2	B	120	LYS
2	B	121	LEU
2	B	124	LYS
2	B	139	SER
2	B	143	LYS
2	B	149	LYS
2	B	156	THR
2	B	159	THR
2	B	166	ARG
2	B	171	VAL
2	B	183	ASP
2	B	204	GLU
2	B	210	LYS
2	B	214	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	HIS
1	A	17	HIS
1	A	41	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	180	GLN
2	B	2	ASN
2	B	29	HIS
2	B	75	HIS
2	B	152	ASN
2	B	158	HIS
2	B	194	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	119	1,3	4,7,8	13.72	2 (50%)	1,8,10	7.02	1 (100%)
1	CSO	A	121	1,3	3,6,7	2.16	1 (33%)	1,6,8	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	119	1,3	-	0/2/6/8	-
1	CSO	A	121	1,3	-	0/1/5/7	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	119	CSD	OD1-SG	27.35	1.72	1.47
1	A	121	CSO	O-C	3.65	1.33	1.20
1	A	119	CSD	O-C	2.03	1.27	1.20

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	CSD	OD1-SG-CB	7.02	118.54	105.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	121	CSO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/216 (92%)	-0.71	0 100 100	17, 26, 42, 49	0
2	B	227/229 (99%)	-0.59	0 100 100	15, 25, 46, 62	0
All	All	427/445 (95%)	-0.65	0 100 100	15, 26, 44, 62	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	119	8/9	0.98	0.05	15,18,19,22	0
1	CSO	A	121	7/8	0.98	0.06	19,20,21,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CO	A	301	1/1	0.99	0.02	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.