



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:21 AM UTC

PDB ID : 4DPR / pdb_00004dpr
Title : Structure of human Leukotriene A4 hydrolase in complex with inhibitor captopril
Authors : Stsiapanava, A.; Haeggstrom, J.Z.; Rinaldo-Matthis, A.
Deposited on : 2012-02-14
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	 82% 16% ..

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Leukotriene A-4 hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	607	5052	3253	831	947	21	0	37	0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	1	1	0	0

- Molecule 3 is L-CAPTOPRIL (CCD ID: X8Z) (formula: C₉H₁₅NO₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	14	9	1	3	1	0	0

- Molecule 4 is YTTERBIUM (III) ION (CCD ID: YB) (formula: Yb).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Yb		
4	A	5	5	5	0	0

- Molecule 5 is ACETIC ACID (CCD ID: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	8	4	4	0	1

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	414	Total	O	0	21
			435	435		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leukotriene A-4 hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.55Å 87.40Å 99.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.69 – 2.02 65.69 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.9 (65.69-2.02) 99.9 (65.69-2.02)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.174 , 0.228 0.176 , 0.231	Depositor DCC
R_{free} test set	2275 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.3	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5527	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, YB, GOL, X8Z, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.31	5/5282 (0.1%)	1.14	10/7174 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	208	ALA	CA-CB	7.71	1.63	1.52
1	A	149	THR	CA-CB	5.72	1.63	1.53
1	A	470	ALA	CA-CB	5.49	1.62	1.53
1	A	214	ILE	N-CA	5.38	1.51	1.46
1	A	550	ALA	CA-CB	5.21	1.62	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	ASN	CA-C-N	-6.34	113.42	119.76
1	A	272	ASN	C-N-CA	-6.34	113.42	119.76
1	A	214	ILE	N-CA-C	5.88	118.84	113.20
1	A	143	ILE	CB-CA-C	-5.46	106.06	111.30
1	A	541	LEU	N-CA-C	-5.42	105.28	111.07
1	A	178	THR	CA-C-N	-5.22	114.41	119.78
1	A	178	THR	C-N-CA	-5.22	114.41	119.78
1	A	598	VAL	N-CA-C	-5.16	105.81	110.82
1	A	405	GLY	N-CA-C	-5.13	106.49	113.37
1	A	56	THR	N-CA-C	-5.05	101.40	109.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5052	0	5092	67	1
2	A	1	0	0	0	0
3	A	14	0	13	3	0
4	A	5	0	0	0	1
5	A	8	0	6	0	0
6	A	12	0	16	1	0
7	A	435	0	0	13	0
All	All	5527	0	5127	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352[A]:SER:OG	7:A:970[A]:HOH:O	1.66	1.13
1:A:360:HIS:HD2	1:A:362:PHE:H	1.30	0.77
1:A:261[A]:LEU:CD1	1:A:277:PHE:HB3	2.21	0.70
1:A:261[B]:LEU:HD22	1:A:265:PHE:CD1	2.26	0.70
1:A:527:ASN:ND2	1:A:538:TRP:HE1	1.90	0.70
3:A:702:X8Z:S	7:A:940:HOH:O	2.49	0.70
1:A:527:ASN:HD22	1:A:538:TRP:HE1	1.38	0.69
1:A:261[A]:LEU:HD11	1:A:277:PHE:HB3	1.76	0.68
1:A:521:GLN:HE22	1:A:527:ASN:H	1.42	0.66
1:A:17:ARG:HD2	1:A:45[A]:GLN:CD	2.24	0.62
1:A:19[B]:LYS:HE3	7:A:849:HOH:O	2.01	0.59
1:A:556:LYS:HE2	1:A:560[A]:GLU:OE2	2.06	0.56
1:A:247:LEU:HD21	1:A:331[A]:ARG:NH2	2.20	0.56
1:A:596:HIS:HE1	7:A:801:HOH:O	1.89	0.55
1:A:324:LEU:HD21	1:A:407:LEU:HD22	1.87	0.54
1:A:484:ASN:HB3	7:A:906:HOH:O	2.08	0.53
1:A:205:VAL:HG11	1:A:259:LEU:HD22	1.91	0.52
1:A:235:SER:OG	1:A:287[A]:LYS:NZ	2.35	0.52
1:A:62[B]:GLU:O	1:A:63:LYS:HB3	2.10	0.52
1:A:564:MET:SD	1:A:602[A]:LEU:HD22	2.50	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:GLY:HA2	1:A:350:GLN:HE21	1.75	0.51
1:A:58:ASP:OD1	1:A:78:ARG:NH1	2.43	0.51
1:A:308:ASN:OD1	1:A:314:PHE:HA	2.11	0.51
1:A:104:ILE:HG21	1:A:145:PRO:HG2	1.93	0.50
1:A:19[B]:LYS:CE	7:A:849:HOH:O	2.59	0.50
1:A:159:GLU:HG3	1:A:190:LYS:HG2	1.95	0.49
1:A:171:SER:HB3	1:A:272:ASN:HB3	1.94	0.49
1:A:95:SER:HB3	7:A:1163:HOH:O	2.11	0.49
1:A:401[B]:GLU:CD	1:A:401[B]:GLU:H	2.21	0.49
1:A:598:VAL:O	1:A:602[A]:LEU:HD13	2.13	0.48
1:A:474:ARG:NH1	7:A:956:HOH:O	2.30	0.48
1:A:128:HIS:CE1	1:A:165:GLU:HG3	2.50	0.47
1:A:360:HIS:CD2	1:A:362:PHE:H	2.21	0.47
1:A:232:TYR:CZ	1:A:287[B]:LYS:HE3	2.51	0.46
1:A:28[A]:ASP:OD2	1:A:31[A]:ARG:NE	2.49	0.46
1:A:261[A]:LEU:HD11	1:A:277:PHE:CG	2.52	0.45
1:A:174[A]:ARG:HG3	7:A:831:HOH:O	2.16	0.45
1:A:90:LEU:HD22	1:A:100:ILE:HD11	1.98	0.45
1:A:292:VAL:O	1:A:296:GLU:HG2	2.17	0.45
1:A:447:TRP:CZ3	1:A:454:PRO:HB3	2.51	0.45
1:A:13:ALA:O	1:A:45[A]:GLN:HG3	2.17	0.45
1:A:261[A]:LEU:HD11	1:A:277:PHE:CB	2.44	0.44
1:A:408:LYS:O	1:A:412[B]:GLU:HG3	2.18	0.44
1:A:23:LEU:O	1:A:158:ALA:HA	2.18	0.44
1:A:220[A]:VAL:HG11	1:A:230:SER:HB3	1.98	0.44
1:A:329:CYS:SG	7:A:964:HOH:O	2.47	0.44
1:A:177:GLU:OE1	1:A:187:LYS:NZ	2.45	0.44
1:A:27:VAL:HG21	1:A:206:VAL:HG21	1.99	0.43
1:A:19[A]:LYS:HD2	7:A:882:HOH:O	2.19	0.42
1:A:265:PHE:CD1	1:A:266:PRO:HD2	2.54	0.42
1:A:305:LEU:HD23	1:A:305:LEU:HA	1.88	0.42
1:A:596:HIS:HD2	1:A:598:VAL:H	1.65	0.42
1:A:261[B]:LEU:HD22	1:A:265:PHE:CG	2.54	0.42
1:A:383:TYR:CE2	3:A:702:X8Z:H52C	2.54	0.42
1:A:247:LEU:HD21	1:A:331[A]:ARG:HH22	1.84	0.41
1:A:601:MET:SD	1:A:602[A]:LEU:HD12	2.60	0.41
1:A:564:MET:HE3	7:A:1066[A]:HOH:O	2.20	0.41
1:A:596:HIS:CD2	1:A:598:VAL:H	2.37	0.41
1:A:95:SER:HB2	1:A:98[B]:GLN:OE1	2.20	0.41
1:A:17:ARG:NE	7:A:1021[B]:HOH:O	2.53	0.41
1:A:175:ASP:OD2	1:A:175:ASP:O	2.38	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:LYS:NZ	1:A:164:LYS:CB	2.84	0.41
1:A:261[A]:LEU:HD11	1:A:277:PHE:CD2	2.55	0.40
1:A:568:ARG:HB2	1:A:569:PRO:HD3	2.03	0.40
1:A:296:GLU:OE2	3:A:702:X8Z:H11C	2.21	0.40
1:A:29:PHE:HB3	6:A:709:GOL:H32	2.03	0.40
1:A:271:GLU:HG3	1:A:299:HIS:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:481:ASP:OD1	4:A:703:YB:YB[1_545]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	642/611 (105%)	628 (98%)	14 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	576/543 (106%)	550 (96%)	26 (4%)	24 17

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	VAL
1	A	31[A]	ARG
1	A	31[B]	ARG
1	A	34	LEU
1	A	75	LEU
1	A	92	ILE
1	A	144	LEU
1	A	154	LEU
1	A	164	LYS
1	A	178	THR
1	A	182	GLU
1	A	192	ILE
1	A	242	LYS
1	A	308	ASN
1	A	332	LEU
1	A	352[A]	SER
1	A	352[B]	SER
1	A	380[A]	SER
1	A	380[B]	SER
1	A	396	LEU
1	A	401[A]	GLU
1	A	401[B]	GLU
1	A	407	LEU
1	A	530[A]	ASN
1	A	530[B]	ASN
1	A	587	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	116	GLN
1	A	213	GLN
1	A	272	ASN
1	A	341	ASN
1	A	350	GLN
1	A	360	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	440	ASN
1	A	441	GLN
1	A	466	ASN
1	A	521	GLN
1	A	525	ASN
1	A	527	ASN
1	A	581	HIS
1	A	591	HIS
1	A	596	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	709	-	5,5,5	0.68	0	5,5,5	0.81	0
3	X8Z	A	702	2	14,14,14	2.00	3 (21%)	18,19,19	1.29	4 (22%)
5	ACY	A	708[B]	4	3,3,3	0.60	0	3,3,3	1.20	0
6	GOL	A	710	-	5,5,5	0.26	0	5,5,5	0.84	0
5	ACY	A	708[A]	4	3,3,3	0.61	0	3,3,3	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	710	-	-	2/4/4/4	-
6	GOL	A	709	-	-	0/4/4/4	-
3	X8Z	A	702	2	-	2/14/24/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	X8Z	C8-C9	-5.36	1.42	1.52
3	A	702	X8Z	C8-N	2.96	1.52	1.47
3	A	702	X8Z	C2-C4	2.34	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	X8Z	C9-C8-N	-2.50	107.53	112.54
3	A	702	X8Z	C7-C8-N	2.37	106.49	103.02
3	A	702	X8Z	C2-C1-S	-2.20	111.19	114.04
3	A	702	X8Z	C5-N-C8	-2.12	108.69	112.01

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	X8Z	S-C1-C2-C4
3	A	702	X8Z	S-C1-C2-C3
6	A	710	GOL	O1-C1-C2-C3
6	A	710	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	709	GOL	1	0
3	A	702	X8Z	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	607/611 (99%)	-0.57	0 100 100	5, 14, 26, 40	37 (6%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	YB	A	706	1/1	0.88	0.11	84,84,84,84	1
6	GOL	A	710	6/6	0.90	0.09	26,35,36,36	0
6	GOL	A	709	6/6	0.92	0.08	26,31,32,33	0
4	YB	A	707	1/1	0.93	0.11	84,84,84,84	1
3	X8Z	A	702	14/14	0.95	0.08	13,22,25,25	0
5	ACY	A	708[B]	4/4	0.97	0.07	13,13,14,16	4
4	YB	A	705	1/1	0.97	0.07	62,62,62,62	1
5	ACY	A	708[A]	4/4	0.97	0.07	5,6,7,8	4
4	YB	A	703	1/1	1.00	0.01	9,9,9,9	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	YB	A	704	1/1	1.00	0.01	10,10,10,10	1
2	ZN	A	701	1/1	1.00	0.02	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.