



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:35 AM UTC

PDB ID : 4DQP / pdb_00004dqp
Title : Ternary complex of Bacillus DNA Polymerase I Large Fragment, DNA duplex, and ddCTP (paired with dG of template)
Authors : Wang, W.; Beese, L.S.
Deposited on : 2012-02-16
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	6% 89% 8%
1	D	592	% 90% 7%
2	B	9	67% 33%
2	E	9	89% 11%
3	C	13	15% 92% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 92%8%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 21755 atoms, of which 10084 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	576	Total	C	H	N	O	S	0	17	0
			9444	2979	4762	810	876	17			
1	D	579	Total	C	H	N	O	S	0	26	0
			9574	3014	4830	824	889	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ALA	ASP	engineered mutation	UNP Q5KWC1
A	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1
D	598	ALA	ASP	engineered mutation	UNP Q5KWC1
D	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1


- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	P	0	0	0
			274	85	100	29	52	8			
2	E	9	Total	C	H	N	O	P	0	0	0
			274	85	100	29	52	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*C*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	12	Total	C	H	N	O	P	0	0	0
			385	119	134	52	69	11			
3	F	12	Total	C	H	N	O	P	0	0	0
			385	119	134	52	69	11			

- Molecule 4 is 2',3'-DIDEOXYCYTIDINE 5'-TRIPHOSPHATE (CCD ID: DCT) (formula:


$$\text{C}_9\text{H}_{16}\text{N}_3\text{O}_{12}\text{P}_3).$$


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 35	C 9	H 8	N 3	O 12	P 3	0	0
4	A	1	Total 35	C 9	H 8	N 3	O 12	P 3	0	0
4	D	1	Total 35	C 9	H 8	N 3	O 12	P 3	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	D	1	Total Mg 1 1	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	388	Total	O	0	0
			388	388		
7	D	710	Total	O	0	0
			710	710		
7	B	32	Total	O	0	0
			32	32		
7	C	57	Total	O	0	0
			57	57		
7	E	37	Total	O	0	0
			37	37		
7	F	78	Total	O	0	0
			78	78		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase




- Molecule 1: DNA polymerase




- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3')




- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3')



- Molecule 3: DNA (5'-D(*C*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain C:  15% 92% 8%



- Molecule 3: DNA (5'-D(*C*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain F:  92% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.22Å 109.34Å 149.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.45 – 1.74 29.45 – 1.74	Depositor EDS
% Data completeness (in resolution range)	87.0 (29.45-1.74) 87.0 (29.45-1.74)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 1.74Å)	Xtriage
Refinement program	PHENIX dev_1026	Depositor
R, R_{free}	0.167 , 0.195 0.168 , 0.196	Depositor DCC
R_{free} test set	6153 reflections (4.46%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.139	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21755	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, DCT, MG, DOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/4824	0.76	0/6520
1	D	0.77	1/4944 (0.0%)	0.85	0/6681
2	B	0.48	0/173	1.30	1/264 (0.4%)
2	E	0.59	0/173	1.28	0/264
3	C	0.50	0/283	1.01	0/437
3	F	0.59	0/283	1.18	0/437
All	All	0.66	1/10680 (0.0%)	0.85	1/14603 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	620	GLU	C-O	8.54	1.27	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	26	DC	N1-C1'-C2'	-7.11	102.83	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4682	4762	4687	33	0
1	D	4744	4830	4704	42	0
2	B	174	100	103	1	0
2	E	174	100	103	0	0
3	C	251	134	136	0	0
3	F	251	134	136	0	0
4	A	54	16	24	0	0
4	D	27	8	12	0	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	D	10	0	0	0	0
7	A	388	0	0	12	0
7	B	32	0	0	0	0
7	C	57	0	0	0	0
7	D	710	0	0	19	0
7	E	37	0	0	0	0
7	F	78	0	0	0	0
All	All	11671	10084	9905	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:423:ARG:NE	7:D:1668:HOH:O	1.97	0.94
1:D:629:ARG:NE	7:D:1666:HOH:O	2.08	0.82
1:D:629:ARG:NH2	7:D:1666:HOH:O	2.11	0.81
1:D:435:ARG:NH1	7:D:1356:HOH:O	2.15	0.80
1:A:445:GLU:OE1	7:A:1368:HOH:O	1.99	0.79
1:D:629:ARG:NH1	7:D:1254:HOH:O	2.15	0.78
1:A:435:ARG:NH1	7:A:1157:HOH:O	2.18	0.73
1:D:459:ARG:NH2	7:D:1688:HOH:O	2.14	0.71
1:D:310:GLU:OE2	7:D:1465:HOH:O	2.10	0.68
1:D:589:GLU:OE1	7:D:1694:HOH:O	2.11	0.68
1:D:602[B]:VAL:HG21	1:D:621:PRO:HG3	1.75	0.66
1:A:654:TYR:HB3	1:A:657:ILE:HB	1.78	0.65
1:A:541:LYS:CE	7:A:1362:HOH:O	2.45	0.64
1:A:692:VAL:HG21	1:A:701:MET:HE1	1.81	0.63
1:D:692:VAL:HB	1:D:696:GLU:HB2	1.83	0.60
1:D:876:LYS:O	7:D:1621:HOH:O	2.16	0.60
1:D:629:ARG:CZ	7:D:1666:HOH:O	2.36	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:664:HIS:O	1:A:859:ARG:NH1	2.35	0.60
1:D:690:PHE:HB2	1:D:701:MET:HE3	1.84	0.59
1:A:846:ARG:CZ	7:A:1387:HOH:O	2.51	0.58
1:D:690:PHE:CB	1:D:701:MET:HE3	2.34	0.58
1:A:552[B]:THR:HG22	2:B:24:DG:OP1	2.05	0.57
1:D:323:VAL:O	1:D:435:ARG:NH2	2.37	0.57
1:D:726:ASN:HB3	7:D:1347:HOH:O	2.04	0.57
1:D:426:GLU:OE2	1:D:431:LYS:HE2	2.05	0.56
1:D:692:VAL:HG21	1:D:701:MET:HE1	1.88	0.56
1:D:466:ARG:NH1	7:D:1237:HOH:O	2.39	0.55
1:A:583[B]:LEU:HD11	1:A:636:ILE:CG1	2.36	0.55
1:A:703:ARG:HG2	7:A:1351:HOH:O	2.06	0.55
1:A:541:LYS:HE3	7:A:1362:HOH:O	2.04	0.55
1:A:846:ARG:NH1	7:A:1387:HOH:O	2.40	0.54
1:D:744:PRO:O	1:D:748[A]:ARG:HG3	2.08	0.54
1:D:690:PHE:CG	1:D:701:MET:HE3	2.43	0.54
1:A:541:LYS:HE2	7:A:1362:HOH:O	2.09	0.53
1:D:325:GLU:CD	7:D:1703:HOH:O	2.51	0.53
1:A:459:ARG:HB3	1:A:460:PRO:HD3	1.91	0.53
1:D:463:ASP:OD2	1:D:466:ARG:NH2	2.42	0.53
1:D:561:LEU:HB3	1:D:571[A]:VAL:HG12	1.91	0.52
1:D:408:ASP:OD1	1:D:408:ASP:N	2.42	0.52
1:A:565:ALA:N	1:A:566:PRO:HD2	2.25	0.51
1:A:738:ARG:O	1:A:741:GLU:HG2	2.10	0.51
1:D:620:GLU:HB3	7:D:1471:HOH:O	2.10	0.50
1:D:459:ARG:NH2	7:D:1361:HOH:O	2.40	0.50
1:A:565:ALA:N	1:A:566:PRO:CD	2.74	0.50
1:A:477[A]:LEU:HD12	1:A:809:ILE:HD12	1.95	0.48
1:A:848:VAL:HB	1:A:849:PRO:HD3	1.96	0.47
1:A:472:ARG:NH1	7:A:1236:HOH:O	2.44	0.47
1:D:690:PHE:CD1	1:D:701:MET:HE3	2.49	0.47
1:A:519:TYR:CD2	1:A:525:GLU:HG2	2.50	0.47
1:A:576:HIS:HD2	7:A:1101:HOH:O	1.97	0.47
1:D:426:GLU:OE1	7:D:1541:HOH:O	2.21	0.47
1:A:334:GLY:HA2	1:A:348:PRO:HD3	1.97	0.46
1:D:764:THR:HA	1:D:769:ARG:O	2.16	0.45
1:D:565:ALA:N	1:D:566:PRO:CD	2.80	0.45
1:A:583[B]:LEU:HD11	1:A:636:ILE:HG13	1.98	0.44
1:D:841:MET:HE2	1:D:844:LEU:HD12	1.99	0.44
1:D:315:LYS:HG3	7:D:1632:HOH:O	2.15	0.44
1:D:738:ARG:HD3	7:D:1419:HOH:O	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:689:ILE:HG13	1:A:690:PHE:CD2	2.52	0.44
1:A:779:ARG:NH2	7:A:1138:HOH:O	2.51	0.43
1:D:754:VAL:HG12	1:D:758:LYS:HE2	2.00	0.43
1:A:672:MET:O	1:A:675:PHE:C	2.62	0.43
1:A:719:TYR:O	1:A:723:GLN:HG2	2.18	0.42
1:D:648:LEU:HD12	1:D:841:MET:HG3	2.00	0.42
1:D:654:TYR:HB3	1:D:657:ILE:HB	2.01	0.42
1:D:534:LEU:HD11	1:D:574:ILE:HD13	2.01	0.42
1:A:576:HIS:CD2	7:A:1101:HOH:O	2.72	0.42
1:A:329:ASP:H	1:A:382:TRP:CD1	2.38	0.41
1:D:836:ALA:HB3	1:D:841:MET:CE	2.50	0.41
1:D:565:ALA:N	1:D:566:PRO:HD2	2.36	0.41
1:A:744:PRO:O	1:A:748:ARG:HG3	2.21	0.41
1:D:564:LEU:C	1:D:566:PRO:HD2	2.46	0.41
1:D:569:GLU:HA	7:D:1466:HOH:O	2.20	0.40
1:A:534:LEU:HD11	1:A:574:ILE:HD13	2.04	0.40
1:A:349:GLU:CD	1:A:349:GLU:H	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/592 (100%)	572 (97%)	16 (3%)	1 (0%)	43	29
1	D	603/592 (102%)	589 (98%)	13 (2%)	1 (0%)	43	29
All	All	1192/1184 (101%)	1161 (97%)	29 (2%)	2 (0%)	43	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	628	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	628	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	507/507 (100%)	503 (99%)	4 (1%)	73	64
1	D	521/507 (103%)	517 (99%)	4 (1%)	73	64
All	All	1028/1014 (101%)	1020 (99%)	8 (1%)	73	64

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	632	GLU
1	A	728	SER
1	A	839	GLU
1	A	844	LEU
1	D	415	LYS
1	D	511	LEU
1	D	561	LEU
1	D	726	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	356	GLN
1	A	418	GLN
1	A	543	GLN
1	A	821	GLN
1	D	341	HIS
1	D	625	ASN
1	D	726	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	DOC	E	29	2,3	16,19,20	0.68	0	20,26,29	0.89	2 (10%)
2	DOC	B	29	2,3	16,19,20	0.48	0	20,26,29	0.81	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	E	29	2,3	-	0/7/18/19	0/2/2/2
2	DOC	B	29	2,3	-	0/7/18/19	0/2/2/2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	29	DOC	C4'-O4'-C1'	-2.62	107.33	109.81
2	E	29	DOC	O4'-C1'-C2'	2.18	109.00	106.41
2	E	29	DOC	C4'-O4'-C1'	-2.15	107.78	109.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	D	903	-	4,4,4	0.19	0	6,6,6	0.65	0
4	DCT	A	901	5	27,28,28	1.06	3 (11%)	37,43,43	1.64	4 (10%)
4	DCT	A	902	-	27,28,28	1.23	4 (14%)	37,43,43	1.50	5 (13%)
6	SO4	D	904	-	4,4,4	0.29	0	6,6,6	0.15	0
4	DCT	D	901	5	27,28,28	1.00	2 (7%)	37,43,43	1.45	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCT	D	901	5	-	4/22/31/31	0/2/2/2
4	DCT	A	902	-	-	6/22/31/31	0/2/2/2
4	DCT	A	901	5	-	5/22/31/31	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	DCT	PA-O3A	2.80	1.62	1.59
4	A	902	DCT	PB-O3B	2.67	1.62	1.59
4	D	901	DCT	C4-N3	2.60	1.39	1.34
4	A	902	DCT	PB-O3A	2.59	1.62	1.59
4	A	901	DCT	PB-O3A	2.41	1.62	1.59
4	A	901	DCT	C4-N3	2.17	1.38	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	901	DCT	PA-O3A	2.06	1.61	1.59
4	A	901	DCT	C6-C5	2.03	1.39	1.35
4	A	902	DCT	C4-N3	2.01	1.38	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	901	DCT	C2'-C1'-N1	-6.06	100.91	112.40
4	A	902	DCT	C2'-C1'-N1	-4.73	103.43	112.40
4	A	901	DCT	O4'-C1'-N1	4.00	114.96	107.86
4	D	901	DCT	C4'-O4'-C1'	3.94	113.53	109.81
4	A	902	DCT	O4'-C4'-C5'	2.98	114.72	109.34
4	A	902	DCT	O4'-C1'-N1	2.98	113.15	107.86
4	A	902	DCT	O2A-PA-O3A	2.72	114.61	107.27
4	A	901	DCT	O2B-PB-O3B	2.71	114.59	107.27
4	A	902	DCT	C3'-C2'-C1'	2.34	105.57	102.87
4	D	901	DCT	C4-N3-C2	-2.27	116.70	120.26
4	D	901	DCT	O3G-PG-O2G	2.23	116.16	107.80
4	A	901	DCT	C4'-O4'-C1'	2.22	111.90	109.81
4	D	901	DCT	C5-C6-N1	-2.04	118.53	121.84

There are no chirality outliers.


All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	902	DCT	C3'-C4'-C5'-O5'
4	A	902	DCT	O4'-C4'-C5'-O5'
4	A	901	DCT	PG-O3B-PB-O1B
4	A	902	DCT	PA-O3A-PB-O3B
4	A	902	DCT	PG-O3B-PB-O2B
4	D	901	DCT	PG-O3B-PB-O1B
4	A	901	DCT	PB-O3B-PG-O1G
4	A	901	DCT	PB-O3A-PA-O2A
4	D	901	DCT	PG-O3B-PB-O2B
4	D	901	DCT	PB-O3B-PG-O1G
4	A	901	DCT	PB-O3A-PA-O1A
4	A	901	DCT	PG-O3B-PB-O2B
4	D	901	DCT	PB-O3A-PA-O2A
4	A	902	DCT	PA-O3A-PB-O1B
4	A	902	DCT	PA-O3A-PB-O2B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/592 (97%)	0.45	35 (6%) 27 33	12, 43, 86, 115	9 (1%)
1	D	579/592 (97%)	-0.45	5 (0%) 81 87	6, 21, 45, 64	13 (2%)
2	B	8/9 (88%)	-0.35	0 100 100	23, 27, 48, 62	0
2	E	8/9 (88%)	-0.72	0 100 100	15, 21, 35, 51	0
3	C	12/13 (92%)	-0.02	2 (16%) 4 6	21, 29, 80, 107	0
3	F	12/13 (92%)	-0.55	0 100 100	11, 19, 58, 82	0
All	All	1195/1228 (97%)	-0.01	42 (3%) 47 56	6, 32, 74, 115	22 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	686	ALA	4.4
1	A	740	PHE	4.0
1	A	675	PHE	3.9
1	A	657	ILE	3.8
1	A	689	ILE	3.8
1	A	697	VAL	3.7
1	A	297	ALA	3.6
1	A	705	ALA	3.5
1	A	724[A]	ASN	3.5
1	A	681	ILE	3.4
1	A	674	ALA	3.4
1	A	690	PHE	3.3
1	D	748[A]	ARG	3.2
1	A	693	SER	3.0
1	A	713	VAL	3.0
1	A	692	VAL	2.9
1	A	742	SER	2.9
1	A	743	PHE	2.8
1	A	685	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	699	PRO	2.6
1	D	513[A]	THR	2.6
1	A	684	LYS	2.6
1	A	477[A]	LEU	2.5
3	C	1	DA	2.5
1	A	714	TYR	2.5
1	A	735	PHE	2.4
1	A	739	TYR	2.4
1	D	703	ARG	2.3
1	A	688	ASP	2.3
1	D	876	LYS	2.3
1	A	672	MET	2.3
1	A	860	VAL	2.3
3	C	2	DT	2.3
1	A	845	CYS	2.2
1	A	708	VAL	2.2
1	A	763[A]	VAL	2.2
1	A	683	THR	2.1
1	A	781	PHE	2.1
1	A	435	ARG	2.1
1	A	694	GLU	2.1
1	D	727	ILE	2.0
1	A	758	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	DOC	B	29	18/19	0.97	0.06	24,29,33,39	0
2	DOC	E	29	18/19	0.99	0.04	11,14,18,21	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DCT	A	902	27/27	0.87	0.10	30,39,87,90	0
6	SO4	D	904	5/5	0.90	0.09	57,59,65,67	0
5	MG	A	903	1/1	0.91	0.09	40,40,40,40	0
4	DCT	A	901	27/27	0.93	0.08	20,31,50,52	0
6	SO4	D	903	5/5	0.96	0.08	25,28,45,49	0
5	MG	D	902	1/1	0.98	0.03	18,18,18,18	0
4	DCT	D	901	27/27	0.99	0.03	8,13,19,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around DCT A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DCT D 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.