



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:20 PM UTC

PDB ID : 2DRC / pdb_00002drc
Title : INVESTIGATION OF THE FUNCTIONAL ROLE OF TRYPTOPHAN-22 IN ESCHERICHIA COLI DIHYDROFOLATE REDUCTASE BY SITE-DIRECTED MUTAGENESIS
Authors : Brown, K.A.; Kraut, J.
Deposited on : 1992-06-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	159	
1	B	159	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MTX	B	161	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2619 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIHYDROFOLATE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	1243	791	210	235	7	0	0	0
1	B	159	1259	801	215	236	7	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	PHE	TRP	conflict	UNP P0ABQ4
A	37	ASP	ASN	conflict	UNP P0ABQ4
B	22	PHE	TRP	conflict	UNP P0ABQ4
B	37	ASP	ASN	conflict	UNP P0ABQ4

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is METHOTREXATE (CCD ID: MTX) (formula: C₂₀H₂₂N₈O₅).


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: DIHYDROFOLATE REDUCTASE

Chain A:  67% 23% 8%



- Molecule 1: DIHYDROFOLATE REDUCTASE

Chain B:  75% 23%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	92.99Å 92.99Å 73.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 1.90	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.161 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2619	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA, MTX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.42	3/1275 (0.2%)	1.97	40/1733 (2.3%)
1	B	1.42	1/1292 (0.1%)	2.04	30/1755 (1.7%)
All	All	1.42	4/2567 (0.2%)	2.01	70/3488 (2.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	158	ARG	CA-C	-6.08	1.45	1.52
1	B	113	THR	CA-C	5.31	1.59	1.52
1	A	22	PHE	N-CA	-5.23	1.39	1.45
1	A	87	ASP	CG-OD2	5.11	1.35	1.25

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	SER	N-CA-C	9.34	122.45	111.71
1	A	87	ASP	CA-CB-CG	-8.47	104.13	112.60
1	A	132	ASP	N-CA-C	-8.18	100.83	113.02
1	B	120	GLU	CA-C-N	-8.08	112.70	122.15
1	B	120	GLU	C-N-CA	-8.08	112.70	122.15
1	A	158	ARG	O-C-N	8.00	133.22	122.82
1	B	45	HIS	CA-CB-CG	-7.77	106.03	113.80
1	A	69	ASP	N-CA-C	7.50	121.91	109.46
1	A	45	HIS	CA-CB-CG	-7.34	106.46	113.80
1	B	87	ASP	CA-CB-CG	-7.11	105.49	112.60
1	B	101	GLU	CB-CA-C	-6.67	99.71	110.79
1	B	94	ILE	N-CA-C	6.61	118.44	112.17
1	B	111	TYR	CA-C-O	6.54	127.27	120.40
1	A	113	THR	CA-C-O	-6.54	113.54	120.40
1	B	148	SER	N-CA-C	6.50	120.47	112.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	79	ASP	O-C-N	6.39	128.65	122.07
1	A	125	PHE	N-CA-C	-6.34	101.33	110.08
1	B	12	ARG	NE-CZ-NH1	6.32	127.81	121.50
1	A	94	ILE	N-CA-C	6.30	118.16	112.17
1	B	46	THR	N-CA-C	-6.28	104.52	111.36
1	A	107	ALA	CA-C-O	6.21	128.11	121.16
1	A	126	PRO	N-CA-CB	6.09	108.43	103.32
1	A	120	GLU	CA-C-N	-6.05	111.13	121.17
1	A	120	GLU	C-N-CA	-6.05	111.13	121.17
1	B	145	ALA	CA-C-N	-6.03	112.76	122.23
1	B	145	ALA	C-N-CA	-6.03	112.76	122.23
1	A	68	THR	CA-C-N	6.01	130.66	122.19
1	A	68	THR	C-N-CA	6.01	130.66	122.19
1	A	12	ARG	CG-CD-NE	5.91	125.00	112.00
1	A	114	HIS	CA-CB-CG	-5.86	107.94	113.80
1	B	138	SER	CA-CB-OG	-5.86	99.38	111.10
1	A	118	GLU	CB-CG-CD	-5.85	102.65	112.60
1	B	143	ALA	N-CA-C	-5.85	102.39	110.35
1	B	82	ILE	N-CA-C	5.82	116.56	110.62
1	B	27	ASP	CB-CG-OD1	5.81	131.77	118.40
1	A	64	SER	CA-CB-OG	-5.66	99.78	111.10
1	B	114	HIS	CA-C-N	-5.62	116.22	123.19
1	B	114	HIS	C-N-CA	-5.62	116.22	123.19
1	A	61	ILE	CA-CB-CG1	5.60	119.92	110.40
1	A	95	GLY	CA-C-N	5.56	131.71	121.70
1	A	95	GLY	C-N-CA	5.56	131.71	121.70
1	A	65	GLN	CB-CA-C	5.53	121.07	110.17
1	A	80	GLU	CB-CG-CD	-5.50	103.25	112.60
1	B	12	ARG	CD-NE-CZ	5.50	132.10	124.40
1	A	54	LEU	N-CA-C	-5.48	102.74	109.65
1	A	79	ASP	CA-C-O	-5.47	115.08	120.82
1	B	24	LEU	N-CA-C	5.44	118.52	108.94
1	A	127	ASP	CA-CB-CG	-5.43	107.17	112.60
1	B	153	PHE	CA-CB-CG	-5.43	108.37	113.80
1	A	130	PRO	N-CA-CB	5.42	108.94	103.25
1	A	23	ASN	CA-CB-CG	-5.37	107.23	112.60
1	B	8	LEU	CB-CA-C	-5.36	99.90	109.71
1	B	136	VAL	O-C-N	-5.32	117.32	122.16
1	A	69	ASP	N-CA-CB	-5.24	101.51	110.16
1	A	107	ALA	N-CA-C	5.23	118.03	110.28
1	B	140	PHE	CA-CB-CG	-5.20	108.60	113.80
1	A	43	GLY	N-CA-C	-5.19	104.66	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	12	ARG	NE-CZ-NH2	-5.15	114.56	119.20
1	A	92	MET	CA-C-N	-5.14	116.32	122.90
1	A	92	MET	C-N-CA	-5.14	116.32	122.90
1	B	12	ARG	CG-CD-NE	-5.12	100.73	112.00
1	B	111	TYR	CB-CA-C	-5.12	102.50	110.74
1	B	111	TYR	N-CA-C	-5.11	99.08	108.02
1	A	39	PRO	N-CA-CB	5.10	108.10	103.15
1	B	68	THR	O-C-N	-5.07	116.10	122.24
1	A	22	PHE	CB-CA-C	5.06	121.33	110.45
1	A	102	GLN	CA-C-N	-5.05	113.53	122.26
1	A	102	GLN	C-N-CA	-5.05	113.53	122.26
1	B	61	ILE	CA-CB-CG1	5.01	118.92	110.40
1	A	12	ARG	CD-NE-CZ	5.01	131.41	124.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1243	0	1177	22	0
1	B	1259	0	1216	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	33	0	20	1	0
3	B	33	0	21	1	0
4	B	1	0	0	0	0
5	A	33	0	0	2	0
5	B	15	0	0	0	0
All	All	2619	0	2434	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:TYR:O	1:A:130:PRO:HD3	1.87	0.75
1:A:67:GLY:HA2	1:A:74:TRP:CE2	2.28	0.68
1:A:10:VAL:HG13	1:A:117:ALA:O	1.95	0.66
1:A:67:GLY:HA2	1:A:74:TRP:CD2	2.32	0.65
1:B:20:MET:HE3	1:B:22:PHE:HE2	1.62	0.64
1:B:104:LEU:HB3	1:B:105:PRO:HD3	1.84	0.58
1:B:20:MET:HE3	1:B:22:PHE:CE2	2.39	0.57
1:A:33:ARG:HG3	5:A:418:HOH:O	2.09	0.53
1:B:64:SER:CB	1:B:65:GLN:NE2	2.75	0.50
1:A:104:LEU:N	1:A:105:PRO:CD	2.76	0.49
1:A:107:ALA:O	1:A:158:ARG:HD2	2.12	0.49
1:A:98:ARG:HB3	1:A:98:ARG:CZ	2.43	0.48
1:A:130:PRO:O	1:A:133:TRP:O	2.31	0.48
1:A:107:ALA:O	1:A:158:ARG:NH1	2.44	0.48
1:B:65:GLN:HB3	1:B:66:PRO:HD2	1.96	0.47
1:B:97:GLY:O	1:B:101:GLU:HG3	2.15	0.47
1:A:12:ARG:HH21	1:A:127:ASP:N	2.13	0.47
1:A:104:LEU:N	1:A:105:PRO:HD2	2.30	0.46
1:B:137:PHE:O	1:B:154:GLU:HA	2.15	0.46
1:A:25:PRO:HD2	1:A:147:ASN:OD1	2.16	0.45
1:B:36:LEU:HD12	1:B:36:LEU:HA	1.84	0.43
3:B:161:MTX:H13	3:B:161:MTX:HM1	1.75	0.43
1:A:129:GLU:O	1:A:131:ASP:N	2.52	0.43
1:A:75:VAL:HB	1:A:80:GLU:HB3	1.99	0.43
3:A:161:MTX:H13	3:A:161:MTX:HM1	1.69	0.42
1:B:129:GLU:HB3	1:B:132:ASP:CG	2.44	0.42
1:A:12:ARG:HH21	1:A:127:ASP:CA	2.33	0.41
1:A:33:ARG:HD3	5:A:418:HOH:O	2.20	0.41
1:A:103:PHE:C	1:A:105:PRO:HD2	2.45	0.41
1:A:44:ARG:NH2	1:A:65:GLN:HB2	2.36	0.41
1:A:59:ASN:HD22	1:A:59:ASN:N	2.19	0.41
1:B:65:GLN:HB3	1:B:66:PRO:CD	2.51	0.41
1:A:108:GLN:HE21	1:A:108:GLN:HB2	1.72	0.40
1:A:129:GLU:C	1:A:131:ASP:H	2.30	0.40
1:B:64:SER:HB3	1:B:65:GLN:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/159 (99%)	150 (96%)	6 (4%)	1 (1%)	21	13
1	B	157/159 (99%)	153 (98%)	4 (2%)	0	100	100
All	All	314/318 (99%)	303 (96%)	10 (3%)	1 (0%)	36	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/136 (95%)	118 (92%)	11 (8%)	10	4
1	B	134/136 (98%)	127 (95%)	7 (5%)	21	13
All	All	263/272 (97%)	245 (93%)	18 (7%)	14	7

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	10	VAL
1	A	33	ARG
1	A	59	ASN
1	A	61	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	64	SER
1	A	80	GLU
1	A	87	ASP
1	A	104	LEU
1	A	132	ASP
1	A	136	VAL
1	B	1	MET
1	B	52	ARG
1	B	59	ASN
1	B	61	ILE
1	B	64	SER
1	B	87	ASP
1	B	156	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	108	GLN
1	B	65	GLN
1	B	108	GLN
1	B	124	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MTX	B	161	-	35,35,35	4.66	21 (60%)	47,49,49	3.33	22 (46%)
3	MTX	A	161	-	35,35,35	4.17	22 (62%)	47,49,49	3.20	19 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MTX	B	161	-	-	3/25/25/25	0/3/3/3
3	MTX	A	161	-	-	2/25/25/25	0/3/3/3

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	161	MTX	C13-C12	-12.99	1.17	1.38
3	B	161	MTX	C4A-C8A	-12.92	1.19	1.40
3	A	161	MTX	OE1-CD	9.62	1.53	1.22
3	A	161	MTX	CG-CD	9.33	1.72	1.50
3	B	161	MTX	C9-C6	7.37	1.63	1.51
3	B	161	MTX	C7-N8	-7.07	1.19	1.31
3	A	161	MTX	C16-C15	-7.01	1.27	1.38
3	A	161	MTX	C16-C11	6.79	1.49	1.39
3	A	161	MTX	CM-N10	6.54	1.56	1.46
3	B	161	MTX	C7-C6	-6.49	1.28	1.39
3	B	161	MTX	C16-C15	-6.46	1.28	1.38
3	A	161	MTX	C15-C14	5.75	1.50	1.39
3	B	161	MTX	O-C	5.66	1.36	1.23
3	A	161	MTX	C13-C12	5.56	1.47	1.38
3	B	161	MTX	C8A-N1	5.54	1.47	1.36
3	B	161	MTX	C8A-N8	5.50	1.45	1.37
3	B	161	MTX	OE1-CD	5.50	1.40	1.22
3	A	161	MTX	C4A-N5	-5.04	1.27	1.37
3	A	161	MTX	C9-N10	4.97	1.56	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	161	MTX	C4A-C8A	-4.87	1.32	1.40
3	A	161	MTX	C7-N8	4.82	1.39	1.31
3	B	161	MTX	C2-N3	4.40	1.43	1.35
3	A	161	MTX	C7-C6	4.37	1.47	1.39
3	A	161	MTX	O1-CT	4.30	1.34	1.22
3	A	161	MTX	OE2-CD	-3.86	1.18	1.30
3	B	161	MTX	C2-NA2	-3.36	1.27	1.33
3	A	161	MTX	C2-NA2	-3.23	1.27	1.33
3	A	161	MTX	C12-C11	-3.14	1.34	1.39
3	B	161	MTX	OE2-CD	3.07	1.41	1.30
3	B	161	MTX	C9-N10	3.06	1.52	1.46
3	A	161	MTX	C4-N3	3.03	1.39	1.33
3	A	161	MTX	CA-N	2.98	1.52	1.45
3	B	161	MTX	C11-C	-2.85	1.44	1.50
3	B	161	MTX	C12-C11	-2.84	1.35	1.39
3	B	161	MTX	C4-NA4	2.75	1.43	1.34
3	A	161	MTX	C14-N10	2.73	1.46	1.38
3	B	161	MTX	O1-CT	2.57	1.29	1.22
3	B	161	MTX	CA-N	-2.45	1.40	1.45
3	A	161	MTX	C13-C14	-2.45	1.34	1.39
3	B	161	MTX	CB-CG	-2.29	1.46	1.52
3	B	161	MTX	C4A-N5	2.17	1.41	1.37
3	A	161	MTX	O-C	2.13	1.28	1.23
3	A	161	MTX	O2-CT	-2.09	1.24	1.30

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	161	MTX	N1-C2-N3	-11.53	112.56	127.21
3	A	161	MTX	C4A-C4-N3	-10.51	108.40	120.84
3	A	161	MTX	C4-C4A-N5	-7.20	114.78	120.33
3	B	161	MTX	C13-C14-N10	-6.92	111.93	121.59
3	A	161	MTX	N1-C2-N3	-6.20	119.33	127.21
3	A	161	MTX	CT-CA-N	-5.99	96.68	110.57
3	A	161	MTX	N8-C8A-N1	-5.82	109.42	115.77
3	B	161	MTX	C16-C15-C14	-5.66	113.13	120.30
3	B	161	MTX	CB-CA-CT	-5.58	97.11	110.35
3	B	161	MTX	C4-C4A-N5	-5.51	116.08	120.33
3	B	161	MTX	CM-N10-C14	5.47	128.65	119.59
3	B	161	MTX	C4A-C4-N3	-5.45	114.39	120.84
3	B	161	MTX	NA2-C2-N3	5.39	125.31	117.22
3	A	161	MTX	C6-C7-N8	-5.26	118.09	123.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	161	MTX	C9-N10-C14	-4.92	114.21	120.17
3	A	161	MTX	C2-N3-C4	4.80	130.42	116.72
3	B	161	MTX	C2-N1-C8A	4.17	119.98	115.48
3	A	161	MTX	C13-C14-N10	-4.07	115.91	121.59
3	A	161	MTX	C4A-C4-NA4	4.03	126.45	120.31
3	A	161	MTX	CM-N10-C9	3.99	125.49	115.11
3	B	161	MTX	OE1-CD-CG	-3.80	111.03	123.09
3	B	161	MTX	C9-N10-C14	-3.61	115.79	120.17
3	B	161	MTX	C2-N3-C4	3.60	127.00	116.72
3	B	161	MTX	C9-C6-N5	-3.47	111.47	117.09
3	B	161	MTX	C6-N5-C4A	-3.17	113.69	118.17
3	A	161	MTX	NA4-C4-N3	2.99	125.14	117.11
3	B	161	MTX	C11-C-N	2.93	122.46	117.04
3	A	161	MTX	OE2-CD-CG	2.92	123.21	114.00
3	A	161	MTX	NA2-C2-N3	2.86	121.52	117.22
3	B	161	MTX	C7-C6-N5	2.78	122.68	120.87
3	B	161	MTX	NA2-C2-N1	2.78	122.13	117.79
3	B	161	MTX	OE2-CD-CG	2.63	122.31	114.00
3	A	161	MTX	C12-C13-C14	-2.62	116.98	120.30
3	B	161	MTX	O-C-N	-2.52	117.67	122.47
3	A	161	MTX	OE1-CD-CG	-2.50	115.16	123.09
3	A	161	MTX	C4A-C8A-N1	2.46	125.56	121.74
3	A	161	MTX	CG-CB-CA	2.46	117.69	113.16
3	B	161	MTX	C9-C6-C7	2.43	125.81	121.38
3	B	161	MTX	NA4-C4-N3	2.41	123.59	117.11
3	B	161	MTX	C15-C14-C13	2.37	123.70	119.18
3	A	161	MTX	C13-C12-C11	2.14	123.09	120.80

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	161	MTX	C6-C9-N10-CM
3	B	161	MTX	C6-C9-N10-CM
3	A	161	MTX	CA-CB-CG-CD
3	B	161	MTX	OE1-CD-CG-CB
3	B	161	MTX	OE2-CD-CG-CB


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	161	MTX	1	0
3	A	161	MTX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.