



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 03:47 AM UTC

PDB ID : 4DRB / pdb_00004drb
Title : The crystal structure of FANCM bound MHF complex
Authors : Tao, Y.; Niu, L.; Teng, M.
Deposited on : 2012-02-17
Resolution : 2.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2053 (2.66-2.62)
Clashscore	190562	2097 (2.66-2.62)
Ramachandran outliers	187476	2066 (2.66-2.62)
Sidechain outliers	187428	2066 (2.66-2.62)
RSRZ outliers	180081	2052 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	 2% 52% 24% 5% 18%
1	B	120	 % 59% 26% • 14%
1	D	120	 2% 59% 21% • 19%
1	E	120	 4% 58% 22% • 19%
1	G	120	 2% 72% 10% 5% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	120	<p>2% 61% 18% 22%</p>
2	C	141	<p>3% 43% 21% 10% 25%</p>
2	F	141	<p>5% 51% 16% 29%</p>
2	I	141	<p>2% 47% 20% 6% 27%</p>
3	J	84	<p>65% 21% 12%</p>
3	K	84	<p>55% 31% 12%</p>
3	L	84	<p>64% 19% 5% 12%</p>
3	M	84	<p>7% 57% 25% 6% 12%</p>
3	N	84	<p>70% 15% 12%</p>
3	O	84	<p>70% 13% 5% 12%</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10508 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Centromere protein S.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	98	758	477	131	145	3	2	0	0	0
1	B	103	769	480	134	150	3	2	0	0	0
1	D	97	736	460	125	146	3	2	0	1	0
1	E	97	699	436	124	134	3	2	0	0	0
1	G	105	793	492	141	155	3	2	0	0	0
1	H	94	731	459	125	142	3	2	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q8N2Z9
A	-4	HIS	-	expression tag	UNP Q8N2Z9
A	-3	HIS	-	expression tag	UNP Q8N2Z9
A	-2	HIS	-	expression tag	UNP Q8N2Z9
A	-1	HIS	-	expression tag	UNP Q8N2Z9
A	0	HIS	-	expression tag	UNP Q8N2Z9
B	-5	HIS	-	expression tag	UNP Q8N2Z9
B	-4	HIS	-	expression tag	UNP Q8N2Z9
B	-3	HIS	-	expression tag	UNP Q8N2Z9
B	-2	HIS	-	expression tag	UNP Q8N2Z9
B	-1	HIS	-	expression tag	UNP Q8N2Z9
B	0	HIS	-	expression tag	UNP Q8N2Z9
D	-5	HIS	-	expression tag	UNP Q8N2Z9
D	-4	HIS	-	expression tag	UNP Q8N2Z9
D	-3	HIS	-	expression tag	UNP Q8N2Z9
D	-2	HIS	-	expression tag	UNP Q8N2Z9
D	-1	HIS	-	expression tag	UNP Q8N2Z9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	HIS	-	expression tag	UNP Q8N2Z9
E	-5	HIS	-	expression tag	UNP Q8N2Z9
E	-4	HIS	-	expression tag	UNP Q8N2Z9
E	-3	HIS	-	expression tag	UNP Q8N2Z9
E	-2	HIS	-	expression tag	UNP Q8N2Z9
E	-1	HIS	-	expression tag	UNP Q8N2Z9
E	0	HIS	-	expression tag	UNP Q8N2Z9
G	-5	HIS	-	expression tag	UNP Q8N2Z9
G	-4	HIS	-	expression tag	UNP Q8N2Z9
G	-3	HIS	-	expression tag	UNP Q8N2Z9
G	-2	HIS	-	expression tag	UNP Q8N2Z9
G	-1	HIS	-	expression tag	UNP Q8N2Z9
G	0	HIS	-	expression tag	UNP Q8N2Z9
H	-5	HIS	-	expression tag	UNP Q8N2Z9
H	-4	HIS	-	expression tag	UNP Q8N2Z9
H	-3	HIS	-	expression tag	UNP Q8N2Z9
H	-2	HIS	-	expression tag	UNP Q8N2Z9
H	-1	HIS	-	expression tag	UNP Q8N2Z9
H	0	HIS	-	expression tag	UNP Q8N2Z9

- Molecule 2 is a protein called Fanconi anemia group M protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	106	Total	C	N	O	S	0	0	0
			860	542	151	161	6			
2	F	100	Total	C	N	O	S	0	1	0
			796	505	141	144	6			
2	I	103	Total	C	N	O	S	0	1	0
			857	542	151	158	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	660	GLY	-	expression tag	UNP Q8IYD8
F	660	GLY	-	expression tag	UNP Q8IYD8
I	660	GLY	-	expression tag	UNP Q8IYD8

- Molecule 3 is a protein called Centromere protein X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	J	74	Total	C	N	O	Se	0	0	0
			581	371	102	107	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	K	74	Total	C	N	O	Se	0	0	0
			570	365	100	104	1			
3	L	74	Total	C	N	O	Se	0	0	0
			558	358	96	103	1			
3	M	74	Total	C	N	O	Se	0	0	0
			529	340	88	100	1			
3	N	74	Total	C	N	O	Se	0	0	0
			567	363	97	106	1			
3	O	74	Total	C	N	O	Se	0	0	0
			584	374	102	107	1			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	-2	GLY	-	expression tag	UNP A8MT69
J	-1	SER	-	expression tag	UNP A8MT69
J	0	HIS	-	expression tag	UNP A8MT69
K	-2	GLY	-	expression tag	UNP A8MT69
K	-1	SER	-	expression tag	UNP A8MT69
K	0	HIS	-	expression tag	UNP A8MT69
L	-2	GLY	-	expression tag	UNP A8MT69
L	-1	SER	-	expression tag	UNP A8MT69
L	0	HIS	-	expression tag	UNP A8MT69
M	-2	GLY	-	expression tag	UNP A8MT69
M	-1	SER	-	expression tag	UNP A8MT69
M	0	HIS	-	expression tag	UNP A8MT69
N	-2	GLY	-	expression tag	UNP A8MT69
N	-1	SER	-	expression tag	UNP A8MT69
N	0	HIS	-	expression tag	UNP A8MT69
O	-2	GLY	-	expression tag	UNP A8MT69
O	-1	SER	-	expression tag	UNP A8MT69
O	0	HIS	-	expression tag	UNP A8MT69

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		
4	B	10	Total	O	0	0
			10	10		
4	C	17	Total	O	0	0
			17	17		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	3	Total O 3 3	0	0
4	E	3	Total O 3 3	0	0
4	F	3	Total O 3 3	0	0
4	G	8	Total O 8 8	0	0
4	H	12	Total O 12 12	0	0
4	I	18	Total O 18 18	0	0
4	J	3	Total O 3 3	0	0
4	K	13	Total O 13 13	0	0
4	L	8	Total O 8 8	0	0
4	M	3	Total O 3 3	0	0
4	N	4	Total O 4 4	0	0
4	O	8	Total O 8 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Centromere protein S




- Molecule 1: Centromere protein S




- Molecule 1: Centromere protein S





- Molecule 1: Centromere protein S








• Molecule 1: Centromere protein S





• Molecule 1: Centromere protein S



• Molecule 2: Fanconi anemia group M protein



• Molecule 2: Fanconi anemia group M protein



• Molecule 2: Fanconi anemia group M protein






• Molecule 3: Centromere protein X




• Molecule 3: Centromere protein X



• Molecule 3: Centromere protein X




• Molecule 3: Centromere protein X



• Molecule 3: Centromere protein X



• Molecule 3: Centromere protein X



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	111.32Å 70.03Å 115.75Å 90.00° 91.41° 90.00°	Depositor
Resolution (Å)	49.67 – 2.63 49.67 – 2.63	Depositor EDS
% Data completeness (in resolution range)	95.1 (49.67-2.63) 99.5 (49.67-2.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.87 (at 2.65Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6_289)	Depositor
R, R_{free}	0.221 , 0.256 0.219 , 0.254	Depositor DCC
R_{free} test set	2693 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtrriage
Anisotropy	0.390	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for l,k,-h 0.020 for h,-k,-l 0.012 for l,-k,h	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10508	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/765	0.65	0/1027
1	B	0.37	0/775	0.70	0/1045
1	D	0.28	0/743	0.66	0/1001
1	E	0.33	0/704	0.72	0/953
1	G	0.27	0/799	0.67	0/1074
1	H	0.29	0/738	0.63	0/991
2	C	0.31	0/882	0.74	2/1195 (0.2%)
2	F	0.24	0/818	0.66	0/1111
2	I	0.25	0/882	0.61	0/1194
3	J	0.27	0/586	0.64	0/788
3	K	0.28	0/575	0.65	0/777
3	L	0.27	0/563	0.64	0/762
3	M	0.26	0/534	0.62	0/728
3	N	0.27	0/572	0.65	0/772
3	O	0.27	0/589	0.61	0/792
All	All	0.29	0/10525	0.66	2/14210 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	705	LEU	CA-C-N	5.68	126.95	119.84
2	C	705	LEU	C-N-CA	5.68	126.95	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	758	0	724	29	0
1	B	769	0	715	27	0
1	D	736	0	673	17	0
1	E	699	0	630	22	0
1	G	793	0	740	15	0
1	H	731	0	698	16	0
2	C	860	0	771	49	0
2	F	796	0	679	22	0
2	I	857	0	781	30	0
3	J	581	0	593	15	0
3	K	570	0	571	22	0
3	L	558	0	547	16	0
3	M	529	0	491	23	0
3	N	567	0	565	10	0
3	O	584	0	602	14	0
4	A	7	0	0	1	0
4	B	10	0	0	0	0
4	C	17	0	0	1	0
4	D	3	0	0	0	0
4	E	3	0	0	0	0
4	F	3	0	0	0	0
4	G	8	0	0	1	0
4	H	12	0	0	0	0
4	I	18	0	0	0	0
4	J	3	0	0	0	0
4	K	13	0	0	0	0
4	L	8	0	0	0	0
4	M	3	0	0	0	0
4	N	4	0	0	0	0
4	O	8	0	0	0	0
All	All	10508	0	9780	266	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 266 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:676:LYS:HG2	2:C:677:ASP:H	1.23	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:771:GLU:HA	2:C:772:GLU:CB	1.94	0.95
1:A:14:SER:H	1:A:17:GLN:HE21	1.19	0.90
1:E:99:LYS:HG2	3:M:40:GLU:HG2	1.54	0.86
1:E:61:PHE:HB3	3:M:38:MSE:HE1	1.58	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/120 (80%)	96 (100%)	0	0	100	100
1	B	101/120 (84%)	97 (96%)	3 (3%)	1 (1%)	12	18
1	D	96/120 (80%)	96 (100%)	0	0	100	100
1	E	95/120 (79%)	91 (96%)	3 (3%)	1 (1%)	11	16
1	G	103/120 (86%)	103 (100%)	0	0	100	100
1	H	92/120 (77%)	92 (100%)	0	0	100	100
2	C	102/141 (72%)	94 (92%)	5 (5%)	3 (3%)	3	4
2	F	95/141 (67%)	91 (96%)	4 (4%)	0	100	100
2	I	98/141 (70%)	92 (94%)	6 (6%)	0	100	100
3	J	72/84 (86%)	71 (99%)	1 (1%)	0	100	100
3	K	72/84 (86%)	72 (100%)	0	0	100	100
3	L	72/84 (86%)	72 (100%)	0	0	100	100
3	M	72/84 (86%)	69 (96%)	3 (4%)	0	100	100
3	N	72/84 (86%)	71 (99%)	1 (1%)	0	100	100
3	O	72/84 (86%)	72 (100%)	0	0	100	100
All	All	1310/1647 (80%)	1279 (98%)	26 (2%)	5 (0%)	30	42

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	105	GLN
2	C	772	GLU
2	C	706	PRO
2	C	707	GLN
1	E	103	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	75/104 (72%)	65 (87%)	10 (13%)	4 5
1	B	74/104 (71%)	72 (97%)	2 (3%)	39 60
1	D	71/104 (68%)	66 (93%)	5 (7%)	14 23
1	E	63/104 (61%)	57 (90%)	6 (10%)	8 12
1	G	77/104 (74%)	70 (91%)	7 (9%)	9 13
1	H	74/104 (71%)	69 (93%)	5 (7%)	14 23
2	C	90/133 (68%)	74 (82%)	16 (18%)	2 2
2	F	77/133 (58%)	67 (87%)	10 (13%)	4 5
2	I	92/133 (69%)	79 (86%)	13 (14%)	3 4
3	J	62/67 (92%)	59 (95%)	3 (5%)	23 37
3	K	59/67 (88%)	53 (90%)	6 (10%)	7 10
3	L	56/67 (84%)	49 (88%)	7 (12%)	4 6
3	M	50/67 (75%)	42 (84%)	8 (16%)	2 2
3	N	59/67 (88%)	56 (95%)	3 (5%)	21 35
3	O	63/67 (94%)	58 (92%)	5 (8%)	11 18
All	All	1042/1425 (73%)	936 (90%)	106 (10%)	7 10

5 of 106 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	38	LYS
2	I	781	VAL
3	N	68	ASP
1	H	75	THR
2	I	712	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	786	GLN
3	L	69	GLN
3	J	76	GLN
3	K	69	GLN
3	M	56	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	96/120 (80%)	0.04	3 (3%) 51 45	23, 40, 81, 95	0
1	B	101/120 (84%)	0.14	1 (0%) 79 76	27, 42, 75, 89	0
1	D	95/120 (79%)	0.32	3 (3%) 50 44	29, 50, 71, 81	1 (1%)
1	E	95/120 (79%)	0.59	5 (5%) 32 26	41, 63, 80, 92	0
1	G	103/120 (85%)	0.09	2 (1%) 66 62	28, 41, 76, 89	0
1	H	92/120 (76%)	-0.05	2 (2%) 62 58	25, 38, 66, 81	0
2	C	106/141 (75%)	0.16	4 (3%) 44 38	26, 39, 68, 72	0
2	F	100/141 (70%)	0.63	7 (7%) 22 18	21, 54, 80, 86	1 (1%)
2	I	103/141 (73%)	0.00	3 (2%) 53 48	20, 36, 58, 70	1 (0%)
3	J	73/84 (86%)	-0.22	0 100 100	25, 34, 52, 59	0
3	K	73/84 (86%)	-0.03	0 100 100	30, 38, 54, 58	0
3	L	73/84 (86%)	0.14	1 (1%) 73 70	30, 43, 59, 72	0
3	M	73/84 (86%)	0.77	6 (8%) 17 14	35, 61, 75, 79	0
3	N	73/84 (86%)	-0.08	1 (1%) 73 70	29, 38, 55, 64	0
3	O	73/84 (86%)	-0.28	0 100 100	23, 33, 50, 54	0
All	All	1329/1647 (80%)	0.16	38 (2%) 53 48	20, 42, 74, 95	3 (0%)

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	708	VAL	4.2
1	G	108	LEU	4.2
3	M	25	ASP	4.1
3	M	28	THR	4.0
2	F	712	SER	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.