



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 10:06 PM UTC

PDB ID : 7DRS / pdb_00007drs
Title : Structure of SspE_40224
Authors : Haiyan, G.; Jinchuan, Z.; Chen, S.; Wang, L.; Wu, G.
Deposited on : 2020-12-29
Resolution : 3.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1210 (3.48-3.36)
Clashscore	190562	1234 (3.48-3.36)
Ramachandran outliers	187476	1222 (3.48-3.36)
Sidechain outliers	187428	1222 (3.48-3.36)
RSRZ outliers	180081	1210 (3.48-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	771	 45% 46% 8% .
1	B	771	 47% 41% 11% .
1	C	771	 44% 41% 12% .
1	D	771	 44% 43% 11% .

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 24548 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SspE protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	771	6137	3872	1079	1166	20	0	0	0
1	A	771	6137	3872	1079	1166	20	0	0	0
1	B	771	6137	3872	1079	1166	20	0	0	0
1	C	771	6137	3872	1079	1166	20	0	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SspE protein




• Molecule 1: SspE protein


Chain A: 45% 46% 8%





• Molecule 1: SspE protein


Chain B: 47% 41% 11%





• Molecule 1: SspE protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	111.08Å 138.15Å 293.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.14 – 3.42 46.14 – 3.42	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.14-3.42) 99.7 (46.14-3.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 3.40Å)	Xtrriage
Refinement program	REFMAC 5.5	Depositor
R, R_{free}	0.212 , 0.264 0.221 , 0.268	Depositor DCC
R_{free} test set	3032 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	121.6	Xtrriage
Anisotropy	0.343	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 126.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24548	wwPDB-VP
Average B, all atoms (Å ²)	156.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/6272	0.91	7/8515 (0.1%)
1	B	0.57	1/6272 (0.0%)	0.88	9/8515 (0.1%)
1	C	0.76	3/6272 (0.0%)	1.06	25/8515 (0.3%)
1	D	0.66	2/6272 (0.0%)	1.00	16/8515 (0.2%)
All	All	0.65	6/25088 (0.0%)	0.97	57/34060 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	5
1	D	0	3
All	All	0	11

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	209	ASP	C-O	6.25	1.27	1.23
1	C	151	MET	CG-SD	-5.72	1.66	1.80
1	D	145	ASN	CA-C	5.49	1.62	1.52
1	C	161	ARG	CB-CG	-5.33	1.36	1.52
1	D	170	TRP	CG-CD2	-5.12	1.34	1.43

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	145	ASN	CA-C-O	8.74	132.69	119.23
1	C	151	MET	CA-C-N	-7.74	108.86	122.26
1	C	151	MET	C-N-CA	-7.74	108.86	122.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	345	GLY	N-CA-C	-7.16	92.53	113.30
1	A	417	ILE	N-CA-C	-6.55	95.72	109.34

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	207	PRO	Peptide
1	A	553	VAL	Peptide
1	D	160	PRO	Peptide
1	D	173	ASN	Peptide
1	D	79	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6137	0	6046	353	1
1	B	6137	0	6046	367	1
1	C	6137	0	6046	369	2
1	D	6137	0	6046	379	2
All	All	24548	0	24184	1444	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 1444 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:223:HIS:HE1	1:C:227:MET:HE2	1.19	1.07
1:C:628:ARG:HA	1:C:631:ASP:HB2	1.44	0.99
1:A:418:VAL:H	1:A:421:ARG:HD2	1.25	0.98
1:B:126:ASP:HB3	1:B:129:SER:HB3	1.45	0.98
1:A:126:ASP:HB3	1:A:129:SER:HB3	1.44	0.96

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:85:SER:N	1:C:211:ASP:OD2[4_457]	1.94	0.26
1:D:84:ARG:N	1:C:211:ASP:OD2[4_457]	2.13	0.07
1:A:236:ARG:NH1	1:B:278:GLU:OE1[4_447]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	769/771 (100%)	613 (80%)	105 (14%)	51 (7%)	1 7
1	B	769/771 (100%)	605 (79%)	117 (15%)	47 (6%)	1 8
1	C	769/771 (100%)	612 (80%)	104 (14%)	53 (7%)	1 6
1	D	769/771 (100%)	609 (79%)	105 (14%)	55 (7%)	1 5
All	All	3076/3084 (100%)	2439 (79%)	431 (14%)	206 (7%)	1 7

5 of 206 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	4	LYS
1	D	36	TRP
1	D	127	GLU
1	D	200	VAL
1	D	205	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	651/651 (100%)	561 (86%)	90 (14%)	3	14
1	B	651/651 (100%)	563 (86%)	88 (14%)	4	15
1	C	651/651 (100%)	547 (84%)	104 (16%)	2	10
1	D	651/651 (100%)	559 (86%)	92 (14%)	3	13
All	All	2604/2604 (100%)	2230 (86%)	374 (14%)	3	13

5 of 374 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	349	TRP
1	C	99	GLN
1	B	396	LEU
1	B	636	THR
1	C	167	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	557	GLN
1	C	279	GLN
1	B	16	GLN
1	C	391	GLN
1	B	586	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	771/771 (100%)	-0.37	1 (0%) 92 92	84, 138, 228, 331	0
1	B	771/771 (100%)	-0.31	5 (0%) 85 75	74, 180, 334, 405	0
1	C	771/771 (100%)	-0.42	4 (0%) 87 78	57, 129, 243, 347	0
1	D	771/771 (100%)	-0.43	3 (0%) 88 81	67, 143, 228, 341	0
All	All	3084/3084 (100%)	-0.38	13 (0%) 88 81	57, 142, 295, 405	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	594	LEU	4.0
1	B	458	ILE	3.1
1	C	714	LEU	2.7
1	D	623	THR	2.3
1	B	595	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.