



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 5, 2026 – 11:51 AM UTC

PDB ID : 9DSX / pdb_00009dsx
Title : Crystal structure of the complex of M. tuberculosis PheRS with cognate precursor tRNA and fragment DDD00107555
Authors : Chang, C.; Michalska, K.; Forte, B.; Baragana, B.; Gilbert, I.H.; Wower, J.; Joachimiak, A.; Center for Structural Biology of Infectious Diseases (CSBID)
Deposited on : 2024-09-30
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	835	15% 88% 8% . .
3	C	77	6% 86% 9% 5%
3	F	77	26% 70% 17% 8% 5%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 22325 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylalanine-tRNA ligase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	Total	C	N	O	S	0	4	0
			2545	1606	458	472	9			
1	D	339	Total	C	N	O	S	0	8	0
			2680	1691	483	497	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ALA	-	expression tag	UNP P9WUFU3
D	0	ALA	-	expression tag	UNP P9WUFU3

- Molecule 2 is a protein called Phenylalanine-tRNA ligase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	834	Total	C	N	O	S	0	11	0
			6321	3971	1146	1182	22			
2	E	804	Total	C	N	O	S	0	10	0
			6088	3824	1109	1135	20			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLN	-	expression tag	UNP P9WUFU1
B	-2	SER	-	expression tag	UNP P9WUFU1
B	-1	ASN	-	expression tag	UNP P9WUFU1
B	0	ALA	-	expression tag	UNP P9WUFU1
E	-3	GLN	-	expression tag	UNP P9WUFU1
E	-2	SER	-	expression tag	UNP P9WUFU1
E	-1	ASN	-	expression tag	UNP P9WUFU1
E	0	ALA	-	expression tag	UNP P9WUFU1

- Molecule 3 is a RNA chain called tRNA(Phe).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	73	Total 1563	C 695	N 283	O 512	P 73	0	0	0
3	F	73	Total 1563	C 695	N 283	O 512	P 73	0	0	0

- Molecule 4 is ethyl (2R)-2-(3-oxo-2,3-dihydro-4H-1,4-benzoxazin-4-yl)propanoate (CCD ID: A1BCB) (formula: C₁₃H₁₅NO₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	Total 18	C 13	N 1	O 4	0	0
4	D	1	Total 18	C 13	N 1	O 4	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	Total 1	Mg 1	0	0
5	C	3	Total 3	Mg 3	0	0
5	D	1	Total 1	Mg 1	0	0

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Na 1 1	0	0
7	E	1	Total Na 1 1	0	0

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	C	1	Total C O 6 3 3	0	0
8	D	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0
8	F	1	Total C O 6 3 3	0	0

- Molecule 9 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	O	S	0	0
			4	2	1	1		
9	B	1	Total	C	O	S	0	0
			4	2	1	1		
9	D	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 10 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			10	6	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			10	6	4		
10	E	1	Total	C	O	0	0
			10	6	4		

- Molecule 11 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
11	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	101	Total	O	0	0
			101	101		
12	B	633	Total	O	0	0
			633	633		
12	C	115	Total	O	0	0
			115	115		
12	D	114	Total	O	0	0
			114	114		
12	E	359	Total	O	0	0
			359	359		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	F	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenylalanine-tRNA ligase alpha subunit




- Molecule 1: Phenylalanine-tRNA ligase alpha subunit





- Molecule 2: Phenylalanine-tRNA ligase beta subunit








● Molecule 2: Phenylalanine-tRNA ligase beta subunit



● Molecule 3: tRNA(Phe)



● Molecule 3: tRNA(Phe)



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	147.36Å 65.03Å 189.49Å 90.00° 110.86° 90.00°	Depositor
Resolution (Å)	48.86 – 2.05 48.86 – 2.05	Depositor EDS
% Data completeness (in resolution range)	94.3 (48.86-2.05) 94.3 (48.86-2.05)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.05Å)	Xtrriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.199 , 0.233 0.199 , 0.233	Depositor DCC
R_{free} test set	9901 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22325	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, ACT, EPE, A1BCB, MG, NA, PGE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.10	0/2604	0.28	0/3546
1	D	0.10	0/2739	0.27	0/3726
2	B	0.11	0/6458	0.28	0/8842
2	E	0.09	0/6218	0.26	0/8513
3	C	0.09	0/1746	0.22	0/2720
3	F	0.07	0/1746	0.18	0/2720
All	All	0.10	0/21511	0.26	0/30067

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2545	0	2484	21	0
1	D	2680	0	2641	22	0
2	B	6321	0	6344	24	0
2	E	6088	0	6118	40	0
3	C	1563	0	792	1	0
3	F	1563	0	792	10	0
4	A	18	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	18	0	0	0	0
5	A	1	0	0	0	0
5	C	3	0	0	0	0
5	D	1	0	0	0	0
6	A	4	0	3	1	0
6	B	12	0	9	0	0
6	E	12	0	9	2	0
7	A	1	0	0	0	0
7	E	1	0	0	0	0
8	B	30	0	40	3	0
8	C	6	0	8	0	0
8	D	6	0	8	2	0
8	E	18	0	24	0	0
8	F	6	0	8	0	0
9	B	8	0	12	0	0
9	D	4	0	6	1	0
10	B	20	0	28	3	0
10	E	10	0	14	3	0
11	B	30	0	36	0	0
12	A	101	0	0	1	0
12	B	633	0	0	6	0
12	C	115	0	0	0	0
12	D	114	0	0	1	0
12	E	359	0	0	3	0
12	F	34	0	0	1	0
All	All	22325	0	19376	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:4:C:H42	3:F:70:G:H1	1.11	0.96
3:C:20:U:H5 ^{''}	1:D:46:GLN:HG2	1.71	0.71
2:E:49:PRO:HB2	2:E:100:LEU:HB2	1.72	0.70
1:D:2:LEU:O	1:D:62:ARG:NH1	2.23	0.69
2:E:27:GLU:HG3	2:E:37:VAL:HG21	1.78	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/342 (96%)	316 (96%)	11 (3%)	1 (0%)	36	30
1	D	343/342 (100%)	335 (98%)	8 (2%)	0	100	100
2	B	843/835 (101%)	824 (98%)	18 (2%)	1 (0%)	48	43
2	E	804/835 (96%)	779 (97%)	22 (3%)	3 (0%)	30	22
All	All	2318/2354 (98%)	2254 (97%)	59 (2%)	5 (0%)	43	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	158	GLN
2	B	373	THR
2	E	88	THR
2	E	373[A]	THR
2	E	373[B]	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/268 (96%)	255 (99%)	2 (1%)	73	77
1	D	273/268 (102%)	270 (99%)	3 (1%)	65	68
2	B	661/652 (101%)	658 (100%)	3 (0%)	81	85
2	E	636/652 (98%)	624 (98%)	12 (2%)	50	50
All	All	1827/1840 (99%)	1807 (99%)	20 (1%)	65	68

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	360	LEU
2	E	564	THR
2	E	607	VAL
2	E	584	VAL
1	D	216	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	170	GLN
1	A	215	GLN
1	D	170	GLN
2	E	132	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	72/77 (93%)	6 (8%)	0
3	F	72/77 (93%)	12 (16%)	0
All	All	144/154 (93%)	18 (12%)	0

5 of 18 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	16	U
3	C	21	A
3	C	22	U
3	C	23	G
3	C	43	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 7 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	A1BCB	A	401	-	18,19,19	2.34	4 (22%)	23,26,26	1.48	4 (17%)
8	GOL	B	904	-	5,5,5	0.34	0	5,5,5	0.42	0
8	GOL	B	913	-	5,5,5	0.34	0	5,5,5	0.42	0
6	ACT	B	912	-	3,3,3	1.14	0	3,3,3	1.20	0
9	DMS	B	907	-	3,3,3	0.58	0	3,3,3	0.14	0
8	GOL	F	101	-	5,5,5	0.33	0	5,5,5	0.39	0
11	EPE	B	909	-	15,15,15	0.84	1 (6%)	19,20,20	0.53	0
6	ACT	E	906	-	3,3,3	1.12	0	3,3,3	1.22	0
8	GOL	B	902	-	5,5,5	0.33	0	5,5,5	0.37	0
6	ACT	E	905	-	3,3,3	1.14	0	3,3,3	1.21	0
8	GOL	D	401	-	5,5,5	0.32	0	5,5,5	0.39	0
8	GOL	B	903	-	5,5,5	0.35	0	5,5,5	0.40	0
9	DMS	D	402	-	3,3,3	0.57	0	3,3,3	0.24	0
10	PGE	B	914	-	9,9,9	0.33	0	8,8,8	0.54	0
4	A1BCB	D	403	-	18,19,19	2.36	5 (27%)	23,26,26	1.56	5 (21%)
8	GOL	E	908	-	5,5,5	0.35	0	5,5,5	0.42	0
8	GOL	E	903	-	5,5,5	0.34	0	5,5,5	0.41	0
8	GOL	C	104	-	5,5,5	0.34	0	5,5,5	0.43	0
6	ACT	E	904	-	3,3,3	1.13	0	3,3,3	1.20	0
6	ACT	B	910	-	3,3,3	1.13	0	3,3,3	1.17	0
8	GOL	E	902	-	5,5,5	0.32	0	5,5,5	0.33	0
11	EPE	B	908	-	15,15,15	0.80	1 (6%)	19,20,20	0.68	0
10	PGE	B	906	-	9,9,9	0.32	0	8,8,8	0.45	0
9	DMS	B	905	-	3,3,3	0.58	0	3,3,3	0.13	0
8	GOL	B	901	-	5,5,5	0.32	0	5,5,5	0.54	0
6	ACT	A	403	-	3,3,3	1.12	0	3,3,3	1.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ACT	B	911	-	3,3,3	1.13	0	3,3,3	1.17	0
10	PGE	E	901	-	9,9,9	0.32	0	8,8,8	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	D	401	-	-	0/4/4/4	-
8	GOL	B	903	-	-	2/4/4/4	-
4	A1BCB	A	401	-	-	5/11/24/24	0/2/2/2
10	PGE	B	914	-	-	4/7/7/7	-
10	PGE	E	901	-	-	4/7/7/7	-
8	GOL	B	904	-	-	0/4/4/4	-
4	A1BCB	D	403	-	-	5/11/24/24	0/2/2/2
8	GOL	B	913	-	-	0/4/4/4	-
8	GOL	E	908	-	-	0/4/4/4	-
8	GOL	E	903	-	-	2/4/4/4	-
8	GOL	C	104	-	-	2/4/4/4	-
8	GOL	E	902	-	-	4/4/4/4	-
8	GOL	F	101	-	-	0/4/4/4	-
11	EPE	B	908	-	-	3/9/19/19	0/1/1/1
11	EPE	B	909	-	-	3/9/19/19	0/1/1/1
10	PGE	B	906	-	-	5/7/7/7	-
8	GOL	B	901	-	-	2/4/4/4	-
8	GOL	B	902	-	-	2/4/4/4	-

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	403	A1BCB	O03-C02	7.44	1.46	1.37
4	A	401	A1BCB	O03-C02	7.37	1.45	1.37
4	D	403	A1BCB	O12-C10	3.43	1.40	1.33
4	A	401	A1BCB	O12-C10	3.41	1.40	1.33
4	D	403	A1BCB	C05-N07	2.98	1.42	1.36

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	403	A1BCB	O12-C10-C08	4.44	121.44	111.14
4	A	401	A1BCB	O12-C10-C08	4.39	121.32	111.14
4	A	401	A1BCB	O12-C10-O11	-2.57	119.43	124.14
4	D	403	A1BCB	C08-N07-C05	2.55	121.50	117.43
4	D	403	A1BCB	O12-C10-O11	-2.46	119.64	124.14

There are no chirality outliers.

5 of 43 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	401	A1BCB	N07-C08-C10-O11
4	A	401	A1BCB	N07-C08-C10-O12
4	D	403	A1BCB	N07-C08-C10-O11
4	D	403	A1BCB	N07-C08-C10-O12
8	B	901	GOL	O1-C1-C2-C3


There are no ring outliers.

10 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	913	GOL	1	0
6	E	906	ACT	1	0
8	D	401	GOL	2	0
8	B	903	GOL	2	0
9	D	402	DMS	1	0
10	B	914	PGE	1	0
6	E	904	ACT	1	0
10	B	906	PGE	2	0
6	A	403	ACT	1	0
10	E	901	PGE	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/342 (95%)	1.02	75 (22%) 2 1	10, 33, 100, 140	4 (1%)
1	D	339/342 (99%)	0.76	46 (13%) 7 6	12, 35, 71, 99	8 (2%)
2	B	834/835 (99%)	-0.14	22 (2%) 57 59	6, 22, 47, 80	11 (1%)
2	E	804/835 (96%)	0.65	126 (15%) 5 4	10, 33, 87, 126	10 (1%)
3	C	73/77 (94%)	-0.17	5 (6%) 23 23	16, 40, 156, 201	0
3	F	73/77 (94%)	1.02	20 (27%) 1 1	28, 63, 205, 235	0
All	All	2451/2508 (97%)	0.43	294 (11%) 9 8	6, 29, 87, 235	33 (1%)

The worst 5 of 294 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	90[A]	LEU	10.5
1	D	91[A]	VAL	8.7
1	A	106[A]	ALA	7.4
1	A	105[A]	PRO	7.2
1	D	341	ALA	6.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	B	902	6/6	0.70	0.19	27,43,52,53	0
8	GOL	B	903	6/6	0.71	0.18	48,49,58,62	0
10	PGE	B	914	10/10	0.72	0.20	27,37,49,57	0
8	GOL	E	902	6/6	0.73	0.17	36,46,54,55	0
8	GOL	B	913	6/6	0.74	0.14	42,47,50,66	0
4	A1BCB	A	401	18/18	0.74	0.17	29,40,60,64	0
4	A1BCB	D	403	18/18	0.74	0.17	31,45,68,69	0
6	ACT	B	912	4/4	0.76	0.17	44,46,49,50	0
8	GOL	F	101	6/6	0.78	0.23	39,43,44,45	0
10	PGE	B	906	10/10	0.79	0.17	15,32,38,42	10
6	ACT	A	403	4/4	0.79	0.21	36,42,55,62	0
10	PGE	E	901	10/10	0.80	0.16	33,46,52,56	0
8	GOL	E	908	6/6	0.82	0.15	49,52,59,61	0
6	ACT	E	906	4/4	0.86	0.13	36,51,54,55	0
8	GOL	D	401	6/6	0.86	0.13	38,42,44,44	0
9	DMS	D	402	4/4	0.87	0.19	42,50,53,68	0
8	GOL	C	104	6/6	0.87	0.27	25,35,49,50	0
6	ACT	E	905	4/4	0.88	0.14	36,40,44,44	0
8	GOL	E	903	6/6	0.88	0.10	41,43,45,57	0
11	EPE	B	908	15/15	0.88	0.14	32,41,54,55	0
6	ACT	E	904	4/4	0.89	0.12	33,37,44,45	0
8	GOL	B	901	6/6	0.89	0.12	22,25,31,35	0
8	GOL	B	904	6/6	0.89	0.10	41,48,51,58	0
11	EPE	B	909	15/15	0.89	0.12	37,44,55,60	0
6	ACT	B	911	4/4	0.92	0.10	23,24,37,38	0
7	NA	E	907	1/1	0.93	0.21	37,37,37,37	0
9	DMS	B	905	4/4	0.93	0.15	28,34,47,58	0
9	DMS	B	907	4/4	0.93	0.12	33,45,52,54	0
6	ACT	B	910	4/4	0.94	0.09	17,27,28,33	0
5	MG	C	101	1/1	0.95	0.07	43,43,43,43	0
5	MG	C	103	1/1	0.96	0.08	33,33,33,33	0
7	NA	A	404	1/1	0.98	0.04	28,28,28,28	0
5	MG	D	404	1/1	0.98	0.03	21,21,21,21	0
5	MG	A	402	1/1	0.98	0.03	16,16,16,16	0
5	MG	C	102	1/1	0.99	0.03	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.