



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:30 PM UTC

PDB ID : 2DTH / pdb\_00002dth  
Title : The Crystal Structure of the Orthorhombic Form of Biotin Protein Ligase From *Pyrococcus Horikoshii* OT3 in Complex with Biotin and ADP  
Authors : Bagautdinov, B.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2006-07-12  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	
1	B	235	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	235	Total	C	N	O	S	0	0	0
			1837	1185	311	336	5			
1	B	235	Total	C	N	O	S	0	0	0
			1837	1185	311	336	5			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total	C	N	O	0	0
			17	9	5	3		
2	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 3 is BIOTIN (CCD ID: BTN) (formula:  $C_{10}H_{16}N_2O_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	16	10	2	3	1	0	0
3	B	1	16	10	2	3	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	312	312	312	0	0
4	B	299	299	299	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase



- Molecule 1: 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.00Å 72.72Å 104.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.24 – 1.95 36.24 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.6 (36.24-1.95) 98.6 (36.24-1.95)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.84 (at 1.95Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.240 , 0.269 0.240 , 0.268	Depositor DCC
$R_{free}$ test set	1940 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.6	Xtrriage
Anisotropy	0.360	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 53.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4351	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9474e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BTN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1865	0.89	7/2515 (0.3%)
1	B	0.35	0/1865	0.84	1/2515 (0.0%)
All	All	0.36	0/3730	0.87	8/5030 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	ARG	N-CA-C	-7.59	103.02	111.82
1	A	30	TYR	N-CA-C	-6.06	100.13	109.76
1	A	210	PHE	N-CA-C	-6.04	106.24	113.97
1	B	11	ARG	N-CA-C	-5.78	105.07	111.71
1	A	47	GLY	CA-C-N	5.63	128.13	120.54
1	A	47	GLY	C-N-CA	5.63	128.13	120.54
1	A	135	LYS	N-CA-C	-5.37	102.93	110.50
1	A	205	GLU	N-CA-C	5.20	116.75	111.14

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1837	0	1934	51	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1837	0	1934	60	0
2	A	17	0	9	0	0
2	B	17	0	9	1	0
3	A	16	0	15	0	0
3	B	16	0	15	0	0
4	A	312	0	0	14	0
4	B	299	0	0	11	0
All	All	4351	0	3916	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:11:ARG:HH11	1:B:11:ARG:HB2	1.23	0.99
1:B:194:ILE:HD13	1:B:225:VAL:HG21	1.56	0.87
1:A:186:MET:HE3	1:A:188:LEU:HD21	1.58	0.85
1:B:206:ASP:OD1	1:B:207:ILE:N	2.16	0.78
1:B:51:ARG:HB2	4:B:1452:HOH:O	1.88	0.73
1:B:11:ARG:HH11	1:B:11:ARG:CB	1.99	0.73
1:A:186:MET:HE1	1:A:207:ILE:HG23	1.71	0.72
1:A:139:GLY:HA3	4:A:1614:HOH:O	1.90	0.72
1:B:180:ASN:ND2	4:B:1440:HOH:O	2.22	0.72
1:A:48:ARG:NH1	4:A:1495:HOH:O	2.25	0.69
1:B:186:MET:HE1	1:B:207:ILE:HG23	1.75	0.68
1:B:230:VAL:HB	4:B:1615:HOH:O	1.94	0.67
1:B:48:ARG:HG2	1:B:49:LEU:HD13	1.77	0.66
1:A:71:PRO:HG2	1:A:74:ASP:OD2	1.96	0.66
1:B:110:LYS:HE3	1:B:141:THR:HB	1.78	0.66
1:B:91:LYS:HD2	1:B:91:LYS:O	1.95	0.65
1:A:79:VAL:HB	4:A:1463:HOH:O	1.96	0.65
1:A:51:ARG:HD3	4:A:1648:HOH:O	1.97	0.64
1:A:179:LEU:HD13	1:A:209:ASP:HA	1.79	0.64
1:B:91:LYS:HD2	1:B:91:LYS:C	2.22	0.63
1:A:177:ASP:O	1:A:180:ASN:ND2	2.32	0.63
1:A:27:LYS:C	1:A:29:SER:H	2.07	0.63
1:B:74:ASP:HA	1:B:77:LYS:NZ	2.13	0.63
1:B:208:ASP:OD2	1:B:224:LYS:HE2	1.99	0.63
1:B:1:MET:HE2	1:B:2:LEU:HD23	1.81	0.62
1:B:194:ILE:HD12	1:B:215:ILE:HG13	1.82	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:LYS:HB3	1:B:104:ASP:HB2	1.83	0.61
1:B:48:ARG:O	1:B:49:LEU:HB2	2.00	0.61
1:A:223:LYS:NZ	4:A:1434:HOH:O	2.35	0.59
1:B:79:VAL:HA	1:B:117:VAL:HG21	1.86	0.58
1:B:195:LEU:HB2	1:B:231:SER:HB3	1.86	0.57
1:A:100:LYS:HB3	1:A:104:ASP:HB2	1.87	0.57
1:A:56:PRO:HG2	1:A:134:ASN:HB2	1.86	0.56
1:A:208:ASP:OD2	1:A:212:ARG:HB2	2.06	0.56
1:B:194:ILE:CD1	1:B:215:ILE:HG13	2.34	0.56
1:B:194:ILE:CD1	1:B:225:VAL:HG21	2.33	0.56
1:B:48:ARG:C	1:B:50:ASN:H	2.15	0.54
1:A:101:TRP:CE2	1:A:102:PRO:HB3	2.42	0.54
1:B:56:PRO:HG2	1:B:134:ASN:HB2	1.88	0.54
1:B:59:GLY:HA3	1:B:61:TRP:CZ2	2.44	0.53
1:B:186:MET:HE1	1:B:207:ILE:CG2	2.39	0.53
1:B:103:ASN:HB2	1:B:115:VAL:HB	1.92	0.52
1:A:179:LEU:CD1	1:A:209:ASP:HA	2.39	0.52
1:A:222:VAL:HG11	4:A:1628:HOH:O	2.10	0.52
1:B:84:VAL:O	1:B:88:GLU:HG3	2.09	0.52
1:B:74:ASP:HA	1:B:77:LYS:HZ3	1.74	0.51
1:A:116:LEU:HA	4:A:1491:HOH:O	2.11	0.51
1:B:91:LYS:HE2	4:B:1598:HOH:O	2.10	0.51
1:B:11:ARG:HB2	1:B:11:ARG:NH1	2.08	0.50
1:B:177:ASP:O	1:B:180:ASN:ND2	2.44	0.50
1:A:24:GLU:O	1:A:28:THR:HG22	2.11	0.49
1:B:75:LEU:HG	1:B:124:ILE:HD11	1.94	0.49
1:B:80:PHE:HB3	1:B:182:VAL:HG21	1.94	0.48
1:B:42:GLN:HB2	1:B:55:SER:HB2	1.96	0.48
1:A:108:ASN:ND2	4:A:1416:HOH:O	2.23	0.48
1:A:180:ASN:ND2	4:A:1681:HOH:O	2.47	0.47
1:B:144:LYS:HE3	4:B:1606:HOH:O	2.12	0.47
1:A:154:LEU:HD22	1:B:154:LEU:HD22	1.95	0.47
1:A:84:VAL:O	1:A:88:GLU:HG3	2.13	0.47
1:B:158:ARG:HH11	1:B:158:ARG:HG3	1.80	0.47
1:B:80:PHE:HB3	1:B:182:VAL:CG2	2.45	0.47
1:B:11:ARG:NH1	4:B:1467:HOH:O	2.43	0.47
1:B:48:ARG:C	1:B:50:ASN:N	2.72	0.46
1:A:27:LYS:C	1:A:29:SER:N	2.70	0.46
1:A:48:ARG:NH2	1:A:230:VAL:O	2.47	0.46
1:B:194:ILE:HD11	1:B:215:ILE:HD11	1.97	0.46
1:A:216:ARG:HG3	1:A:222:VAL:HG12	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:TRP:CE2	1:B:102:PRO:HB3	2.51	0.46
1:B:152:PRO:HG2	1:B:155:SER:OG	2.16	0.46
1:A:191:ARG:NH1	4:A:1669:HOH:O	2.48	0.45
1:A:212:ARG:HG2	1:A:226:ILE:HG22	1.98	0.45
1:B:48:ARG:O	4:B:1452:HOH:O	2.21	0.45
1:B:77:LYS:HE3	1:B:209:ASP:O	2.17	0.45
1:A:176:MET:HE3	1:A:209:ASP:O	2.16	0.45
1:A:30:TYR:O	1:A:30:TYR:CD2	2.70	0.45
1:B:42:GLN:HB2	1:B:55:SER:CB	2.47	0.44
1:A:30:TYR:CD1	1:A:30:TYR:N	2.77	0.44
1:A:68:PRO:HG2	1:A:75:LEU:HD11	2.00	0.44
1:A:232:LEU:HB2	4:A:1577:HOH:O	2.17	0.44
1:A:73:LYS:NZ	4:A:1553:HOH:O	2.51	0.44
1:A:79:VAL:HA	1:A:117:VAL:HG21	1.99	0.44
1:A:206:ASP:OD1	1:A:207:ILE:N	2.46	0.43
1:A:214:ILE:HD13	1:A:224:LYS:HG2	2.00	0.43
1:A:232:LEU:CB	4:A:1577:HOH:O	2.66	0.43
1:B:51:ARG:HD2	2:B:1302:ADP:O3'	2.17	0.43
1:B:226:ILE:HG22	1:B:227:TYR:N	2.33	0.43
1:A:62:LEU:C	1:A:62:LEU:HD12	2.44	0.43
1:A:186:MET:HE1	1:A:207:ILE:CG2	2.46	0.43
1:A:225:VAL:HG11	1:A:230:VAL:HG11	2.00	0.42
1:B:173:LYS:NZ	4:B:1462:HOH:O	2.36	0.42
1:A:223:LYS:HD2	4:A:1689:HOH:O	2.19	0.42
1:A:27:LYS:O	1:A:29:SER:N	2.52	0.42
1:A:103:ASN:HB2	1:A:115:VAL:HB	2.02	0.42
1:B:232:LEU:HD23	1:B:232:LEU:C	2.44	0.42
1:A:19:ILE:O	1:A:42:GLN:HA	2.20	0.42
1:A:101:TRP:HA	1:A:102:PRO:HA	1.70	0.42
1:B:19:ILE:HD11	1:B:25:PHE:HB2	2.02	0.42
1:A:23:ASN:O	1:A:27:LYS:HG3	2.20	0.42
1:B:51:ARG:HD3	4:B:1417:HOH:O	2.19	0.42
1:B:11:ARG:HH11	1:B:11:ARG:CG	2.33	0.41
1:B:62:LEU:C	1:B:62:LEU:HD12	2.46	0.41
1:B:72:GLN:HA	1:B:72:GLN:OE1	2.20	0.41
1:B:210:PHE:HB3	4:B:1635:HOH:O	2.20	0.41
1:A:232:LEU:C	1:A:232:LEU:HD23	2.45	0.41
1:A:75:LEU:HD12	1:A:75:LEU:HA	1.88	0.41
1:B:57:GLU:CD	1:B:57:GLU:C	2.89	0.41
1:B:108:ASN:OD1	4:B:1627:HOH:O	2.22	0.41
1:B:144:LYS:O	1:B:148:GLY:N	2.54	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLN:HB2	1:A:55:SER:HB2	2.03	0.41
1:A:75:LEU:N	1:A:76:PRO:CD	2.84	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/235 (99%)	225 (97%)	7 (3%)	1 (0%)	30	21
1	B	233/235 (99%)	228 (98%)	5 (2%)	0	100	100
All	All	466/470 (99%)	453 (97%)	12 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/205 (100%)	200 (98%)	5 (2%)	43	36
1	B	205/205 (100%)	198 (97%)	7 (3%)	32	23
All	All	410/410 (100%)	398 (97%)	12 (3%)	37	29

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	67	SER
1	A	75	LEU
1	A	102	PRO
1	A	176	MET
1	B	11	ARG
1	B	49	LEU
1	B	50	ASN
1	B	75	LEU
1	B	91	LYS
1	B	177	ASP
1	B	180	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	180	ASN
1	B	50	ASN
1	B	138	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ADP	A	1301	-	19,19,29	2.52	7 (36%)	27,28,45	1.44	4 (14%)
3	BTN	B	1402	-	17,17,17	1.04	1 (5%)	23,23,23	1.30	2 (8%)
3	BTN	A	1401	-	17,17,17	1.15	1 (5%)	23,23,23	1.24	2 (8%)
2	ADP	B	1302	-	19,19,29	2.49	7 (36%)	27,28,45	1.41	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1301	-	-	0/4/17/32	0/3/3/3
3	BTN	B	1402	-	-	2/7/28/28	0/2/2/2
3	BTN	A	1401	-	-	2/7/28/28	0/2/2/2
2	ADP	B	1302	-	-	2/4/17/32	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1302	ADP	O4'-C4'	-7.42	1.28	1.43
2	A	1301	ADP	O4'-C4'	-7.39	1.28	1.43
2	A	1301	ADP	C2-N1	4.22	1.41	1.33
2	B	1302	ADP	C2-N1	4.18	1.41	1.33
3	A	1401	BTN	C2-S1	-3.49	1.77	1.82
2	A	1301	ADP	C1'-N9	3.21	1.55	1.46
2	B	1302	ADP	C1'-N9	3.21	1.55	1.46
3	B	1402	BTN	C2-S1	-3.05	1.77	1.82
2	A	1301	ADP	C5-C6	2.54	1.48	1.41
2	B	1302	ADP	C5-C6	2.36	1.47	1.41
2	A	1301	ADP	C8-N7	2.35	1.36	1.31
2	A	1301	ADP	C4-N3	2.23	1.38	1.34
2	B	1302	ADP	C5-N7	-2.16	1.35	1.39
2	B	1302	ADP	C4-N3	2.14	1.38	1.34
2	A	1301	ADP	C5-N7	-2.09	1.35	1.39
2	B	1302	ADP	C8-N7	2.07	1.35	1.31

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1302	ADP	N3-C2-N1	-3.97	122.57	128.58
2	A	1301	ADP	N3-C2-N1	-3.88	122.70	128.58
3	B	1402	BTN	C5-C6-S1	-3.50	101.25	106.06
3	B	1402	BTN	C2-C4-C5	-3.41	104.73	108.89
3	A	1401	BTN	C5-C6-S1	-3.24	101.61	106.06
2	B	1302	ADP	O3'-C3'-C2'	3.17	117.67	111.43
2	A	1301	ADP	C2-N1-C6	3.09	123.81	118.73
3	A	1401	BTN	C2-C4-C5	-3.02	105.20	108.89
2	A	1301	ADP	O3'-C3'-C2'	3.02	117.37	111.43
2	B	1302	ADP	C2-N1-C6	3.01	123.68	118.73
2	A	1301	ADP	C4'-O4'-C1'	2.66	114.76	107.53
2	B	1302	ADP	C4'-O4'-C1'	2.62	114.66	107.53

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1302	ADP	C2'-C1'-N9-C8
3	A	1401	BTN	C9-C10-C11-O11
3	B	1402	BTN	C9-C10-C11-O11
3	A	1401	BTN	C9-C10-C11-O12
3	B	1402	BTN	C9-C10-C11-O12
2	B	1302	ADP	O4'-C1'-N9-C8


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1302	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	235/235 (100%)	1.00	41 (17%) <b>4</b> <b>4</b>	16, 29, 52, 60	0
1	B	235/235 (100%)	0.92	41 (17%) <b>4</b> <b>4</b>	17, 32, 50, 66	0
All	All	470/470 (100%)	0.96	82 (17%) <b>4</b> <b>4</b>	16, 30, 52, 66	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	TYR	6.6
1	A	119	GLY	6.0
1	A	28	THR	4.7
1	A	219	SER	4.0
1	A	120	LYS	3.8
1	A	148	GLY	3.7
1	B	219	SER	3.7
1	A	49	LEU	3.6
1	A	196	GLY	3.5
1	B	148	GLY	3.4
1	B	217	LEU	3.4
1	B	198	GLY	3.3
1	B	227	TYR	3.2
1	B	78	ILE	3.2
1	A	228	GLY	3.2
1	A	73	LYS	3.1
1	A	176	MET	3.1
1	A	224	LYS	3.1
1	B	121	GLY	3.1
1	B	122	ASP	3.0
1	B	232	LEU	3.0
1	A	227	TYR	3.0
1	B	180	ASN	3.0
1	A	32	GLU	2.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	180	ASN	2.9
1	B	44	MET	2.9
1	B	197	ASP	2.8
1	B	49	LEU	2.8
1	B	195	LEU	2.8
1	A	220	GLY	2.8
1	B	204	ALA	2.7
1	B	210	PHE	2.7
1	B	228	GLY	2.7
1	A	194	ILE	2.6
1	B	223	LYS	2.6
1	A	218	ASP	2.5
1	B	235	LEU	2.5
1	B	222	VAL	2.5
1	A	214	ILE	2.5
1	B	194	ILE	2.5
1	A	75	LEU	2.5
1	A	121	GLY	2.5
1	B	32	GLU	2.5
1	A	210	PHE	2.4
1	B	215	ILE	2.4
1	A	70	VAL	2.4
1	B	179	LEU	2.4
1	B	221	GLU	2.3
1	A	78	ILE	2.3
1	B	225	VAL	2.3
1	B	108	ASN	2.3
1	A	199	SER	2.3
1	A	229	ASP	2.3
1	A	230	VAL	2.3
1	A	50	ASN	2.3
1	B	50	ASN	2.3
1	B	51	ARG	2.3
1	A	231	SER	2.3
1	B	30	TYR	2.2
1	B	200	PHE	2.2
1	A	122	ASP	2.2
1	A	69	LYS	2.2
1	A	77	LYS	2.2
1	B	138	ASN	2.2
1	B	79	VAL	2.2
1	B	57	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	217	LEU	2.1
1	A	225	VAL	2.1
1	A	145	LEU	2.1
1	B	203	ILE	2.1
1	B	226	ILE	2.1
1	B	142	SER	2.1
1	B	101	TRP	2.1
1	B	181	LEU	2.1
1	A	203	ILE	2.0
1	B	231	SER	2.0
1	A	76	PRO	2.0
1	A	222	VAL	2.0
1	B	192	VAL	2.0
1	A	232	LEU	2.0
1	A	200	PHE	2.0
1	A	198	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ADP	B	1302	17/27	0.83	0.15	40,43,55,57	0
2	ADP	A	1301	17/27	0.86	0.13	29,35,53,55	0
3	BTN	A	1401	16/16	0.96	0.07	15,18,27,28	0
3	BTN	B	1402	16/16	0.96	0.07	17,19,25,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.