



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:27 PM UTC

PDB ID : 6DTL / pdb_00006dtl
Title : Mitogen-activated protein kinase 6
Authors : Ruble, J.
Deposited on : 2018-06-17
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	13% 77% 20% .
1	B	364	10% 74% 24% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	0	0
			2896	1851	499	531	15			
1	B	357	Total	C	N	O	S	0	0	0
			2915	1867	497	535	16			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	60	Total	O	0	0
			60	60		
2	B	58	Total	O	0	0
			58	58		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 6



- Molecule 1: Mitogen-activated protein kinase 6



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	151.59Å 151.59Å 85.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.98 – 2.75 42.98 – 2.75	Depositor EDS
% Data completeness (in resolution range)	93.7 (42.98-2.75) 93.7 (42.98-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.05 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.205 , 0.256 0.204 , 0.253	Depositor DCC
R_{free} test set	2002 reflections (6.71%)	wwPDB-VP
Wilson B-factor (Å ²)	43.1	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5929	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/2964	0.44	0/4020
1	B	0.20	0/2985	0.44	2/4049 (0.0%)
All	All	0.20	0/5949	0.44	2/8069 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	LYS	CA-C-N	5.90	126.45	120.38
1	B	63	LYS	C-N-CA	5.90	126.45	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2896	0	2877	51	1
1	B	2915	0	2887	59	1
2	A	60	0	0	3	0
2	B	58	0	0	8	0
All	All	5929	0	5764	110	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230:ARG:NH2	2:B:405:HOH:O	2.11	0.82
1:A:337:ILE:O	2:A:401:HOH:O	1.98	0.80
1:B:277:MET:HE1	1:B:299:ILE:HD12	1.66	0.77
1:A:285:GLU:HA	1:A:288:LEU:HB2	1.69	0.73
1:B:115:ARG:NH1	1:B:385:GLU:OE1	2.22	0.73
1:A:283:PRO:HG2	1:A:288:LEU:HD21	1.70	0.72
1:B:278:GLU:OE1	2:B:402:HOH:O	2.08	0.72
1:B:326:GLU:OE2	2:B:403:HOH:O	2.08	0.71
1:A:163:GLU:OE2	1:A:347:TYR:OH	2.09	0.70
1:B:307:ARG:O	2:B:404:HOH:O	2.11	0.69
1:B:134:ARG:HE	1:B:379:LYS:HE2	1.57	0.68
1:B:131:PRO:O	1:B:393:TYR:OH	2.07	0.66
1:A:212:ARG:HE	1:A:216:GLU:HB3	1.59	0.66
1:B:232:PRO:HG3	1:B:276:LEU:HD23	1.77	0.65
1:B:256:GLU:OE1	2:B:406:HOH:O	2.13	0.65
1:A:258:MET:HE2	1:A:317:VAL:HG13	1.82	0.61
1:A:369:GLU:HG3	1:A:381:LEU:HD11	1.83	0.60
1:B:236:LEU:HD13	1:B:291:LEU:HD21	1.82	0.60
1:A:279:LEU:HD21	1:A:310:ILE:HD11	1.82	0.59
1:A:188:ARG:HD2	1:A:212:ARG:HB2	1.84	0.58
1:A:159:GLN:O	1:A:260:ARG:NH2	2.38	0.57
1:B:134:ARG:NH2	1:B:380:GLU:OE1	2.34	0.57
1:B:104:ALA:O	1:B:108:LEU:HB3	2.05	0.56
1:B:255:MET:SD	1:B:325:ILE:HD13	2.47	0.55
1:B:98:PHE:HZ	1:B:379:LYS:HB2	1.72	0.55
1:B:380:GLU:HG3	1:B:384:ARG:CZ	2.37	0.55
1:B:376:GLU:O	1:B:379:LYS:HB3	2.06	0.54
1:A:288:LEU:HD22	1:A:299:ILE:HD13	1.89	0.54
1:A:189:ASP:O	1:A:194:ASN:ND2	2.40	0.54
1:A:151:LEU:HD11	1:A:257:LEU:HD13	1.89	0.54
1:B:109:ARG:O	1:B:113:LEU:HB2	2.07	0.54
1:A:255:MET:HE2	1:A:325:ILE:HG21	1.89	0.53
1:B:105:LYS:O	1:B:109:ARG:HB2	2.09	0.53
1:B:117:MET:HE1	1:B:180:ILE:HA	1.89	0.53
1:B:135:ASN:OD1	1:B:136:ALA:N	2.42	0.53
1:B:42:HIS:O	1:B:45:ARG:HG2	2.09	0.53
1:A:37:PRO:HG2	1:A:50:ASN:HB3	1.90	0.52
1:A:87:GLU:OE1	2:A:402:HOH:O	2.19	0.52
1:B:45:ARG:HG3	1:B:46:PHE:HD1	1.74	0.51
1:A:330:THR:O	1:A:336:ARG:NH1	2.44	0.51
1:B:68:PRO:HA	1:B:78:CYS:HA	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:ARG:HG3	1:B:46:PHE:CD1	2.46	0.50
1:B:222:GLU:OE1	2:B:408:HOH:O	2.20	0.49
1:B:55:ILE:O	1:B:93:LYS:NZ	2.45	0.49
1:B:112:LYS:HB3	1:B:364:PHE:HZ	1.78	0.49
1:A:293:GLU:HA	1:A:296:LYS:HB2	1.94	0.49
1:A:40:LEU:HD23	1:A:47:ILE:HG12	1.95	0.48
1:A:273:LEU:O	1:A:277:MET:HG2	2.13	0.48
1:B:127:ASP:HB3	1:B:143:ALA:HB3	1.96	0.48
1:B:130:PRO:HD3	1:B:390:ASN:OD1	2.14	0.47
1:B:305:TYR:OH	2:B:407:HOH:O	2.19	0.47
1:B:105:LYS:HG3	1:B:378:MET:HE1	1.96	0.47
1:A:285:GLU:HA	1:A:288:LEU:CB	2.40	0.46
1:B:178:LYS:HE2	1:B:359:GLU:HG3	1.97	0.46
1:A:197:LEU:HD21	1:A:203:LEU:HD12	1.97	0.46
1:A:103:ASP:O	1:A:106:ARG:HG2	2.16	0.46
1:A:227:ARG:NE	1:A:269:HIS:HA	2.30	0.46
1:B:98:PHE:HZ	1:B:379:LYS:CB	2.29	0.46
1:A:320:LEU:HD13	1:A:346:PRO:HB2	1.96	0.46
1:B:220:MET:O	1:B:224:VAL:HG22	2.16	0.46
1:A:33:ILE:HG12	1:A:65:PRO:HD3	1.98	0.46
1:A:117:MET:HB3	1:A:179:TYR:HE1	1.81	0.46
1:A:227:ARG:HE	1:A:269:HIS:HA	1.80	0.46
1:B:104:ALA:O	1:B:108:LEU:CB	2.64	0.46
1:A:127:ASP:HB3	1:A:143:ALA:HB3	1.98	0.45
1:A:288:LEU:CD2	1:A:299:ILE:HD13	2.46	0.45
1:A:245:ILE:HG13	1:A:246:ASP:N	2.31	0.45
1:A:255:MET:HE2	1:A:325:ILE:HD13	1.97	0.45
1:A:272:GLN:O	1:A:276:LEU:HG	2.16	0.45
1:B:310:ILE:HG22	1:B:322:ILE:HD12	1.99	0.45
1:B:220:MET:SD	1:B:224:VAL:HG21	2.56	0.45
1:A:346:PRO:HA	1:A:349:ASN:CG	2.42	0.45
1:B:126:ARG:CZ	1:B:145:GLU:HG2	2.47	0.45
1:B:155:ILE:HD13	1:B:256:GLU:HG2	1.99	0.45
1:B:277:MET:HE2	1:B:331:PHE:CE2	2.51	0.45
1:A:227:ARG:HH21	1:A:268:ASP:C	2.25	0.44
1:A:292:ASN:O	1:A:296:LYS:N	2.40	0.44
1:A:67:MET:HE3	1:A:68:PRO:HD2	2.00	0.44
1:A:285:GLU:CD	1:A:300:ARG:HH22	2.25	0.43
1:A:288:LEU:HD21	1:A:299:ILE:HG21	1.99	0.43
1:A:188:ARG:CD	1:A:212:ARG:HB2	2.49	0.43
1:B:230:ARG:HD3	1:B:234:LEU:HB3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181:HIS:CE1	1:B:243:ALA:HB1	2.54	0.42
1:A:298:TYR:O	1:A:301:GLN:HB3	2.19	0.42
1:B:189:ASP:O	1:B:194:ASN:ND2	2.52	0.42
1:B:349:ASN:HB3	2:B:413:HOH:O	2.18	0.42
1:A:33:ILE:H	1:A:33:ILE:HG13	1.65	0.42
1:A:267:ARG:HH21	1:A:274:ARG:HH22	1.67	0.42
1:B:129:ILE:HB	1:B:141:TYR:HB2	2.00	0.42
1:B:94:ILE:HG22	1:B:97:ALA:HB3	2.00	0.42
1:B:96:ASN:HA	1:B:138:ASN:O	2.19	0.42
1:B:289:GLU:HB2	1:B:334:ARG:HH12	1.85	0.42
1:B:56:PHE:CE1	1:B:93:LYS:HB2	2.55	0.42
1:B:70:GLY:HA3	1:B:77:VAL:HB	2.02	0.42
1:B:295:ALA:O	1:B:299:ILE:HG12	2.20	0.42
1:A:380:GLU:O	1:A:384:ARG:HG2	2.19	0.42
1:B:234:LEU:HA	1:B:241:TYR:OH	2.18	0.42
1:A:231:ALA:HA	1:A:248:TRP:CD1	2.54	0.42
1:B:113:LEU:O	1:B:117:MET:HG2	2.20	0.42
1:B:272:GLN:O	1:B:276:LEU:HD13	2.19	0.42
1:A:267:ARG:HH21	1:A:274:ARG:NH2	2.17	0.41
1:A:181:HIS:CE1	1:A:243:ALA:HB1	2.56	0.41
1:B:244:ALA:HB1	1:B:336:ARG:HH11	1.85	0.41
1:B:294:ASN:OD1	1:B:297:ARG:NH2	2.52	0.41
1:B:123:VAL:HG23	1:B:205:ILE:O	2.21	0.41
1:B:109:ARG:HH22	1:B:366:PHE:HE1	1.68	0.41
1:A:283:PRO:HG3	1:A:331:PHE:CD1	2.56	0.41
1:A:248:TRP:HE3	1:A:336:ARG:NH1	2.19	0.40
1:A:302:LEU:HD23	1:A:303:PRO:HD2	2.02	0.40
1:A:310:ILE:N	2:A:418:HOH:O	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:ARG:NH2	1:B:304:PRO:O[6_455]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/364 (97%)	335 (95%)	17 (5%)	0	100	100
1	B	353/364 (97%)	333 (94%)	19 (5%)	1 (0%)	36	56
All	All	705/728 (97%)	668 (95%)	36 (5%)	1 (0%)	48	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	394	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/327 (98%)	317 (99%)	2 (1%)	78	88
1	B	322/327 (98%)	320 (99%)	2 (1%)	78	88
All	All	641/654 (98%)	637 (99%)	4 (1%)	78	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	368	PHE
1	A	369	GLU
1	B	234	LEU
1	B	269	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	121	ASN
1	B	301	GLN
1	B	308	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/364 (97%)	0.50	49 (13%) 6 5	13, 49, 123, 143	0
1	B	357/364 (98%)	0.33	38 (10%) 11 10	15, 44, 115, 146	0
All	All	713/728 (97%)	0.42	87 (12%) 8 7	13, 47, 121, 146	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	102	ILE	6.8
1	B	74	TYR	6.2
1	A	272	GLN	5.1
1	A	291	LEU	5.1
1	A	213	VAL	4.9
1	B	98	PHE	4.9
1	B	97	ALA	4.8
1	A	225	VAL	4.2
1	A	288	LEU	4.1
1	B	210	LEU	4.0
1	B	104	ALA	4.0
1	B	209	GLY	4.0
1	B	34	GLU	4.0
1	A	318	HIS	4.0
1	A	302	LEU	3.9
1	A	33	ILE	3.8
1	B	221	THR	3.8
1	A	280	ILE	3.6
1	A	299	ILE	3.6
1	A	369	GLU	3.6
1	A	290	PHE	3.6
1	B	381	LEU	3.5
1	B	33	ILE	3.3
1	B	241	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	103	ASP	3.3
1	B	72	GLY	3.2
1	A	270	VAL	3.2
1	A	317	VAL	3.2
1	B	133	LEU	3.2
1	A	277	MET	3.1
1	B	76	ILE	3.0
1	B	384	ARG	3.0
1	A	283	PRO	2.9
1	A	364	PHE	2.9
1	A	236	LEU	2.9
1	A	279	LEU	2.9
1	B	100	ASN	2.9
1	A	276	LEU	2.9
1	A	333	PRO	2.9
1	A	214	THR	2.9
1	B	368	PHE	2.9
1	B	71	LYS	2.8
1	A	295	ALA	2.8
1	A	34	GLU	2.8
1	A	298	TYR	2.8
1	A	366	PHE	2.8
1	B	105	LYS	2.7
1	B	95	ALA	2.7
1	B	75	GLY	2.6
1	A	32	GLY	2.6
1	B	101	LYS	2.6
1	A	287	GLU	2.6
1	A	307	ARG	2.6
1	A	306	PRO	2.6
1	A	331	PHE	2.5
1	A	303	PRO	2.5
1	A	73	ALA	2.4
1	A	372	ALA	2.4
1	B	52	PHE	2.4
1	B	73	ALA	2.4
1	A	392	GLU	2.4
1	B	223	TYR	2.4
1	A	235	LEU	2.3
1	B	138	ASN	2.3
1	B	106	ARG	2.3
1	B	366	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	373	LEU	2.3
1	A	293	GLU	2.3
1	B	55	ILE	2.3
1	B	136	ALA	2.3
1	B	378	MET	2.2
1	A	349	ASN	2.2
1	A	316	THR	2.2
1	A	72	GLY	2.2
1	A	232	PRO	2.2
1	A	337	ILE	2.1
1	A	314	PHE	2.1
1	A	237	ASN	2.1
1	A	274	ARG	2.1
1	A	300	ARG	2.1
1	A	334	ARG	2.1
1	B	387	LEU	2.1
1	B	217	SER	2.0
1	B	239	SER	2.0
1	A	71	LYS	2.0
1	B	391	PRO	2.0
1	A	368	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.