



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:43 PM UTC

PDB ID : 7DT2 / pdb_00007dt2
Title : Strategic design of catalytic lysine-targeting reversible covalent BCR-ABL Inhibitors
Authors : Anantharajan, J.; Baburajendran, N.
Deposited on : 2021-01-04
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	4% 82% 15%
1	B	310	14% 78% 7% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	0
			2110	1363	343	388	16			
1	B	267	Total	C	N	O	S	0	0	0
			2120	1367	349	387	17			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	201	MET	-	initiating methionine	UNP P00519
A	202	GLY	-	expression tag	UNP P00519
A	203	SER	-	expression tag	UNP P00519
A	204	SER	-	expression tag	UNP P00519
A	205	HIS	-	expression tag	UNP P00519
A	206	HIS	-	expression tag	UNP P00519
A	207	HIS	-	expression tag	UNP P00519
A	208	HIS	-	expression tag	UNP P00519
A	209	HIS	-	expression tag	UNP P00519
A	210	HIS	-	expression tag	UNP P00519
A	211	SER	-	expression tag	UNP P00519
A	212	SER	-	expression tag	UNP P00519
A	213	GLY	-	expression tag	UNP P00519
A	214	LEU	-	expression tag	UNP P00519
A	215	VAL	-	expression tag	UNP P00519
A	216	PRO	-	expression tag	UNP P00519
A	217	ARG	-	expression tag	UNP P00519
A	218	GLY	-	expression tag	UNP P00519
A	219	SER	-	expression tag	UNP P00519
A	220	HIS	-	expression tag	UNP P00519
A	221	MET	-	expression tag	UNP P00519
A	222	GLU	-	expression tag	UNP P00519
A	223	ASN	-	expression tag	UNP P00519
A	224	LEU	-	expression tag	UNP P00519
A	225	TYR	-	expression tag	UNP P00519

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	226	PHE	-	expression tag	UNP P00519
A	227	GLN	-	expression tag	UNP P00519
A	228	SER	-	expression tag	UNP P00519
B	201	MET	-	initiating methionine	UNP P00519
B	202	GLY	-	expression tag	UNP P00519
B	203	SER	-	expression tag	UNP P00519
B	204	SER	-	expression tag	UNP P00519
B	205	HIS	-	expression tag	UNP P00519
B	206	HIS	-	expression tag	UNP P00519
B	207	HIS	-	expression tag	UNP P00519
B	208	HIS	-	expression tag	UNP P00519
B	209	HIS	-	expression tag	UNP P00519
B	210	HIS	-	expression tag	UNP P00519
B	211	SER	-	expression tag	UNP P00519
B	212	SER	-	expression tag	UNP P00519
B	213	GLY	-	expression tag	UNP P00519
B	214	LEU	-	expression tag	UNP P00519
B	215	VAL	-	expression tag	UNP P00519
B	216	PRO	-	expression tag	UNP P00519
B	217	ARG	-	expression tag	UNP P00519
B	218	GLY	-	expression tag	UNP P00519
B	219	SER	-	expression tag	UNP P00519
B	220	HIS	-	expression tag	UNP P00519
B	221	MET	-	expression tag	UNP P00519
B	222	GLU	-	expression tag	UNP P00519
B	223	ASN	-	expression tag	UNP P00519
B	224	LEU	-	expression tag	UNP P00519
B	225	TYR	-	expression tag	UNP P00519
B	226	PHE	-	expression tag	UNP P00519
B	227	GLN	-	expression tag	UNP P00519
B	228	SER	-	expression tag	UNP P00519

- Molecule 2 is [4-[5-[5-(dimethylcarbamoyl)pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridin-3-yl]-2-methanoyl-5-methoxy-phenyl]boronic acid (CCD ID: HJ9) (formula: C₂₃H₂₁BN₄O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	B	C	N	O	0	0
			32	1	23	4	4		
2	B	1	Total	B	C	N	O	0	0
			32	1	23	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	108	Total	O	0	0
			108	108		
3	B	56	Total	O	0	0
			56	56		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase ABL1



• Molecule 1: Tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.69Å 132.30Å 56.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.08 – 2.30 49.08 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.08-2.30) 99.9 (49.08-2.30)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, R_{free}	0.206 , 0.237 0.214 , 0.245	Depositor DCC
R_{free} test set	2000 reflections (5.57%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4458	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HJ9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	2/2164 (0.1%)	0.72	5/2936 (0.2%)
1	B	0.29	2/2172 (0.1%)	0.70	6/2943 (0.2%)
All	All	0.29	4/4336 (0.1%)	0.71	11/5879 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	270	VAL	C-N	7.51	1.43	1.33
1	B	271	LYS	C-N	7.43	1.43	1.33
1	A	271	LYS	C-N	7.23	1.43	1.33
1	B	270	VAL	C-N	6.99	1.43	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	VAL	O-C-N	20.38	144.80	123.20
1	A	270	VAL	CA-C-N	-18.37	97.65	123.00
1	A	270	VAL	C-N-CA	-18.37	97.65	123.00
1	B	271	LYS	O-C-N	16.52	144.71	123.23
1	B	270	VAL	CA-C-N	-14.82	100.71	122.94
1	B	270	VAL	C-N-CA	-14.82	100.71	122.94
1	B	270	VAL	O-C-N	13.11	137.10	123.20
1	B	271	LYS	CA-C-N	-10.45	104.33	123.03
1	B	271	LYS	C-N-CA	-10.45	104.33	123.03
1	A	271	LYS	CA-C-N	-6.76	111.46	122.82
1	A	271	LYS	C-N-CA	-6.76	111.46	122.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2110	0	2029	6	0
1	B	2120	0	2052	15	0
2	A	32	0	0	0	0
2	B	32	0	0	0	0
3	A	108	0	0	0	0
3	B	56	0	0	0	0
All	All	4458	0	4081	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:VAL:HG22	1:B:314:ILE:HG12	1.73	0.69
1:A:327:LEU:HD21	1:A:343:MET:HE1	1.81	0.63
1:A:459:GLU:CD	1:A:459:GLU:H	2.08	0.61
1:B:301:LEU:HD11	1:B:304:VAL:HG23	1.90	0.54
1:B:361:HIS:HA	1:B:384:LEU:HD23	1.93	0.50
1:B:273:LEU:HD21	1:B:280:VAL:HG22	1.94	0.50
1:B:304:VAL:CG1	1:B:306:THR:HG23	2.42	0.48
1:B:361:HIS:NE2	1:B:380:ALA:O	2.43	0.48
1:B:292:GLU:CD	1:B:386:ARG:HH22	2.23	0.47
1:A:242:ILE:HG21	1:A:270:VAL:HG21	1.97	0.47
1:A:286:GLU:OE2	1:A:382:PHE:HB3	2.15	0.45
1:B:381:ASP:OD2	1:B:381:ASP:N	2.50	0.44
1:B:241:ASP:CG	1:B:261:TRP:HE1	2.25	0.44
1:A:306:THR:HA	1:A:311:PHE:HD2	1.82	0.43
1:B:355:GLU:HG3	1:B:418:ILE:HG12	2.00	0.43
1:B:236:GLU:HG3	1:B:307:ARG:HE	1.83	0.43
1:B:243:THR:OG1	1:B:262:LYS:NZ	2.47	0.42
1:B:460:ARG:HB2	1:B:469:TYR:CG	2.55	0.41
1:B:418:ILE:HD12	1:B:418:ILE:HA	1.94	0.41
1:A:285:LYS:HA	1:B:498:GLN:HG2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/310 (84%)	253 (97%)	8 (3%)	0	100	100
1	B	263/310 (85%)	253 (96%)	10 (4%)	0	100	100
All	All	524/620 (84%)	506 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/273 (80%)	218 (100%)	1 (0%)	81	90
1	B	219/273 (80%)	218 (100%)	1 (0%)	81	90
All	All	438/546 (80%)	436 (100%)	2 (0%)	81	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	271	LYS
1	B	271	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	491	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HJ9	B	601	1	35,35,36	2.55	7 (20%)	49,51,52	1.68	13 (26%)
2	HJ9	A	601	1	35,35,36	2.51	7 (20%)	49,51,52	1.75	13 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HJ9	B	601	1	-	0/22/22/24	0/4/4/4
2	HJ9	A	601	1	-	0/22/22/24	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	HJ9	C02-N03	10.29	1.49	1.34
2	A	601	HJ9	C02-N03	10.11	1.49	1.34
2	B	601	HJ9	C18-C17	5.96	1.59	1.49
2	A	601	HJ9	C18-C17	5.95	1.59	1.49
2	B	601	HJ9	C06-C02	5.20	1.58	1.50
2	A	601	HJ9	C06-C02	4.93	1.58	1.50
2	B	601	HJ9	C11-C10	3.73	1.57	1.49
2	A	601	HJ9	C11-C10	3.60	1.57	1.49
2	B	601	HJ9	C16-C17	2.96	1.40	1.37
2	A	601	HJ9	O01-C02	-2.82	1.16	1.22
2	B	601	HJ9	O01-C02	-2.80	1.16	1.22
2	A	601	HJ9	C16-C17	2.79	1.40	1.37
2	B	601	HJ9	O38-C37	2.17	1.40	1.37
2	A	601	HJ9	O38-C37	2.13	1.40	1.37

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HJ9	C40-C14-N13	-3.94	121.53	126.31
2	B	601	HJ9	C40-C14-N13	-3.91	121.58	126.31
2	A	601	HJ9	C14-C40-C17	3.68	107.50	106.22
2	A	601	HJ9	C14-N15-C16	3.42	110.09	108.48
2	B	601	HJ9	B33-C32-C20	-3.36	122.21	125.34
2	B	601	HJ9	C14-N15-C16	3.03	109.91	108.48
2	B	601	HJ9	C14-C40-C17	3.02	107.27	106.22
2	B	601	HJ9	C12-N13-C14	2.70	121.48	116.19
2	A	601	HJ9	C09-N08-C07	2.68	121.15	117.51
2	B	601	HJ9	C09-N08-C07	2.67	121.14	117.51
2	A	601	HJ9	C41-C11-C12	2.64	119.93	117.10
2	A	601	HJ9	B33-C32-C20	-2.63	122.89	125.34
2	A	601	HJ9	C12-N13-C14	2.60	121.29	116.19
2	A	601	HJ9	O38-C37-C18	2.52	120.21	116.55
2	B	601	HJ9	C41-C11-C12	2.49	119.77	117.10
2	B	601	HJ9	O38-C37-C18	2.41	120.06	116.55
2	A	601	HJ9	C11-C12-N13	-2.39	120.48	124.28
2	A	601	HJ9	N15-C14-N13	2.38	130.55	126.69
2	B	601	HJ9	C11-C12-N13	-2.37	120.52	124.28
2	A	601	HJ9	O38-C37-C36	-2.31	120.10	124.08
2	A	601	HJ9	C10-C42-C06	-2.24	118.64	121.04
2	B	601	HJ9	O38-C37-C36	-2.20	120.28	124.08
2	B	601	HJ9	C10-C42-C06	-2.18	118.70	121.04
2	B	601	HJ9	N15-C14-N13	2.18	130.21	126.69
2	B	601	HJ9	C06-C02-N03	2.17	121.47	119.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HJ9	C42-C10-C11	-2.05	117.41	120.84


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/310 (85%)	0.16	11 (4%) 40 42	23, 37, 57, 74	0
1	B	267/310 (86%)	0.85	44 (16%) 4 5	28, 46, 85, 93	0
All	All	532/620 (85%)	0.51	55 (10%) 12 13	23, 41, 75, 93	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	444	ASP	5.8
1	B	275	GLU	5.7
1	B	277	THR	4.5
1	B	240	THR	4.2
1	A	274	LYS	4.1
1	B	246	HIS	4.1
1	B	311	PHE	4.0
1	B	242	ILE	4.0
1	B	243	THR	3.6
1	B	445	LEU	3.2
1	B	250	GLY	3.2
1	B	261	TRP	3.2
1	B	238	GLU	3.1
1	B	390	GLY	3.1
1	B	257	TYR	2.8
1	B	244	MET	2.7
1	B	398	GLY	2.7
1	B	310	PRO	2.7
1	A	282	GLU	2.7
1	B	262	LYS	2.6
1	B	274	LYS	2.6
1	B	362	ARG	2.6
1	B	237	MET	2.6
1	B	251	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	264	TYR	2.6
1	B	278	MET	2.6
1	B	446	SER	2.6
1	B	306	THR	2.6
1	A	246	HIS	2.6
1	A	249	GLY	2.5
1	B	272	THR	2.5
1	B	280	VAL	2.5
1	B	273	LEU	2.5
1	B	233	ASP	2.4
1	B	383	GLY	2.4
1	A	278	MET	2.4
1	B	307	ARG	2.4
1	B	279	GLU	2.4
1	A	308	GLU	2.3
1	B	397	ALA	2.3
1	B	309	PRO	2.3
1	B	293	ILE	2.3
1	B	391	ASP	2.2
1	A	470	GLU	2.2
1	A	264	TYR	2.1
1	A	390	GLY	2.1
1	B	413	TYR	2.1
1	B	286	GLU	2.1
1	A	237	MET	2.1
1	B	360	ILE	2.1
1	B	396	HIS	2.1
1	B	401	PHE	2.1
1	B	282	GLU	2.0
1	B	266	LEU	2.0
1	A	309	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HJ9	B	601	32/33	0.90	0.12	46,52,66,67	0
2	HJ9	A	601	32/33	0.91	0.12	28,40,61,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HJ9 B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HJ9 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.