



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 01:26 AM UTC

PDB ID : 3DUH / pdb_00003duh
Title : Structure of Interleukin-23
Authors : Lupardus, P.J.; Garcia, K.C.
Deposited on : 2008-07-17
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	4% 79% 15% • •
1	B	314	6% 75% 17% • 6%
2	C	177	4% 66% 10% • 23%
2	D	177	9% 59% 16% • 23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	313	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	9	0	0
			2355	1488	385	470	12			
1	B	294	Total	C	N	O	S	6	0	0
			2314	1467	373	462	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	LEU	-	expression tag	UNP P29460
A	0	GLU	-	expression tag	UNP P29460
A	307	HIS	-	expression tag	UNP P29460
A	308	HIS	-	expression tag	UNP P29460
A	309	HIS	-	expression tag	UNP P29460
A	310	HIS	-	expression tag	UNP P29460
A	311	HIS	-	expression tag	UNP P29460
A	312	HIS	-	expression tag	UNP P29460
B	-1	LEU	-	expression tag	UNP P29460
B	0	GLU	-	expression tag	UNP P29460
B	307	HIS	-	expression tag	UNP P29460
B	308	HIS	-	expression tag	UNP P29460
B	309	HIS	-	expression tag	UNP P29460
B	310	HIS	-	expression tag	UNP P29460
B	311	HIS	-	expression tag	UNP P29460
B	312	HIS	-	expression tag	UNP P29460

- Molecule 2 is a protein called Interleukin-23 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	137	Total	C	N	O	S	0	0	0
			1049	669	192	183	5			
2	D	137	Total	C	N	O	S	0	0	0
			1064	676	198	185	5			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	LEU	-	expression tag	UNP Q9NPF7
C	171	HIS	-	expression tag	UNP Q9NPF7
C	172	HIS	-	expression tag	UNP Q9NPF7
C	173	HIS	-	expression tag	UNP Q9NPF7
C	174	HIS	-	expression tag	UNP Q9NPF7
C	175	HIS	-	expression tag	UNP Q9NPF7
C	176	HIS	-	expression tag	UNP Q9NPF7
D	0	LEU	-	expression tag	UNP Q9NPF7
D	171	HIS	-	expression tag	UNP Q9NPF7
D	172	HIS	-	expression tag	UNP Q9NPF7
D	173	HIS	-	expression tag	UNP Q9NPF7
D	174	HIS	-	expression tag	UNP Q9NPF7
D	175	HIS	-	expression tag	UNP Q9NPF7
D	176	HIS	-	expression tag	UNP Q9NPF7

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	96	Total 96	O 96	0	0
4	B	77	Total 77	O 77	0	0
4	C	20	Total 20	O 20	0	0
4	D	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-12 subunit beta




- Molecule 1: Interleukin-12 subunit beta



- Molecule 2: Interleukin-23 subunit alpha



- Molecule 2: Interleukin-23 subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.09Å 59.97Å 160.37Å 90.00° 90.46° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 30.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.30) 99.9 (30.00-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.268 0.227 , 0.267	Depositor DCC
R_{free} test set	2502 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7010	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	1/2413 (0.0%)	0.93	7/3285 (0.2%)
1	B	0.67	1/2370 (0.0%)	0.86	0/3225
2	C	0.76	0/1073	0.97	0/1454
2	D	0.62	0/1090	1.05	3/1477 (0.2%)
All	All	0.70	2/6946 (0.0%)	0.93	10/9441 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	101	PRO	N-CD	6.67	1.57	1.47
1	B	92	THR	CA-CB	5.50	1.57	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	212	LYS	CA-C-N	-8.37	112.18	120.21
1	A	212	LYS	C-N-CA	-8.37	112.18	120.21
1	A	41	ASP	CB-CA-C	-7.61	105.69	116.34
1	A	100	GLU	N-CA-C	7.04	125.36	109.81
2	D	8	SER	CA-C-N	6.65	128.15	119.84
2	D	8	SER	C-N-CA	6.65	128.15	119.84
1	A	58	LYS	N-CA-C	-5.63	108.20	114.62
2	D	7	SER	CB-CA-C	5.36	118.27	109.90
1	A	215	PRO	O-C-N	5.20	123.70	121.31
1	A	40	LEU	CB-CA-C	-5.19	101.78	110.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2355	0	2224	27	0
1	B	2314	0	2190	34	0
2	C	1049	0	1031	14	0
2	D	1064	0	1050	17	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
4	A	96	0	0	1	0
4	B	77	0	0	2	0
4	C	20	0	0	0	0
4	D	7	0	0	1	0
All	All	7010	0	6521	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:LYS:O	1:B:105:THR:HG22	1.51	1.10
2:D:51:HIS:O	2:D:73:ARG:NH1	1.89	1.06
1:B:100:GLU:CB	1:B:101:PRO:HD3	1.95	0.96
1:A:32:GLU:HB2	4:A:410:HOH:O	1.71	0.90
1:B:102:LYS:O	1:B:105:THR:CG2	2.21	0.87
1:A:0:GLU:HB3	1:A:12:GLU:HB2	1.59	0.82
1:B:97:ASP:OD1	1:B:97:ASP:C	2.30	0.74
2:D:138:GLN:HG3	2:D:139:ARG:H	1.51	0.74
1:B:100:GLU:CB	1:B:101:PRO:CD	2.66	0.73
2:D:8:SER:HB2	2:D:9:PRO:HD2	1.74	0.67
2:C:93:GLU:CB	2:C:94:PRO:HD2	2.27	0.65
2:C:89:ILE:O	2:C:143:ARG:HD2	1.97	0.64
2:C:141:LEU:O	2:C:145:LYS:HG2	1.98	0.63
2:C:21:LEU:HB3	2:C:154:VAL:HG21	1.81	0.61
2:C:0:LEU:HD12	2:C:0:LEU:N	2.14	0.61
1:A:263:LYS:O	1:A:279:ARG:NH2	2.34	0.60
1:A:133:SER:HB3	1:A:191:ASP:HB2	1.85	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:VAL:HB	1:A:265:ASP:HB2	1.85	0.58
2:C:137:TRP:HB3	2:C:139:ARG:HG3	1.86	0.58
1:A:177:CYS:SG	1:A:180:ALA:HB2	2.45	0.56
1:A:270:ASP:OD1	1:A:291:ARG:HD2	2.06	0.56
1:B:30:THR:O	1:B:51:LYS:HE3	2.05	0.56
2:C:89:ILE:HG22	2:C:143:ARG:HG3	1.89	0.55
2:D:138:GLN:HG3	2:D:139:ARG:N	2.21	0.55
2:D:90:PHE:CE1	2:D:143:ARG:HB2	2.42	0.55
1:B:285:SER:HB3	1:B:300:TRP:CE3	2.43	0.54
1:A:100:GLU:CB	1:A:101:PRO:HD3	2.39	0.53
1:A:156:GLU:OE2	1:A:165:TYR:CE2	2.62	0.53
1:B:256:GLN:HG3	1:B:264:LYS:HB3	1.91	0.52
1:B:186:ILE:O	1:B:204:SER:HA	2.10	0.52
1:B:63:ALA:HB1	1:B:80:LEU:O	2.10	0.52
1:A:127:SER:HB3	1:A:165:TYR:CZ	2.45	0.51
2:C:0:LEU:HD12	2:C:0:LEU:H3	1.73	0.50
1:B:97:ASP:HB2	1:B:106:PHE:CE2	2.47	0.50
1:A:156:GLU:OE2	1:A:165:TYR:HE2	1.95	0.50
1:A:107:LEU:HD12	1:A:203:SER:HB2	1.92	0.50
2:D:11:TRP:CZ3	2:D:164:GLY:HA3	2.46	0.50
2:D:137:TRP:HA	2:D:140:LEU:HB2	1.94	0.50
1:B:63:ALA:HB3	1:B:196:LEU:HD13	1.94	0.50
1:A:228:ARG:HH11	1:A:228:ARG:H	1.57	0.49
1:A:255:VAL:HG22	1:A:284:ILE:HD12	1.93	0.49
2:D:8:SER:CB	2:D:9:PRO:HD2	2.41	0.49
1:B:229:GLN:HE21	1:B:277:ILE:HD11	1.78	0.49
2:C:21:LEU:CB	2:C:154:VAL:HG21	2.43	0.48
1:A:63:ALA:HB1	1:A:80:LEU:O	2.13	0.48
2:D:22:CYS:SG	2:D:154:VAL:HG12	2.54	0.47
1:B:280:LYS:HD3	1:B:280:LYS:HA	1.65	0.47
1:B:224:LEU:HD12	1:B:229:GLN:HB3	1.96	0.47
1:A:87:ASP:OD2	1:A:87:ASP:N	2.47	0.47
1:A:285:SER:HB3	1:A:300:TRP:CE3	2.50	0.47
1:B:222:LYS:HG2	4:B:390:HOH:O	2.15	0.47
2:D:100:SER:OG	2:D:102:VAL:HG22	2.15	0.46
2:D:11:TRP:CE3	2:D:164:GLY:HA3	2.51	0.46
1:B:289:GLN:HG2	1:B:290:ASP:N	2.32	0.45
1:B:138:ARG:NH1	1:B:184:LEU:O	2.49	0.44
2:C:78:LEU:HD11	2:C:112:LEU:HD13	1.98	0.44
1:B:94:ILE:HG23	1:B:130:LEU:HD11	2.00	0.44
1:B:278:CYS:SG	1:B:284:ILE:HD11	2.57	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:GLN:O	1:B:99:LYS:CB	2.64	0.44
1:B:229:GLN:HE21	1:B:277:ILE:CD1	2.30	0.44
2:D:8:SER:HB2	2:D:9:PRO:CD	2.47	0.44
1:A:37:THR:HB	1:A:46:VAL:HG13	2.00	0.43
1:B:86:GLU:O	1:B:89:ILE:HG22	2.19	0.43
1:A:142:ASP:HA	1:A:143:PRO:HD2	1.84	0.43
1:B:57:VAL:HG11	1:B:82:LEU:HD21	1.99	0.43
2:D:1:ARG:HH11	2:D:1:ARG:HB3	1.84	0.43
2:D:78:LEU:HD11	2:D:112:LEU:HD13	2.00	0.43
1:A:41:ASP:HB3	1:A:42:GLN:H	1.56	0.43
1:A:64:GLY:HA2	1:A:196:LEU:HD11	2.01	0.43
2:C:64:ARG:HH11	2:C:64:ARG:HB3	1.84	0.43
1:A:97:ASP:OD2	1:A:99:LYS:N	2.51	0.42
2:D:8:SER:CB	2:D:9:PRO:CD	2.97	0.42
1:A:100:GLU:CB	1:A:101:PRO:CD	2.96	0.42
1:A:270:ASP:CG	1:A:291:ARG:HD2	2.44	0.42
1:B:141:SER:HB3	4:B:362:HOH:O	2.19	0.42
1:B:16:TYR:CD1	1:B:16:TYR:N	2.87	0.42
1:A:122:TRP:NE1	1:A:167:TYR:HB2	2.35	0.42
1:A:276:VAL:HG21	1:A:284:ILE:HD12	2.02	0.42
1:B:96:LYS:O	1:B:97:ASP:C	2.61	0.42
2:D:13:GLN:HG3	4:D:182:HOH:O	2.19	0.41
1:B:246:TYR:CG	2:D:60:PRO:HG3	2.55	0.41
2:C:90:PHE:CE1	2:C:143:ARG:HG2	2.54	0.41
1:B:237:PRO:HG2	1:B:240:TRP:HB2	2.02	0.41
1:B:281:ASN:HD22	1:B:281:ASN:HA	1.56	0.41
2:C:0:LEU:N	2:C:0:LEU:CD1	2.84	0.41
1:A:95:LEU:HD21	1:A:122:TRP:CD1	2.57	0.40
1:B:70:LYS:HB2	1:B:75:LEU:HD11	2.03	0.40
1:B:16:TYR:CD2	1:B:18:ASP:O	2.74	0.40
1:B:121:TRP:HA	1:B:167:TYR:O	2.21	0.40
2:C:163:HIS:CE1	2:C:167:THR:HG21	2.57	0.40
1:B:287:ARG:HB2	1:B:300:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/314 (94%)	276 (93%)	18 (6%)	2 (1%)	18	23
1	B	286/314 (91%)	273 (96%)	13 (4%)	0	100	100
2	C	129/177 (73%)	124 (96%)	5 (4%)	0	100	100
2	D	129/177 (73%)	118 (92%)	10 (8%)	1 (1%)	16	20
All	All	840/982 (86%)	791 (94%)	46 (6%)	3 (0%)	30	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	98	PRO
1	A	141	SER
1	A	101	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/285 (92%)	250 (95%)	12 (5%)	24	36
1	B	259/285 (91%)	243 (94%)	16 (6%)	16	24
2	C	109/151 (72%)	102 (94%)	7 (6%)	16	23
2	D	114/151 (76%)	103 (90%)	11 (10%)	8	10
All	All	744/872 (85%)	698 (94%)	46 (6%)	16	24

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	GLU
1	A	86	GLU
1	A	101	PRO
1	A	121	TRP
1	A	127	SER
1	A	135	LYS
1	A	226	ASN
1	A	228	ARG
1	A	267	VAL
1	A	291	ARG
1	A	295	SER
1	A	306	SER
1	B	0	GLU
1	B	16	TYR
1	B	32	GLU
1	B	41	ASP
1	B	59	GLU
1	B	84	LYS
1	B	87	ASP
1	B	97	ASP
1	B	105	THR
1	B	153	LEU
1	B	202	THR
1	B	256	GLN
1	B	264	LYS
1	B	278	CYS
1	B	281	ASN
1	B	291	ARG
2	C	0	LEU
2	C	64	ARG
2	C	82	GLU
2	C	105	LEU
2	C	112	LEU
2	C	125	THR
2	C	138	GLN
2	D	0	LEU
2	D	3	VAL
2	D	15	GLN
2	D	73	ARG
2	D	82	GLU
2	D	84	LEU
2	D	97	LEU
2	D	112	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	119	GLU
2	D	141	LEU
2	D	150	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	65	GLN
1	A	226	ASN
1	B	144	GLN
1	B	172	GLN
1	B	220	GLN
1	B	229	GLN
1	B	281	ASN
2	C	104	GLN
2	D	15	GLN
2	D	121	HIS
2	D	172	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	313	1	14,14,15	0.61	0	17,19,21	1.49	4 (23%)
3	NAG	A	313	1	14,14,15	0.55	0	17,19,21	1.77	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	313	1	-	4/6/23/26	0/1/1/1
3	NAG	A	313	1	1/1/5/7	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	313	NAG	C1-O5-C5	4.97	118.85	112.19
3	A	313	NAG	C2-N2-C7	3.54	127.64	122.90
3	B	313	NAG	C1-O5-C5	3.49	116.86	112.19
3	B	313	NAG	O5-C1-C2	3.04	115.99	111.29
3	B	313	NAG	C2-N2-C7	2.29	125.97	122.90
3	B	313	NAG	C3-C4-C5	2.11	114.06	110.23
3	A	313	NAG	O7-C7-N2	2.04	125.58	121.98

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	313	NAG	C1

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	313	NAG	C8-C7-N2-C2
3	B	313	NAG	O7-C7-N2-C2
3	B	313	NAG	O5-C5-C6-O6
3	B	313	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	313	NAG	C4-C5-C6-O6
3	A	313	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/314 (95%)	0.33	14 (4%) 36 38	22, 42, 67, 80	3 (1%)
1	B	294/314 (93%)	0.39	18 (6%) 27 29	24, 44, 67, 76	2 (0%)
2	C	137/177 (77%)	0.36	7 (5%) 33 35	22, 41, 70, 83	0
2	D	137/177 (77%)	0.81	16 (11%) 9 10	31, 54, 76, 92	0
All	All	868/982 (88%)	0.43	55 (6%) 26 28	22, 45, 70, 92	5 (0%)

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	137	TRP	4.9
2	D	0	LEU	4.1
2	D	140	LEU	4.0
2	D	6	GLY	3.9
2	C	48	ASP	3.7
1	B	102	LYS	3.6
1	B	103	ASN	3.5
2	D	141	LEU	3.4
2	C	169	SER	3.3
1	B	13	LEU	3.0
2	C	28	ALA	3.0
1	B	18	ASP	3.0
1	A	100	GLU	3.0
2	C	137	TRP	3.0
1	A	103	ASN	2.9
1	B	98	GLN	2.9
1	B	200	ASN	2.8
2	C	29	HIS	2.8
1	B	100	GLU	2.8
1	A	306	SER	2.7
2	C	0	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	47	ASN	2.7
1	B	303	VAL	2.7
1	B	305	CYS	2.6
1	B	144	GLN	2.6
1	A	18	ASP	2.6
2	D	97	LEU	2.6
2	D	7	SER	2.6
1	B	229	GLN	2.4
1	A	19	ALA	2.4
1	A	101	PRO	2.4
2	D	171	HIS	2.4
2	D	144	PHE	2.3
1	A	102	LYS	2.3
1	B	223	PRO	2.3
1	A	21	GLY	2.3
1	A	121	TRP	2.3
2	D	173	HIS	2.3
1	B	255	VAL	2.3
1	A	99	LYS	2.3
2	D	98	PRO	2.2
2	D	5	GLY	2.2
1	B	41	ASP	2.2
1	B	279	ARG	2.2
2	D	118	PRO	2.2
1	B	265	ASP	2.1
2	D	8	SER	2.1
2	D	49	VAL	2.1
1	A	88	GLY	2.1
1	B	224	LEU	2.0
2	D	1	ARG	2.0
1	B	99	LYS	2.0
1	A	41	ASP	2.0
1	A	20	PRO	2.0
1	A	263	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	313	14/15	0.82	0.13	54,57,60,61	0
3	NAG	A	313	14/15	0.89	0.10	54,55,56,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.